

sulaimanIJA1-4-2012-2.pdf

Subgroups Lattice of Symmetric Group S_4

R. Sulaiman

Department of Mathematics, Faculty of Mathematics and Sciences
Universitas Negeri Surabaya, Surabaya 60231 Indonesia
sulaimanraden@yahoo.com

Abstract

In this paper, we determine all of subgroups of symmetric group S_4 by applying Lagrange theorem and Sylow theorem. First, we observe the multiplication table of S_4 , then we determine all possibilities of every subgroup of order n , with n is the factor of order S_4 . We found 30 subgroups of S_4 . The diagram of lattice subgroups of S_4 is then presented.

Mathematics Subject Classification: 20B30, 20B35, 03G10

Keywords: Lagrange theorem, Sylow theorem, Sylow p -subgroup, symmetric group, lattice

1 Introduction

For an arbitrary nonempty set S , define $A(S)$ to be the set of all one-to-one mapping of the set S onto itself. The set $A(S)$ with composition function operation is a group. If the set S contains n elements, then group $A(S)$ are denoted by S_n . Group S_n has $n!$ elements and will be called the symmetric group. There are many references on subgroups of S_2 and S_3 ([2],[4] and [5]). In this paper, we determine all subgroups of S_4 and then draw diagram of lattice subgroups of S_4 .

One of the most important problem of fuzzy group theory is to classify the fuzzy subgroup of a finite group. Many papers have treated the particular case of finite cyclic group. Laszlo [3] studied about the construction of fuzzy subgroup of group of order one to six. Zhang and Zou [13] have determined the number of fuzzy subgroup of cyclic groups of order p^n where p is a prime number. Murali and Makamba ([11],[12]) considered a similar problem. They have determined the number of fuzzy subgroups of $Z_{p^n}, Z_p \times Z_p, Z_{p^n} \times Z_q$ and abelian groups of order $p^n q^m$ where p and q are different primes. Tarnauceanu

and Bentea [6] have determined this number for finite abelian groups. Sulaiman and Abd Ghafur [8] have counted the number of fuzzy subgroups of finite cyclic groups. All of them have treated the particular case of finite abelian group. It is interesting to investigate the number of fuzzy subgroups of nonabelian groups. Sulaiman and Abd Ghafur [7] have counted the number of fuzzy subgroups of nonabelian symmetric groups S_2, S_3 and alternating group A_4 . They [9] have counted for group defined by a presentation, too.

There is a very close relationship between the number of chain of lattice subgroup of group G and the number of fuzzy subgroup of G . Therefore, the result of this paper, that is a diagram of lattice subgroups of S_4 is very important to determine the number of fuzzy subgroup of S_4 .

2 Preliminary

We recall some definitions and results that will be used later.

Definition 2.1 A partial ordered on a nonempty set P is a binary relation $\leq$ on P that is reflexive, antisymmetric and transitive. The pair $\langle P, \leq \rangle$ is called a partially ordered set or poset. Poset $\langle P, \leq \rangle$ is totally ordered if $\forall x, y \in P$ are comparable, that is $x \leq y$ or $y \leq x$. A nonempty subset S of P is a chain in P if S is totally ordered by $\leq$. (Roman, [10])

Definition 2.2 Let $\langle P, \leq \rangle$ be a poset and let $S \subseteq P$. An upper bound for S is an element $x \in P$ for which $s \leq x \forall s \in S$. The least upper bound of S is called the supremum or join of S . A lower bound for S is an element $x \in P$ for which $x \leq s \forall s \in S$. The greatest lower bound of S is called the infimum or meet of S . Poset $\langle P, \leq \rangle$ is called a lattice if every pair x, y elements of P has a supremum and an infimum (Roman, [10]).

Note that the set of all of subgroups G under the "subgroup" relation is a lattice. This lattice is called the lattice subgroup of G .

Theorem 2.3 (Lagrange's theorem) If G is a finite group and H is a subgroup of G , then order of H is a divisor of order G (Herstein, [4]).

Theorem 2.4 If G is a finite group and $a \in G$, then order of a is a divisor of order G (Herstein, [4]).

Theorem 2.5 (The First Sylow Theorem) Let G be a finite group and let $|G| = p^n m$ where $n \geq 1$, p is a prime number and $(p, m) = 1$. Then G contains a subgroup of order p^i for each i where $1 \leq i \leq n$ (Fraleigh, [5]).

2

Definition 2.6 Let G be a finite group and let $|G| = p^n m$ where $n \geq 1$, p is a prime number and $(p, m) = 1$. The subgroup of G of order p^n is called the Sylow p -subgroup of G (Gardiner, [2]).

2

Theorem 2.7 (The Third Sylow Theorem) Let G be a finite group and let $|G| = p^n m$ where $n \geq 1$, p is a prime number and $(p, m) = 1$. Then the number of Sylow p -subgroup is of the form $(1 + kp)$, where k is a non-negative integer, and $(1 + kp)$ divides the order of G (Fraleigh, [5]).

3

Definition 2.8 A subgroup N of G is said to be a normal subgroup of G if for every $g \in G$ and $n \in N$, $gng^{-1} \in N$ (Herstein, [4]).

17

Theorem 2.9 There is a unique Sylow p -subgroup of the finite group G if only if it is normal (Gardiner, [2]).

11

Theorem 2.10 Let G be a group of order pq , where p and q distinct primes and $p < q$. Then G has only one subgroup of order q . This subgroup of order q is normal in G (Gardiner, [2]).

3 Main Results

Let $S_4 = \{i = id, \sigma_1, \sigma_2, \dots, \sigma_9, \tau_1, \tau_2, \dots, \tau_8, \alpha_1, \alpha_2, \dots, \alpha_6\}$ with $\sigma_1 = (1234), \sigma_2 = (13)(24), \sigma_3 = (1432), \sigma_4 = (1243), \sigma_5 = (14)(23), \sigma_6 = (1342), \sigma_7 = (1324), \sigma_8 = (12)(34), \sigma_9 = (1432), \tau_1 = (234), \tau_2 = (243), \tau_3 = (134), \tau_4 = (143), \tau_5 = (124), \tau_6 = (142), \tau_7 = (123), \tau_8 = (132), \alpha_1 = (12), \alpha_2 = (13), \alpha_3 = (14), \alpha_4 = (23), \alpha_5 = (24), \alpha_6 = (34)$. Multiplication table of S_4 can be seen in [1]. The order of elements of S_4 are shown as follows.

Order	Elements
1	i
9	$\sigma_2, \sigma_5, \sigma_8, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6$
3	$\tau_1, \tau_2, \tau_3, \tau_4, \tau_5, \tau_6, \tau_7, \tau_8$
4	$\sigma_1, \sigma_3, \sigma_4, \sigma_6, \sigma_7, \sigma_9$

According to the Lagrange theorem, nontrivial subgroups of S_4 must have order 2, 4, 6, 8 or 12. We will determine all of the subgroups of S_4 . Clearly, the subgroup of S_4 of order 1 is the trivial subgroup $H_1 = \{i\}$.

22

Subgroups of order 2:

Let K be arbitrary subgroup of S_4 of order 2. Since 2 is a prime number, then K is cyclic. Therefore K is generated by an element of S_4 of order 2. Thus, all of subgroups of S_4 of order 2 are $K_2 = \{i, \sigma_2\}$, $K_3 = \{i, \sigma_5\}$, $K_4 = \{i, \sigma_8\}$, $K_5 = \{i, \alpha_1\}$, $K_6 = \{i, \alpha_2\}$, $K_7 = \{i, \alpha_3\}$, $K_8 = \{i, \alpha_4\}$, $K_9 = \{i, \alpha_5\}$, $K_{10} =$

$\{i, \alpha_6\}$.

Subgroups of order 3:

Similar to order 2, subgroups of S_4 of order 3 is generated by a 21 element of S_4 of order 3. Thus, all of subgroups of S_4 of order 3 are $L_{11} = \langle \tau_1 \rangle = \langle \tau_2 \rangle = \{i, \tau_1, \tau_2\}$, $L_{12} = \langle \tau_3 \rangle = \langle \tau_4 \rangle = \{i, \tau_3, \tau_4\}$, $L_{13} = \langle \tau_5 \rangle = \langle \tau_6 \rangle = \{i, \tau_5, \tau_6\}$, $L_{14} = \langle \tau_7 \rangle = \langle \tau_8 \rangle = \{i, \tau_7, \tau_8\}$.

Subgroups of order 4:

Let M be any arbitrary subgroup of S_4 of order 4. According to the 10 theorem 2.4, then elements of M must have order 1, 2 or 4. If M consists an element of order 4, then M is generated by an element of order 4. The subgroups are $M_{15} = \langle \sigma_1 \rangle = \langle \sigma_2 \rangle = \langle \sigma_3 \rangle = \{i, \sigma_1, \sigma_2, \sigma_3\}$, $M_{16} = \langle \sigma_4 \rangle = \langle \sigma_5 \rangle = \langle \sigma_6 \rangle = \{i, \sigma_4, \sigma_5, \sigma_6\}$, $M_{17} = \langle \sigma_7 \rangle = \langle \sigma_8 \rangle = \langle \sigma_9 \rangle = \{i, \sigma_7, \sigma_8, \sigma_9\}$. If M consists an element of order 2, then M does not have element of order 4. Therefore the order of all of the elements of M is 2, except identity element. We will try to multiply all of the combination of elements of order 2. The following is the multiplication table of the elements of order 2.

	σ_2	σ_5	σ_8	α_1	α_2	α_3	α_4	α_5	α_6
σ_2	i	σ_8	σ_5	σ_7	α_5	σ_6	σ_4	α_2	σ_9
σ_5	σ_8	i	σ_2	σ_9	σ_3	α_4	α_3	σ_1	σ_7
σ_8	σ_5	σ_2	i	α_6	σ_1	σ_4	σ_6	σ_3	α_1
α_1	σ_9	σ_7	α_6	i	τ_7	τ_5	τ_8	τ_6	σ_8
α_2	α_5	σ_1	σ_3	τ_8	i	τ_3	τ_7	σ_2	τ_4
α_3	σ_4	α_4	σ_6	τ_6	τ_4	i	σ_5	τ_5	τ_3
α_4	σ_2	α_3	σ_4	τ_7	τ_8	σ_5	i	τ_1	τ_2
α_5	α_2	σ_3	σ_1	τ_5	σ_2	τ_6	τ_2	i	τ_1
α_6	σ_7	σ_9	α_1	σ_8	τ_3	τ_4	τ_1	τ_2	i

We observe the table above by row, from the second row. We conclude that $\{\sigma_2, \alpha_1\}$, $\{\sigma_2, \alpha_3\}$, $\{\sigma_2, \alpha_4\}$, $\{\sigma_2, \alpha_6\}$ are not subsets of M , because the order of the result of their product is not 2. We get the subgroups of order 4 containing σ_2 are $M_{18} = \{i, \sigma_2, \sigma_5, \sigma_8\}$ and $M_{19} = \{i, \sigma_2, \alpha_2, \alpha_5\}$. Using similar argument to the other rows, we obtain the other subgroups of order 4, those are $M_{20} = \{i, \sigma_5, \alpha_3, \alpha_4\}$ and $M_{21} = \{i, \sigma_8, \alpha_1, \alpha_6\}$. Thus, we got seven subgroups of order 4.

Subgroups of order 6:

Let N be any arbitrary subgroup of S_4 of order 6. Since $|N| = 2 \times 3$, according to the Theorem 2.7 (The Third Sylow Theorem) then N has only one subgroup M of order 3. According to the Theorem 2.9, this subgroup M is normal in N . There are four possibilities subgroups of order 3, those are L_{11}, L_{12}, L_{13} or L_{14} .

Using normality M in N , we got four subgroups of order 6, those are $N_{22} = \{i, \tau_1, \tau_2, \alpha_4, \alpha_5, \alpha_6\}$, $N_{23} = \{i, \tau_3, \tau_4, \alpha_2, \alpha_3, \alpha_6\}$, $N_{24} = \{i, \tau_5, \tau_6, \alpha_1, \alpha_3, \alpha_5\}$ and $N_{25} = \{i, \tau_7, \tau_8, \alpha_1, \alpha_2, \alpha_4\}$.

Subgroups of order 8:

Let P be arbitrary subgroup of S_4 of order 8. Since there is no element of S_4 of order 8, then every element of P must have order 1, 2 or 4. Also, since $|S_4| = 2^3 \cdot 3$, then by Theorem 2.7, the number of subgroups of S_4 of order 8 is 1 or 3. We will try to construct all of the possibilities and we will stop if we get three subgroups. We start from the table of orbit of element of order 4.

Element	Orb
σ_1, σ_3	$\{\sigma_1, \sigma_2, \sigma_3, i\}$
σ_4, σ_6	$\{\sigma_4, \sigma_5, \sigma_6, i\}$
σ_7, σ_9	$\{\sigma_7, \sigma_8, \sigma_9, i\}$

Considering the orbit, we can conclude that P have only three possibilities:

- i) Consists two elements of order 4
- ii) Consists four elements of order 4
- iii) No element of order 4.

First, we will determine subgroup of order 8 containing two elements of order 4. The following is a multiplication table between elements of order 4 (which have same orbit) and elements of order 2.

	σ_2	σ_5	σ_8	α_1	α_2	α_3	α_4	α_5	α_6
σ_1	σ_3	α_2	α_5	τ_1	σ_8	τ_7	τ_3	σ_5	τ_5
σ_3	σ_1	α_5	α_2	τ_4	σ_5	τ_2	τ_6	σ_8	τ_8

Considering the table above, we conclude that $\alpha_1, \alpha_3, \alpha_4$ or α_6 can not be added to $\{i, \sigma_1, \sigma_1\sigma_3\}$ in order to be a subgroup of order 8. We get a subgroup of order 6 if we add the other elements of order 2 namely, $P_{26} = \{i, \sigma_1, \sigma_2, \sigma_3, \sigma_5, \sigma_8, \alpha_2, \alpha_5\}$. Using similar argument, we get the other subgroups of S_4 of order 8, those are $P_{27} = \{i, \sigma_2, \sigma_4, \sigma_5, \sigma_6, \sigma_8, \alpha_3, \alpha_4\}$ and $P_{28} = \{i, \sigma_2, \sigma_5, \sigma_7, \sigma_8, \sigma_9, \alpha_1, \alpha_6\}$.

Subgroups of order 12:

Obviously the alternating group $A_4 = \{i, \sigma_2, \sigma_5, \tau_1, \tau_2, \tau_3, \tau_4, \tau_5, \tau_6, \tau_7, \tau_8\}$ is a subgroup of S_4 of order 12. We will prove that A_4 is the unique subgroup of S_4 of order 12. Let $H \leq S_4$ where $|H| = 12$ and $H \neq A_4$. We conclude that (from Theorem 2.7), there are three possibilities:

- i) The number of subgroups H of order 4 is 3 and of order 3 is 4
- ii) The number of subgroups H of order 4 is 1 and of order 3 is 1

- iii) ⁸ The number of subgroups H of order 4 is 3 and of order 3 is 1.

The case i) cannot occur, because three subgroups of order 3 together with ²⁸ subgroups of order 4 consist more than 12 elements.

Case ii): Let the subgroup of order 3 is K and let the subgroup of order 4 is L . According to the Theorem 5, then K and L are normal in H . We can show that $H = KL$ and $kl = lk, \forall k \in K, l \in L$. Hence, H must be abelian. Checking all of possibilities of K and L , we get a conclusion that this case cannot occur.

Case iii): Let $\mathcal{A}$ is a collection of the three subgroups of order 4 and K is the subgroup of order 3. According to the Theorem 5, then K is normal in H . Therefore $h^{-1}kh \in K, \forall h \in H, k \in K$. Giving counter examples for every possibility of K we conclude this condition is not held for every subgroup K of order 3. This means that this case cannot occur. Therefore we got only one subgroup of order 12 namely alternating group, A_4 .

According to this result, we have the diagram of lattice subgroups of S_4 is shown as figure.1 below.




Figure 1: Subgroup lattice of S_4 .

References

- [1] B. Baumslag, *Theory and Problems of Group Theory :Schaum's outline Series*, Mc Graw-Hill, New York, 1968.
- [2] C.F. Gardiner, *A First Course in Group Theory*, Springer-Verlag, Berlin, 1997.
- [3] F. Laszlo, Structure and construction of fuzzy subgroup of a group, *Fuzzy Set and System.*, **51** (1992), 105 - 109.
- [4] I.N. Herstein, *Topic in Algebra*, John Wiley and Sons, New York, 1975.
- [5] J.B. Fraleigh, *A First Course in Abstract Algebra*, Addison-Wesley, London, 1992.
- [6] M. Tarnauceanu and L. Bentea, On the number of fuzzy subgroups of finite abelian groups, *Fuzzy Set and System.*, **159** (2008), 1084 - 1096.
- [7] R. Sulaiman and Abd Ghafur Ahmad, Counting fuzzy subgroups of symmetric groups S_2, S_3 and alternating group A_4 , *Journal of Quality Measurement and Analysis.*, **6** (2010), 57 - 63.
- [8] R. Sulaiman and Abd Ghafur Ahmad, The number of fuzzy subgroups of finite cyclic groups, *International Mathematical Forum.*, **6 no.20** (2011), 987 - 994.
- [9] R. Sulaiman and Abd Ghafur Ahmad, The number of fuzzy subgroups of group defined by apresentation, *International Journal of Algebra.*, **5 no.8** (2011), 375 - 382.
- [10] S. Roman, *Lattice and Ordered Set*, Springer, New York, 2008.
- [11] V. Murali and B.B. Makamba, On an equivalence of fuzzy subgroups I, *Fuzzy Set and System.*, **123** (2001), 259 - 264.
- [12] V. Murali and B.B. Makamba, Counting the number of fuzzy subgroups of an abelian group of order $p^n q^m$, *Fuzzy Set and System.*, **144** (2004), 459 - 470.
- [13] Y. Zhang and K. Zou, A note on an equivalence relation on fuzzy subgroups, *Fuzzy Set and System.*, **95** (1998), 243 - 247.

Received: August, 2011

22%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|---|---------------|
| 1 | B. Humera, Z. Raza. "On fuzzy subgroups of finite Abelian groups", International Mathematical Forum, 2013
<small>Crossref</small> | 90 words — 4% |
| 2 | Shen, Rulin, Xuan Zou, and Wujie Shi. "A characterization of $\mathbb{Z}_5$ by same-order type", Monatshefte für Mathematik, 2015.
<small>Crossref</small> | 54 words — 2% |
| 3 | Studies in Fuzziness and Soft Computing, 2015.
<small>Crossref</small> | 30 words — 1% |
| 4 | nigam.info
<small>Internet</small> | 28 words — 1% |
| 5 | www.pphmj.com
<small>Internet</small> | 24 words — 1% |
| 6 | John Mordeson. "Equivalence of Fuzzy Subgroups of Finite Abelian Groups", Studies in Fuzziness and Soft Computing, 2005
<small>Crossref</small> | 24 words — 1% |
| 7 | www.win.tue.nl
<small>Internet</small> | 20 words — 1% |
| 8 | Tarnauceanu, M.. "On the number of fuzzy subgroups of finite abelian groups", Fuzzy Sets and Systems, 20080501
<small>Crossref</small> | 18 words — 1% |
| 9 | www.gecafs.org | |

17 words — 1%

10 T. G. Ostrom. "Elementary abelian 2-groups on the line at infinity of translation planes", Journal of Geometry, 12/1981
Crossref 14 words — 1%

11 Griggs, J.R.. "Spanning subset sums for finite Abelian groups", Discrete Mathematics, 20010228
Crossref 14 words — 1%

12 www.imo2001.org
Internet 13 words — 1%

13 A. Matta. "Tool management in flexible manufacturing systems with network part program", International Journal of Production Research, 9/1/2004
Crossref 12 words — 1%

14 "Mathematics Is Not a Spectator Sport", Springer Nature, 2005
Crossref 12 words — 1%

15 dspace.cc.tut.fi
Internet 11 words — < 1%

16 Churchill, R.C.. "Two Generator Subgroups of $SL(2, \mathbb{C})$ and the Hypergeometric, Riemann, and Lamé Equations", Journal of Symbolic Computation, 199910
Crossref 10 words — < 1%

17 V. A. Artamonov. "Integral group rings: Groups of units and classical K-theory", Journal of Soviet Mathematics, 11/1991
Crossref 10 words — < 1%

18 Sergio Rajsbaum. "ACM SIGACT News distributed computing column 12", ACM SIGACT News, 10 words — < 1%

9/1/2003

Crossref

19 Schneider. "Spatial data types — A survey",
Lecture Notes in Computer Science, 1997 10 words — < 1%

Crossref

20 Kim, K.H.. "A layered workflow knowledge Grid/P2P
architecture and its models for future generation
workflow systems", Future Generation Computer Systems,
200703 9 words — < 1%

Crossref

21 Lecture Notes in Computer Science, 2009. 8 words — < 1%

Crossref

22 repository.essex.ac.uk 8 words — < 1%

Internet

23 Zhen Cai Shen. "Finite groups with some pronormal
subgroups", Acta Mathematica Sinica English
Series, 04/2011 8 words — < 1%

Crossref

24 Marinos Kavouras. "A method for the formalization
and integration of geographical categorizations",
International Journal of Geographical Information Science,
7/1/2002 8 words — < 1%

Crossref

25 Petros Maragos. "Lattice Image Processing: A
Unification of Morphological and Fuzzy Algebraic
Systems", Journal of Mathematical Imaging and Vision, 05/2005 8 words — < 1%

Crossref

26 XUAN-HA VU. "A UNIFYING FRAMEWORK FOR
GENERALIZED CONSTRAINT ACQUISITION",
International Journal on Artificial Intelligence Tools, 2008 7 words — < 1%

Crossref

27 Ordered Sets, 2016. 7 words — < 1%

Crossref

28	Warren J. Wong. "Determination of a class of primitive permutation groups", Mathematische Zeitschrift, 1967 Crossref	6 words — < 1%
29	Journal of Enterprise Information Management, Volume 29, Issue 3 (2016) Publications	6 words — < 1%
30	Lujin Zhu. "WEAKLY c-NORMAL SUBGROUPS OF FINITE GROUPS AND THEIR PROPERTIES", Communications in Algebra, 1/1/2003 Crossref	6 words — < 1%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF