

Computing the number of fuzzy subgroups by expansion method

By raden sulaiman

COMPUTING THE NUMBER OF FUZZY SUBGROUPS BY EXPANSION METHOD

Raden Sulaiman^{1 §}, Badi Priyo Prawoto²

^{1,2}Department of Mathematics
Faculty of Mathematics and Sciences
Universitas Negeri Surabaya
Surabaya, 60231, INDONESIA

Abstract: The purpose of this paper is to construct the method to find the number of fuzzy subgroups for rectangle groups. The method is formulated by using the pattern of their lattice diagram. We construct the expansion methods, row and column expansion.

AMS Subject Classification: 06B05, 20N25, 01-08

Key Words: fuzzy subgroup, rectangle group, expansion method

1. Introduction

In 1965, Zadeh [1] introduced the concept of fuzzy set. Rosenfeld in 1971 [2] was started study of fuzzy algebraic structures. He introduced the concept of fuzzy subgroups. One the most important problem of fuzzy theory is to classify the fuzzy subgroups of finite group [3]. There are many papers have written. Lazlo (see [4]) constructed of fuzzy subgroups of groups of order one to six. Murali and Makamba (see [5], [6]) found the number of fuzzy subgroups of abelian groups of order $3^n q^n$ where p and q are different primes. Sulaiman (see [7] and [8]) constructed the fuzzy subgroups of symmetric group S_4 and alternating group A_4 .

The definition of equivalence relation on fuzzy subgroups introduced in [9]. Some

Received: June 30, 2014

© 2014 Academic Publications, Ltd.

[§]Correspondence author

definitions and theorems in [9] is used in this paper. Sulaiman [9] developed a method to count the number of fuzzy subgroups of finite group. The method use the subgroup lattice of group. This method can be used not only for abelian group, but also for non-abelian as well. By using the method, we [10] constructed some formulas to count the number of fuzzy subgroups of "rectangle groups". In this paper, we develop the results to find some formulas. The formulas are constructed by expansion methods, row and column expansion.

2. Preliminaries

In this paper, a group G is assumed to be a finite group. First of all, we present some basic notions and results that will be used later. Definition of equivalence relation on fuzzy subgroups use The Definition 6 in [9].

Definition 1. Let X be a nonempty set. A fuzzy subset of X is a function from X into $[0, 1]$.

Definition 2. (Rosenfeld, see [2]). Let G be a group. A fuzzy subset μ of G is called a fuzzy subgroup of G if:

- (1) $\mu(xy) \geq \min\{\mu(x), \mu(y)\}, \forall x, y \in G,$
- (2) $\mu(x^{-1}) \geq \mu(x), \forall x \in G.$

Theorem 3. (Sulaiman, see Theorem 11 in [9]) Let H be a subgroup of G , and let the set of all subgroups of G which contain H (but are not equal to H) be $\{H_1, H_2, H_3, \dots, H_k\}$. Then $n(F_{P_1=H}) = \sum_{i=1}^k n(F_{P_1=H_i})$.

Definition 4. (Sulaiman and Budi Priyo, see [10]). Let G be a group and the number of its subgroups is finite. The diagram of lattice subgroups of G is called "rectangle" and group G is called "rectangle group" if satisfies these conditions: The subgroups of G can be labeled by K_j^i where $1 \leq i \leq m, 1 \leq j \leq s$ for some $m, s \in \mathbb{N}$ with $K_1^1 = G, K_s^m = \{e\}$ such that: (i) for fixed $i, 1 \leq i \leq m, K_k^i < K_{k-1}^i, \forall k, 2 \leq k \leq s$, (ii) for fixed $j, 1 \leq j \leq m, K_j^t < K_j^{t-1}, \forall t, 2 \leq t \leq m$. In this case the size of the diagram is $m \times s$. The diagram of "rectangle lattice" is shown in Figure 1.

3. Expansion Method

In this section we find the number of fuzzy subgroups of "rectangle group" by row and column expansion.




Figure 1: Rectangle Lattice

Theorem 5. Let G be a rectangle group with $m, s \in N$. We have

$$n(F_{P_1=K_s^m=\{e\}}) = 2 \cdot n(F_{P_1=K_{s-1}^m}) + \sum_{i=1}^{m-1} n(F_{P_1=K_s^i}).$$

Proof. Let $m, s \in N$. By analyzing Figure 1 and applying Theorem 3, we have,

$$\begin{aligned} n(F_{P_1=K_s^m=\{e\}}) &= \sum_{i=1}^{s-1} n(F_{P_1=K_i^m}) + \sum_{i=1}^s n(F_{P_1=K_i^{m-1}}) + \sum_{i=1}^s n(F_{P_1=K_i^{m-2}}) \\ &\quad + \dots + \sum_{i=1}^s n(F_{P_1=K_i^2}) + \sum_{i=1}^s n(F_{P_1=K_i^1}). \end{aligned}$$

We can write

$$\begin{aligned} n(F_{P_1=K_s^m=\{e\}}) &= \sum_{i=1}^{s-1} [n(F_{P_1=K_i^m}) + n(F_{P_1=K_i^{m-1}}) + n(F_{P_1=K_i^{m-2}}) + \dots + n(F_{P_1=K_i^2}) \\ &\quad + n(F_{P_1=K_i^1})] + n(F_{P_1=K_s^{m-1}}) + n(F_{P_1=K_s^{m-2}}) + \dots + n(F_{P_1=K_s^2}) + n(F_{P_1=K_s^1}). \end{aligned} \quad (1)$$

Analogue with this way we have

$$n(F_{P_1=K_{s-1}^m}) = \sum_{i=1}^{s-1} [n(F_{P_1=K_i^m}) + n(F_{P_1=K_i^{m-1}}) + n(F_{P_1=K_i^{m-2}}) + \dots + n(F_{P_1=K_i^2}) + n(F_{P_1=K_i^1})] + n(F_{P_1=K_{s-1}^{m-1}}) + n(F_{P_1=K_{s-1}^{m-2}}) + \dots + n(F_{P_1=K_{s-1}^2}) + n(F_{P_1=K_{s-1}^1}). \quad (2)$$

Eq. (1) can be written as

$$n(F_{P_1=K_s^m}) = \sum_{i=1}^{s-2} [n(F_{P_1=K_i^m}) + n(F_{P_1=K_i^{m-1}}) + n(F_{P_1=K_i^{m-2}}) + \dots + n(F_{P_1=K_i^2}) + n(F_{P_1=K_i^1})] + n(F_{P_1=K_{s-1}^m}) + n(F_{P_1=K_{s-1}^{m-1}}) + n(F_{P_1=K_{s-1}^{m-2}}) + \dots + n(F_{P_1=K_{s-1}^2}) + n(F_{P_1=K_{s-1}^1}) + n(F_{P_1=K_s^2}) + n(F_{P_1=K_s^1}). \quad (3)$$

Substitute (2) to (3) to get

$$n(F_{P_1=K_s^m}) = 2.n(F_{P_1=K_{s-1}^m}) + n(F_{P_1=K_{s-1}^{m-1}}) + n(F_{P_1=K_{s-1}^{m-2}}) + \dots + n(F_{P_1=K_{s-1}^2}) + n(F_{P_1=K_{s-1}^1}).$$

Finally we have

$$n(F_{P_1=K_s^m}) = 2.n(F_{P_1=K_{s-1}^m}) + \sum_{i=1}^{m-1} [n(F_{P_1=K_s^i})]. \quad \square$$

Theorem 6. (Raw Expansion) Let G be a rectangle group with $m, s \in N$ and $s \geq 3$. We have

$$n(F_{P_1=K_s^m}) = 2.n(F_{P_1=K_{s-1}^m}) + 2.n(F_{P_1=K_{s-1}^{m-1}}) + 2^2.n(F_{P_1=K_{s-1}^{m-2}}) + 2^3.n(F_{P_1=K_{s-1}^{m-3}}) + \dots + 2^{m-3}.n(F_{P_1=K_{s-1}^3}) + 2^{m-2}.n(F_{P_1=K_{s-1}^2}) + 2^{m-1}.n(F_{P_1=K_{s-1}^1}).$$

Proof. According to the Theorem 5 we have:

$$n(F_{P_1=K_s^2}) = 2.n(F_{P_1=K_{s-1}^2}) + n(F_{P_1=K_s^1}), \quad (4)$$

$$n(F_{P_1=K_s^3}) = 2.n(F_{P_1=K_{s-1}^3}) + n(F_{P_1=K_s^1}) + n(F_{P_1=K_s^2}), \quad (5)$$

$$n(F_{P_1=K_s^4}) = 2.n(F_{P_1=K_{s-1}^4}) + n(F_{P_1=K_s^1}) + n(F_{P_1=K_s^2}) + n(F_{P_1=K_s^3}). \quad (6)$$

Substitute (4) to (5) to get

$$n(F_{P_1=K_s^3}) = 2.n(F_{P_1=K_{s-1}^3}) + 2.n(F_{P_1=K_{s-1}^2}) + 2.n(F_{P_1=K_s^1}). \quad (7)$$

Substitute (7) to (6) to get

$$n(F_{P_1=K_s^4}) = 2.n(F_{P_1=K_{s-1}^4}) + 2.n(F_{P_1=K_{s-1}^3}) + 2^2.n(F_{P_1=K_{s-1}^2}) + 2^2.n(F_{P_1=K_{s-1}^1}).$$

By recursive process we have

$$n(F_{P_1=K_s^{m-2}}) = 2.n(F_{P_1=K_{s-1}^{m-2}}) + 2.n(F_{P_1=K_{s-1}^{m-3}}) + 2^2.n(F_{P_1=K_{s-1}^{m-4}}) + 2^3.n(F_{P_1=K_{s-1}^{m-5}}) \\ + \dots + 2^{m-4}.n(F_{P_1=K_{s-1}^2}) + 2^{m-4}.n(F_{P_1=K_{s-1}^1}),$$

and

$$n(F_{P_1=K_s^{m-1}}) = 2.n(F_{P_1=K_{s-1}^{m-1}}) + 2.n(F_{P_1=K_{s-1}^{m-2}}) + 2^2.n(F_{P_1=K_{s-1}^{m-3}}) + 2^3.n(F_{P_1=K_{s-1}^{m-4}}) \\ + \dots + 2^{m-3}.n(F_{P_1=K_{s-1}^2}) + 2^{m-3}.n(F_{P_1=K_{s-1}^1}),$$

$$n(F_{P_1=K_s^m}) = 2.n(F_{P_1=K_{s-1}^m}) + 2.n(F_{P_1=K_{s-1}^{m-1}}) + 2^2.n(F_{P_1=K_{s-1}^{m-2}}) + 2^3.n(F_{P_1=K_{s-1}^{m-3}}) \\ + \dots + 2^{m-2}.n(F_{P_1=K_{s-1}^2}) + 2^{m-2}.n(F_{P_1=K_{s-1}^1}).$$

Therefore, we have

$$n(F_{P_1=K_s^m}) = 2.n(F_{P_1=K_{s-1}^m}) + (1 + 1 + 2 + 2^2 + 2^3 + \dots + 2^{m-4} + 2^{m-3}).n(F_{P_1=K_{s-1}^1}) \\ + 2(1 + 1 + 2 + 2^2 + 2^3 + \dots + 2^{m-5} + 2^{m-4}).n(F_{P_1=K_{s-1}^2}) + 2(1 + 1 + 2 + 2^2 + 2^3 + \dots + 2^{m-6} \\ + 2^{m-5}).n(F_{P_1=K_{s-1}^3}) + \dots + 2(1 + 1 + 2).n(F_{P_1=K_{s-1}^{m-3}}) + 2(1 + 1).n(F_{P_1=K_{s-1}^{m-2}}) \\ + 2.n(F_{P_1=K_{s-1}^{m-1}}).$$

By manipulation algebraic, use the fact that

$$(1 + 1 + 2 + 2^2 + 2^3 + \dots + 2^{k-1} + 2^k) = 2^{k+1},$$

and Corollary 10 in [9] we have

$$n(F_{P_1=K_s^m}) = 2.n(F_{P_1=K_{s-1}^m}) + 2.n(F_{P_1=K_{s-1}^{m-1}}) + 2^2.n(F_{P_1=K_{s-1}^{m-2}}) + 2^3.n(F_{P_1=K_{s-1}^{m-3}}) \\ + \dots + 2^{m-3}.n(F_{P_1=K_{s-1}^3}) + 2^{m-2}.n(F_{P_1=K_{s-1}^2}) + 2^{m-1}.n(F_{P_1=K_{s-1}^1}). \quad \square$$

Theorem 7. Let G be a rectangle group with $m \times n$ in size. Then

$$n(F_{P_1=K_s^m=\{e\}}) = 2.n(F_{P_1=K_{s-1}^{m-1}}) + \sum_{i=1}^{s-1} n(F_{P_1=K_i^m}).$$

Proof. Analogue with the proof of Theorem 5. $\square$

Theorem 8. (Column Expansion) Let G be a rectangle group with $m, s \in \mathbb{N}$ and $m \geq 3$. We have

$$\begin{aligned} n(F_{P_1=K_s^m}) &= 2.n(F_{P_1=K_s^{m-1}}) + 2.n(F_{P_1=K_{s-1}^{m-1}}) + 2^2.n(F_{P_1=K_{s-2}^{m-1}}) + 2^3.n(F_{P_1=K_{s-3}^{m-1}}) \\ &\quad + \dots + 2^{s-3}.n(F_{P_1=K_3^{m-1}}) + 2^{s-2}.n(F_{P_1=K_2^{m-1}}) + 2^{s-1}.n(F_{P_1=K_1^{m-1}}). \end{aligned}$$

Proof. Analogue with the proof of Theorem 6. $\square$

References

- [1] L. A. Zadeh, Fuzzy sets, *Inform. And Control.*, **8** (1965), 338-353.
- [2] A. Rosenfeld, Fuzzy groups, *J. Math. Anal. Appl.*, **35** (1971), 512-517.
- [3] M. Tarnauceanu, The number of fuzzysubgroups of finite cyclic groups and Delanoy numbers, *European Journal of Combinatorics.*, **51** (1992), 105-109.
- [4] F. Laszlo, Structure and construction of fuzzy subgroup of a group, *Fuzzy Sets and System.*, **51** (1992), 105-109.
- [5] V. Murali and B.B. Makamba, On an equivalence of fuzzy subgroups I, *Fuzzy Sets and System.*, **123** (2001), 259-264.
- [6] V. Murali and B.B. Makamba, Counting the number of fuzzy subgroups of an abelian group of order $p^n q^m$, *Fuzzy Sets and System.*, **144** (2004), 459-470.
- [7] R. Sulaiman, Constructing fuzzy subgroups of symmetric group S_4 , *Int. Journal of Algebra.*, **6,1** (2012), 23-28.
- [8] R. Sulaiman and Abd. Ghafur, Constructing fuzzy subgroups of symmetric group S_2, S_3 and alternating group A_4 , *Journal of Quality Measurement and Analysis.*, **6,1** (2010), 57-63.
- [9] Raden Sulaiman, Fuzzy subgroups computation of finite group by using their lattice, *Int. Journal of Pure and Applied Mathematics.*, **78** (2012), 479-489.
- [10] R. Sulaiman and Budi Priyo, The number of fuzzy subgroups of rectangle groups, *Int. Journal of Algebra.*, **8,1**(2014), 17-23.

Computing the number of fuzzy subgroups by expansion method

ORIGINALITY REPORT

24%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|---|-----------------|
| 1 | R. Sulaiman, Budi Priyo Prawoto. "The number of fuzzy subgroups of rectangle groups", International Journal of Algebra, 2014
<small>Crossref</small> | 167 words — 11% |
| 2 | Nesmith, E., and J.M. Jean-Michel. "THE EFFECT OF IN-FLIGHT MASS DEPLETION ON BIRD MIGRATION RANGE", International Electronic Journal of Pure and Applied Mathematics, 2014.
<small>Crossref</small> | 37 words — 2% |
| 3 | jacr.iausari.ac.ir
<small>Internet</small> | 35 words — 2% |
| 4 | www.m-hikari.com
<small>Internet</small> | 31 words — 2% |
| 5 | jmua.fmipa.unand.ac.id
<small>Internet</small> | 20 words — 1% |
| 6 | Pilev, D.. "WEB-BASED INFORMATION SYSTEM SECURITY", International Electronic Journal of Pure and Applied Mathematics, 2014.
<small>Crossref</small> | 16 words — 1% |
| 7 | Studies in Fuzziness and Soft Computing, 2015.
<small>Crossref</small> | 12 words — 1% |
| 8 | www.mayfeb.com
<small>Internet</small> | 12 words — 1% |

-
- 9 Abou-Zaid, S.. "On generalized characteristic fuzzy subgroups of a finite group", Fuzzy Sets and Systems, 19910925 11 words — 1%
Crossref
-
- 10 www.e.ijpam.eu 10 words — 1%
Internet
-
- 11 Cernea, Aurelian. "ON A HIGHER ORDER DIFFERENTIAL INCLUSION WITH FOUR-POINT INTEGRAL BOUNDARY CONDITIONS", International Electronic Journal of Pure and Applied Mathematics, 2014. 10 words — 1%
Crossref
-
- 12 Songfang Jia, , Yanheng Chen, Jinkui Liu, and Youyi Jiang. "On the number of fuzzy subgroups of finite abelian p -groups with type $(p^{\sup n}, p^{\sup m})$ ", 2011 3rd International Conference on Computer Research and Development, 2011. 10 words — 1%
Crossref
-
- 13 Marian, S.L., M.R. Sagayaraj, A.G.M. Selvam, and M.P. Loganathan. "OSCILLATORY BEHAVIOR OF FORCED FRACTIONAL DIFFERENCE EQUATIONS", International Electronic Journal of Pure and Applied Mathematics, 2014. 9 words — 1%
Crossref
-

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF