

3.The efficacy of Salvinia molesta Mitch. and Marsilea crenata Presl. as phytoremediators of lead pollution.pdf

by

Submission date: 02-Aug-2019 09:52AM (UTC+0700)

Submission ID: 1156925570

File name: 3.The efficacy of Salvinia molesta Mitch. and Marsilea crenata Presl. as phytoremediators of lead pollution.pdf (501.65K)

Word count: 3889

Character count: 18173

The efficacy of *Salvinia molesta* Mitch. and *Marsilea crenata* Presl. as phytoremediators of lead pollution

F. Rachmadiarti* and G. Trimulyono

Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Surabaya, Ketintang Campus, Surabaya, Jawa Timur, Indonesia. *E-mail: fidarachmadiarti@unesa.ac.id, capa.journal61@gmail.com

Abstract

This study focused on the absorption of lead (Pb) by two species of ferns, *Salvinia molesta* Mitch and *Marsilea crenata* Presl. The aim of this research was to describe the accumulation of lead in *S. molesta* and *M. crenata* roots and leaves, and in addition, to find whether these are accumulators or excluders. Both of these ferns were grown in a hydroponic system to which there was added Pb at 0, 5, 10, and 15 ppm concentration, exposed for 7 and 14 days. After the treatment, determination of the lead amounts in the roots and leaves used the AAS method and the RGR calculation. The data were analyzed by ANOVA using SPSS Program 16 edition. The results showed that the accumulation of lead was higher in the roots than the leaves. Both of the ferns had a bioconcentration factor of more than 1 and the factor of translocation was low, i.e., less than 1. It is concluded that *S. molesta* and *M. crenata* are appropriate species for phytoremediation, especially as excluders.

Key words: *Salvinia molesta*, *Marsilea crenata*, lead, phytoremediation

Introduction

Water pollution by heavy metals is an emerging issue (Garbisu and Alkorta, 2003; Kachenko *et al.*, 2007b). One of the causes of heavy metal pollution is human activity, including the use of chemical fertilizers in the agriculture sector and industrial emissions. According to the United States Environmental Protection Agency (EPA), lead (Pb) is one of 'the big three' heavy metal pollutants, along with mercury (Hg) and cadmium (Cd), these having the highest level of danger to human health because of their very high level of toxicity even at low concentration levels. Lead can cause teratogenicity effects. In addition, the lead may also inhibit the synthesis of hemoglobin, causing kidney malfunction and affect the reproductive system and the cardiovascular system. It also may cause acute and chronic damage to the central nervous system and peripheral nervous system (Dunibe *et al.*, 2007). Lead is not essential for metabolism and may cause cancer in humans. Therefore, there should be an effort to decrease lead contamination in water.

One method that has been used to reduce the levels of heavy metals is through chemical physics and remediation that is expensive and may degrade the environment in its own right. Another method that can be used is phytoremediation. Phytoremediation is an environmentally friendly and economically feasible technology for mitigating water/soil contamination using plants (Cheng *et al.*, 2015; Ghosh & Singh, 2005; Gonzaga *et al.*, 2008). The ability of plants to reduce the metal content in the water can be calculated as the bioaccumulation factor (BAF = metal in shoots/metal in media) and the bioconcentration factor (BCF = metal in roots/metal in media), while the ability of plants to move metal from roots to shoots is defined as the translocation factor (TF = metal in shoot/metal in root) (Yoon *et al.*, 2006). Basically, from the values derived from these parameters, plants that extract metal are known as excluders or accumulators. Excluders, with TF < 1, and BAF and BCF > 1 can absorb metals in the roots, termed as

phytostabilization (Comara *et al.*, 2007). Accumulators (TF > 1), also known as hyper accumulators, move the metal from the roots to shoots. Hyper accumulators can accumulate more than 10,000 mg/kg in parts of the plant above the surface without causing any adverse effects.

Few ferns, including *Pteridium* sp are hyper accumulators along with some higher plants (Ghosh *et al.*, 2005). A few ferns have been researched in terms of their absorption capacity; however, the potential of water ferns *Marsilea crenata* (water clover) and *Salvinia molesta* (salvinia) as hyper accumulators or excluders in the tropical waters has not been reported.

The results of an initial study showed that *S. molesta*, from Porong, Sidoarjo, East Java, growing in polluted water contained lead concentrations of 0.5 ppm and *M. crenata*, from Kendung, Sidoarjo, East Java, lead concentrations of 0.8 ppm. Each of these species showed good growth and did not show any adverse signs growing in the polluted water. In this study, an exploration of the mechanisms of lead accumulation into both of these species was conducted. The presence of lead in both of these species could be observed by the color changes of the leaves. Based on this, the aim of this study was to describe the distribution of the lead in the roots and shoots of these water ferns; in addition, the study was also intended to evaluate the possibility of using them for phytoremediation of polluted waters in tropical areas.

Materials and methods

To study the potency of *S. molesta* and *M. crenata* as the phytoremediation agents for the lead, the plants were grown in a medium containing Pb solution. To the growth medium Pb solution of various concentrations was added, i.e.: (1) 0 ppm of Pb as the control; and (2) 5 ppm; (3) 10 ppm; (4) 15 ppm of Pb(NO₃)₂ solution. These treatments were arranged in a randomized block design through three replications.

S. molesta was collected from Porong, Sidoarjo, East Java, and *M. crenata* from Kendung, Surabaya, East Java. Both of these ferns were adapted and grown in a hydroponic system by using a batch system in Hoagland's solution during seven days. After adapting them to the procedure, the plants were cleaned with distilled water, and then put in glass aquarium of 40 cm length, 30 cm width and 35 cm depth containing 5 L distilled water and Hoagland's solution or the plants were moved into an aquarium containing 5 L of lead/Hoagland's solution (0, 5, 10 and 15 ppm Pb) and exposed for 7 and 14 days. The ferns were grown under a 12-hour light-dark cycle. The plant samples from each aquarium were harvested after either 7 and 14 days. The pH of the medium varied from 5.4-6.7 at the beginning of the experiment, to 6.5 -7.0 at the end of the experiment.

Overall biomass, yield and metal content were measured as specified below. Harvested plants were divided into two parts: roots and leaves. Plant biomass yield was measured on dry weight basis. Each part was oven-dried for 2 hours at 70 °C on successive days and weighed to determine dry weight. Dried plant tissues were milled into a powder. Powder samples weighing 0.5 g were digested with 5 mL of HNO₃ and diluted to 50 mL with de-ionized double distilled water. The lead analysis was conducted through an extraction method. Both media samples (50 mL) and digested plant samples were analyzed for lead by using an atomic absorption spectrophotometer. The total accumulation and partitioning of the metals by the plants were calculated.

Relative growth rate (RGR) is stated in grams per kilogram per day, and it was calculated as follows:

$$\text{RGR} = \frac{(\text{Final dry} - \text{Initial dry weight})}{\text{Initial weight} \times \text{days}} \quad (1)$$

The bioconcentration factor (BCF) for the lead of the root system and the upper parts (stem/peduncles leaves) were calculated according to the formula as proposed by Wang *et al.* (2007), i.e.:

$$\text{BCF} = C_{\text{roots}} / C_{\text{medium}} \quad (2)$$

$$\text{BAF} = C_{\text{shoots}} / C_{\text{medium}} \quad (3)$$

Where, C_{roots} = concentration of lead in the roots (mg Pb kg⁻¹ dry weight)

C_{shoots} = total of the concentration of lead in stems or peduncles (mg Pb kg⁻¹ dry weight)

C_{medium} = concentration of lead in growth medium (mg Pb L⁻¹).

The translocation factor (TF) was calculated to evaluate the translocation of Pb from the *M. crenata* and *S. molesta* roots to shoots (stem/petiole and leaves) by using equations (2) and (3) as follows:

$$\text{TF} = \text{BAF}_{\text{shoots}} / \text{BCF}_{\text{roots}} \quad (4)$$

Analysis of variance (ANOVA) was performed using SPSS 16 software package to analyze the data. If there were significant differences, the Tukey test ($P=0.05$) was used.

Results and discussion

There were no significant differences in growth between fern species or the duration of exposure to lead (7 days vs. 14 days) among treatments. The growth of *S. molesta* and *M. crenata* was

Table 1. RGR of *S. molesta* and *M. crenata* treatments after 7 and 14 days planted ($P=0.05$)

Species	Concentration Pb (ppm)	RGR	
		7 days	14 days
<i>Salvinia molesta</i>	0	0.08±0.00b	0.04±0.00b
	5	0.00±0.00a	0.00±0.00a
	10	0.00±0.00a	0.00±0.00a
	15	-0.02±0.04a	0.00±0.00a
<i>Marsilea crenata</i>	0	0.03±0.08b	0.01±0.0b
	5	0.01±0.00a	0.00±0.00a
	10	0.01±0.00a	0.00±0.00a
	15	0.01±0.00a	0.00±0.00a

Table 2. Percentage removal of Pb by *S. molesta* and *M. crenata*, with Pb concentration (ppm) ($P=0.05$)

Species	Concentration Pb (ppm)	Pb removal	
		7 days	14 days
<i>Salvinia molesta</i>	0	0.00±0.00a	0.00±0.00a
	5	97.33±0.12b	97.33±0.12b
	10	98.50±0.00c	98.57±0.12c
	15	98.96±0.04b	98.89±0.13b
<i>Marsilea crenata</i>	0	0.00±0.00a	0.00±0.0a
	5	94.93±2.55b	86.67±1.14b
	10	96.27±0.47c	91.47±1.50c
	15	97.22±0.65b	80.07±5.5b

influenced by the concentration of the lead in the growth media (Table 1). The growth of both water fern species at days 7 and 14 was characterized by a positive value of RGR, generally equal to or higher than 0, although the RGR values of the treatment groups tended to be smaller than the control group.

Based on the results presented in Table 1, it can be concluded that both water fern species were able to grow adequately well in the presence of Pb. The favorable growth of these plants was accompanied by the removal of lead metal from the media (Table 2).

The concentration of the lead in the root tissues of both species was relatively constant in accord with the age of the plants. In root tissues of *S. molesta* and *M. crenata*, the concentration of lead increased from 5 to 15 ppm at 7 and 14 days after planting. Corroborating the data presented in Table 3, it seems that the increase of Pb occurred at each treatment concentration. For both species, on the other hand, the concentration of the lead in the stem and leave tissues increased as the plants growing older. The results presented in Table 4 show that until the 14th day, TF of both *S. molesta* and *M. crenata* were less than 1.0, showing that the metal ion was largely retained in the roots. The results support previous data which show that the highest concentration of the lead was in the root tissue.

S. molesta and *M. crenata* were able to grow in a solution of up to 15 ppm Pb, but the ability to displace lead from media was at best at concentrations of 10 ppm. The removal of lead by both species at concentrations of 5 and 15 ppm were not significantly different. From data presented in Table 2, both of these plants were able to remove lead metal from media up to 98 %. More

Table 3. Mean and Sd of Pb content in roots and leaves (ppm) of *S. molesta* and *M. crenata* in different Pb treatments after 7 and 14 days of planting

Species	Concentration Pb (ppm)	7 days		14 days	
		Roots	Leaves	Roots	Leaves
<i>Salvinia molesta</i>	0	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.00±0.00a
	5	0.59±0.02b	0.29±0.00b	0.64±0.00b	0.39±0.00b
	10	0.63±0.00c	0.33±0.01c	0.67±0.02c	0.43±0.00c
	15	0.64±0.00d	0.41±0.01d	0.73±0.01d	0.47±0.01d
<i>Marsilea crenata</i>	0	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.00±0.00a
	5	0.68±0.14b	0.18±0.01b	0.94±0.06b	0.22±0.07b
	10	1.45±0.08c	0.19±0.02e	1.69±0.11c	0.58±0.05c
	15	1.63±0.35d	0.78±0.02d	2.92±0.43d	0.41±0.01d

lead in the media resulted in an increasingly higher concentration of lead in both *S. molesta* and *M. crenata*, both in the roots and leaves. However, the concentration of lead in the roots was significantly higher than in the leaves.

In order to help characterize the ability of the plants to concentrate lead from the media, the bioconcentration factor (BCF), bioaccumulation factor (BAF) and translocation factor (TF) were calculated. The results are presented in Table 4, which shows the higher the concentrations of lead in the media, the higher the TF value; the value of TF at 14 days, however, was very low and less than 1.0. Similarly, for the BCF and BAF, the higher concentrations of the treatment, the higher value of the BCF and BAF. Table 4 shows that the value of BCF for the roots is greater than the value of BAF for the shoots.

The greatest growth of *S. molesta* and *M. crenata* occurred in the control group, in which the results showed that the biomass increased rapidly in the treatment for 7 and 14 days compared with the treatment group. With increased concentrations of lead in the growing media, growth was slightly slower; both of the species had a positive growth rate even up to concentrations of lead up to 15 ppm in the growth media. Positive values of RGR indicated that both species were able to have adapted morphologically to the pressure of lead present in the media. The adaptation could be exhibited by these plants by establishing specific amino acids.

There was an increased displacement of lead from the growing media, which seemed to be caused by the ability of the roots of both species to absorb Pb ions from the growing media, and then translocate them to other parts of the plant, like the shoot. Roots of both species are fibrous and have a modified layer form of epidermal layer similar to root hairs that have a high ability to absorb nutrients and other substances. The higher concentration of lead may decrease the level of absorption significantly from

the growing media, with an optimum concentration of 10 ppm. At concentrations of 15 ppm, the roots of plants may have been in a saturated condition such that the displacement of metals from the growing media was decreased.

The experimental results presented in Table 3 show that the highest lead concentrations in *S. molesta* and *M. crenata* was found in the roots tissues. The mechanism behind the higher lead concentration in the roots may include the binding process of the positively charged heavy metal ions to negative charges in the cell walls (Grill *et al.*, 1995). Gothberg *et al.* (2002) suggested that in most plants, a larger proportion of metal is retained in the roots, and thereby it may be prevented from interfering with sensitive metabolic reactions in the shoots. The results are similar to other species *S. minima* (Alvarado *et al.*, 2016). This is probably an internal mechanism to avoid toxic metal concentrations in the shoots.

Both, *S. molesta* and *M. crenata* could be categorized as accumulator plants. In addition, Yoon *et al.* (2006) stated that the plants can be classified as excluders and accumulators if the value of TF < 1 and BCF > 1. The higher values of BCF and TF of the tested species may enable them to accumulate large amount hazardous metals in the harvested parts. The proper disposal of the harvested plant parts is the final and most important step in any kind of plant-based remediation technology. If it is not disposed off properly, the accumulated heavy metals may turn to the system or may enter into the food chain. However, bioaccumulation of the trace metals in *M. crenata* tissues was not a concern because the measured levels were lower than the permissible levels set by FAO and WHO (0.5 mg kg⁻¹) for consumption.

The results of this study showed that both *S. molesta* and *M. crenata* have relatively high values of BCF and relatively low value of TF. Patterns of accumulation of both ferns were in accord

Table 4. Mean and Sd of translocation factors (TF), bioconcentration factors (BCF), bioaccumulation factor (BAF) for the different Pb treatments after 7 and 14 days of planting

Species	Concentration Pb (ppm)	7 days			14 days		
		BCF	BAF	TF	BCF	BAF	TF
<i>Salvinia molesta</i>	0	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.00±0.00a
	5	4.40±0.05b	2.20±0.05b	0.50±0.01b	4.81±0.20b	2.95±0.08b	0.61±0.01b
	10	4.24±0.03c	2.22±0.08b	0.52±0.02b	4.73±0.50c	2.99±0.15	0.62±0.01b
	15	3.85±0.03b	2.11±0.08b	0.62±0.01c	4.40±0.49b	2.85±0.03b	0.63±0.01c
<i>Marsilea crenata</i>	0	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.00±0.00a
	5	3.71±0.87b	0.77±0.05b	0.00±0.00b	1.85±0.02b	0.35±0.04b	0.00±0.00b
	10	6.13±0.54c	0.53±0.06b	0.00±0.00b	3.24±0.06c	0.68±0.07b	0.00±0.00b
	15	6.56±0.56b	0.51±0.08b	0.09±0.15c	1.77±0.04b	0.01±0.00b	0.13±0.00c

with previous research that ferns have high value of BCF and low value of T_{trans} and adapted to high concentrations of metal by immobilizing metals and metalloids in the root system, reducing the translocation of these metals to the shoots (Kachenko *et al.*, 2007a). This strategy seems to occur in ferns generally, as sequestration in the root system of various metals and metalloids were also observed in some species in Australia (Kachenko *et al.*, 2007b). Tolerant mechanism of the metal accumulation is a result of continuous flow through phloem tissue and resulting saturation sequestration in root/rhizome of plants. This is a collector-type system in these plants (metal excluders), contrasting with the mechanism that has been reported in the genus *Pteridium*, which is capable of transferring metal from root to shoot tissue. The bioconcentration factor of lead in the root system/rhizome has been also found in *Adiantum* species such as water spinach and yellow velvetleaf. This is supported by investigations, which show that many wetland plants can collect heavy metals from polluted waters, and are found to tolerate a wide range of heavy metals (Pb, Zn, Cu, and Cd), and have demonstrated their potential use in phytoremediation (Leguizamo *et al.*, 2017). Further research is required to increase our understanding of the potential of local plant species in wetlands as phytoremediators.

The results of this study show that the water ferns *S. molesta* and *M. crenata* primarily accumulated lead in their roots rather than leaves. Both ferns have shown a bioconcentration factor greater than 1.0, and a low translocation factor, less than 1.0. In short, *S. molesta* and *M. crenata* are suitable as excluder phytoremediators in heavy metal-polluted waters, as these ferns were able to regulate the accumulation of Pb and limit the metal transport to the shoots of the plant, allowing for continued growth and thus easier harvest at the water surface, in order to reduce overall heavy metal levels in a polluted water body.

Acknowledgments

We would like to extend our gratitude to Ministry of Research and Technology of Indonesia that has funded this research through Competitive Research Grants in 2015-2016. We also thank the students involved in this research, especially Maria Ulfa and Rizki.

References

Alvarado, D.A.L., F.E. Gil, L.S. Carbonell, C.T. May and J.M. Santamaria, 2016. Lead accumulation reduces photosynthesis in the lead hyper-accumulator *Salvinia minima* Baker by affecting the cell membrane and inducing stomatal closure. *Aquat. Toxicol.*, 171: 37-47.

- Cheng, S.F., C.Y. Huang, Y.C. Lin, S.C. Lin and K.L. Chen, 2015. Phytoremediation of lead using corn in contaminated agricultural land-An in situ study and benefit assessment. *Ecotoxicol. Environ. Saf.*, 111: 72-77.
- Comara, L., E. Rocciotiello, V. Minganti, G. Drava, R. De Pellegrini and M.G. Mariotti, 2007. The level of trace elements in plants growing on serpentine and metalliferous soils. *J. Plant Nut. Soil Sci.*, 170: 781-787.
- Dunibe, J.O., M.O.C. Ogwuebu and J.N. Ekwurugwu, 2007. Heavy metal pollution and human biotoxic effects. *Int. J. Phys. Sci.*, 2(5): 112-118.
- Garbisu, C. and I. Alkorta, 2003. Basic concepts on heavy metal soil bioremediation. *Eur. J. Miner. Process Environ. Prot.*, 3: 58-66. Available from: http://www.ejmpep.com/garbisu_and_alkorta.pdf
- Ghosh, M. and S.P. Singh, 2005. A review on phytoremediation of heavy metals and utilization of it's by products. *Appl. Ecol. Environ. Res.*, 2005, 3(01): 1-18. Available from: http://www.aloki.hu/pdf/0301_001018.pdf
- Grill, E., L. Winnacker and M.H. Zenk, 1985. Phytochelatin, the principal heavy-metal complexing peptides of higher plants. *Science*, 230(4726): 674-676.
- Gonzaga, M.I.S., J.A.G. Santos and L.Q. Ma, 2006. Arsenic phytoextraction and hyperaccumulation by fern species. *Scientia Agricola*, 63(1): 90-101.
- Göthberg, A., M. Greger, K. Holm and B.E. Bengtsson, 2002. Influence of nutrient levels on uptake and effect of mercury, cadmium, and lead in water spinach. *J. Environ. Qual.*, 33(4): 1247-55.
- Kachenko, A.G., S. Balwan and P.B. Naveen, 2007a. Heavy metal tolerance in common fern species. *Aus. J. Bot.*, 55: 63-73.
- Kachenko, A.G., N.P. Bhatia, B. Singh and R. Siegel, 2007b. Arsenic hyperaccumulation and localization in the pinnule and stipe tissues of the gold-dust fern (*Pytyrogramma calomelanos* (L.) Link var. *austroamericana* (Domin) Farw.) using quantitative micro-PIXE spectroscopy. *Plant Soil*, 300: 207-219.
- Leguizamo, M.A.O., W.D.F. Gomez and M.C.G. Sarmiento, 2017. Native herbaceous plant species with potential use in phytoremediation of heavy metals, spotlight on wetlandsd- A review. *Chemosphere*, 168: 1230-1247.
- Wang, M., J. Zou and X. Duan, 2007. Cadmium accumulation and its effects on metal uptake in maize (*Zea mays* L.). *Bioresource Technol.*, 98: 82-88.
- Yoon, J., X. Cao, Q. Zhou and L.Q. Ma, 2006. Accumulation of Pb, Cu, and Zn in native plants growing on a contaminated Florida site. *Sci. Total Environ.*, 368: 456-64

Received: March, 2017; Revised: June, 2017; Accepted: June, 2017

3.The efficacy of *Salvinia molesta* Mitch. and *Marsilea crenata* Presl. as phytoremediators of lead pollution.pdf

ORIGINALITY REPORT

20%

SIMILARITY INDEX

%

INTERNET SOURCES

20%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

Huan, Y.. "Changes in flavor compounds of dry-cured Chinese Jinhua ham during processing", *Meat Science*, 200510

Publication

4%

2

P.C. Abhilash, Vimal Chandra Pandey, Pankaj Srivastava, P.S. Rakesh, Smitha Chandran, Nandita Singh, A.P. Thomas. "Phytofiltration of cadmium from water by *Limnocharis flava* (L.) Buchenau grown in free-floating culture system", *Journal of Hazardous Materials*, 2009

Publication

3%

3

Enrica Roccotiello, Alice Manfredi, Giuliana Drava, Vincenzo Minganti, Mauro Giorgio Mariotti, Graziella Berta, Laura Cornara. "Zinc tolerance and accumulation in the ferns *Polypodium cambricum* L. and *Pteris vittata* L.", *Ecotoxicology and Environmental Safety*, 2010

Publication

3%

4

Oyuela Leguizamo, Mayerly Alexandra, Wilmar Darío Fernández Gómez, and Martha Cecilia

1%

Gutiérrez Sarmiento. "Native herbaceous plant species with potential use in phytoremediation of heavy metals, spotlight on wetlands — A review", Chemosphere, 2016.

Publication

5

M.H. Hu, Y.S. Ao, X.E. Yang, T.Q. Li. "Treating eutrophic water for nutrient reduction using an aquatic macrophyte (*Ipomoea aquatica* Forsskal) in a deep flow technique system", Agricultural Water Management, 2008

Publication

6

Lotfy, S.M., and A.Z. Mostafa. "Phytoremediation of contaminated soil with cobalt and chromium", Journal of Geochemical Exploration, 2013.

Publication

7

"Phytoremediation", Springer Nature, 2017

Publication

8

T. S. T. Mansor, Y. B. Che Man, M. Shuhaimi. "Employment of Differential Scanning Calorimetry in Detecting Lard Adulteration in Virgin Coconut Oil", Journal of the American Oil Chemists' Society, 2011

Publication

9

Cai, Q.Y.. "Potential of *Ipomoea aquatica* cultivars in phytoremediation of soils contaminated with di-n-butyl phthalate",

1 %

1 %

1 %

1 %

1 %

10

Phytoremediation, 2015.

Publication

<1 %

11

Demim, S., N. Drouiche, A. Aouabed, T. Benayad, O. Dendene-Badache, and S. Semsari. "Cadmium and nickel: Assessment of the physiological effects and heavy metal removal using a response surface approach by L. gibba", Ecological Engineering, 2013.

Publication

<1 %

12

Castro, Paula M. L., Marques, Ana P. G. C. and Rangel, António O. S. S.. "Zinc accumulation in plant species indigenous to a portuguese polluted site: relation with soil contamination", American Society of Agronomy, 2011.

Publication

<1 %

13

Shu-Fen Cheng, Chin-Yuan Huang, Yung-Cheng Lin, Sheng-Chien Lin, Kuo-Lin Chen. "Phytoremediation of lead using corn in contaminated agricultural land—An in situ study and benefit assessment", Ecotoxicology and Environmental Safety, 2015

Publication

<1 %

14

Ali Seid Mohammed. "Heavy Metal Pollution: Source, Impact, and Remedies", Environmental

<1 %

15

Valencia, I.. "Development of dry fermented sausages rich in docosaehaenoic acid with oil from the microalgae Schizochytrium sp.: Influence on nutritional properties, sensorial quality and oxidation stability", Food Chemistry, 2007

Publication

<1 %

16

Mamdouh A. Eissa, Yaser A. Almaroai. "Phytoremediation Capacity of Some Forage Plants Grown on a Metals-Contaminated Soil", Soil and Sediment Contamination: An International Journal, 2019

Publication

<1 %

17

Ekmekci, Y.. "Effects of cadmium on antioxidant enzyme and photosynthetic activities in leaves of two maize cultivars", Journal of Plant Physiology, 20080418

Publication

<1 %

18

Chirawee Sangthong, Kunchaya Setkit, Benjaphorn Prapagdee. "Improvement of cadmium phytoremediation after soil inoculation with a cadmium-resistant Micrococcus sp.", Environmental Science and Pollution Research, 2015

Publication

<1 %

19

Abhilash, P.C.. "Phytofiltration of cadmium from water by *Limnocharis flava* (L.) Buchenau grown in free-floating culture system", *Journal of Hazardous Materials*, 20091030

Publication

<1 %

20

Ky, Isabelle, and Pierre-Louis Teissedre. "Characterisation of Mediterranean Grape Pomace Seed and Skin Extracts: Polyphenolic Content and Antioxidant Activity", *Molecules*, 2015.

Publication

<1 %

21

Carattino, M.D.. "Effects of long-term exposure to Cu^{2+} and Cd^{2+} on the pentose phosphate pathway dehydrogenase activities in the ovary of adult *Bufo arenarum*: possible role as biomarker for Cu^{2+} toxicity", *Ecotoxicology and Environmental Safety*, 200403

Publication

<1 %

22

Saud A Alamri, Manzer H Siddiqui, Mutahhar YY Al-Khaishany, M. Nasir Khan et al. "Ascorbic acid improves the tolerance of wheat plants to lead toxicity", *Journal of Plant Interactions*, 2018

Publication

<1 %

23

"Detoxification of Heavy Metals", Springer Science and Business Media LLC, 2011

Publication

<1 %

Claudia S. Benimeli. "Bioaccumulation of

24

Copper by Zea mays: Impact on Root, Shoot and Leaf Growth", Water Air and Soil Pollution, 11/10/2009

Publication

<1 %

25

Gülsün Özyurt, Yesim Ozogul, Esmeray Kuley Boga, A. Serhat Özkütük, Mustafa Durmuş, Yılmaz Uçar, Fatih Ozogul. "The Effects of Fermentation Process with Acid and Lactic Acid Bacteria Strains on the Biogenic Amine Formation of Wet and Spray-Dried Fish Silages of Discards", Journal of Aquatic Food Product Technology, 2019

Publication

<1 %

26

Kay, S.H.. "Effects of heavy metals on water hyacinths (*Eichhornia crassipes* (Mart.) Solms)", Aquatic Toxicology, 198405

Publication

<1 %

27

Bruno Barbosa, Sara Boléo, Sarah Sidella, Jorge Costa et al. "Phytoremediation of Heavy Metal-Contaminated Soils Using the Perennial Energy Crops *Miscanthus* spp. and *Arundo donax* L.", BioEnergy Research, 2015

Publication

<1 %

28

Pankaj Kumar, Madhusudan Hiranman Fulekar. "Rhizosphere Bioremediation of Heavy Metals (Copper and Lead) by *Cenchrus ciliaris*", Research Journal of Environmental Sciences,

<1 %

2018

Publication

Exclude quotes On

Exclude bibliography On

Exclude matches Off