

Turnitin_SOME PROPERTIES OF PRIMITIVE HENSTOCK OF INTEGRABLE FUNCTION IN LOCALLY COMPACT METRIC SPACE OF VECTOR VALUED FUNCTION

by Manuharawati Manuharawati

Submission date: 15-Oct-2019 02:40PM (UTC+0700)

Submission ID: 1193180382

File name: Extracted_pages_from_paper_Manuharawati.pdf (608.19K)

Word count: 1853

Character count: 10795

SOME PROPERTIES OF PRIMITIVE ϑ –HENSTOCK OF INTEGRABLE FUNCTION IN LOCALLY COMPACT METRIC SPACE OF VECTOR VALUED FUNCTION

Manuharawati

Mathematics Department
Faculty of Mathematics and Natural Science-The State University of Surabaya
manuhara1@yahoo.co.id

Abstract

By abstracting the domain and range of Henstock integrable function and measures (the volume function) used by mathematicians, we had constructed ϑ –Henstock integral in locally compact metric space of vector valued function. Based on the result of previous researchers, this paper discuss properties of primitive ϑ –Henstock integrable function in locally compact metric space of vector valued function.

Key words ϑ –Henstock integrable function , locally compact metric space, vector.

1. INTRODUCTION

1.1 Background

In 1989, Lee had constructed Henstock integral of real valued function on a cell $[a, b] \subset \mathbb{R}$. Manuharawati (2002) generalized this integral by reduced a cell $[a, b] \subset \mathbb{R}$ to a cell in a locally compact metric space and Cao (1992) generalized this integral by reduced the range of function from $\mathbb{R}$ to Banach space.

Based on the result of these researchers, Manuharawati and Yunianti (2013) had constructed a Henstock integral of vector valued function (the range of function is a Banach space) in locally compact metric space. In this integral, we used a volume function ϑ that defined on the set of all elementary sets in a locally compact metric space. In this paper, we discuss some properties of the primitive this integral.

1.2 Fundamental Concepts

Let X be a locally compact metric space, $\mathbb{R}^*$ be an extended real numbers, $\mathbb{R}^+$ be a positive real numbers, and $E(X)$ be a collection of elementary sets in X .

A nonvoid collection $S \subset 2^X$ is called a **system of intervals** if it satisfies: (Manuharawati, 2002: 11-12):

- (i) for every $p \in X$, $\{p\} \in S$,
- (ii) for every $p \in X$, if $d(N(p, r))$ compact then $N(p, r) \in S$,

- (iii) if $A \in S$ then A konvex, $d(A)$ compact, $d(A) \in S$, and $int(A) \in S$,
 (iv) for every $A, B \in S$, $A \cap B \in S$.
 (v) for every $A, B \in S$ there exists a collection of nonoverlapping sets $C_i \in S$, $1 \leq i \leq n$ such that

$$A - B = \bigcup_{i=1}^n C_i.$$

Every an element of S is called an **interval**. Interval $A \in S$ is said to be **degenerate** if $int(A) = \emptyset$ and **nondegenerate** if $int(A) \neq \emptyset$. A **cell** is a nondegenerate compact interval. A set $A \subset X$ is called **elementary set** if A is a finite union of intervals.

A Volume function $\vartheta: E(X) \rightarrow \mathbb{R}$ is called a measure on $E(X)$ if:

- (i) $\vartheta(A) = 0$ if $int(A) = \emptyset$ and $\vartheta(A) > 0$ if $int(A) \neq \emptyset$
 (ii) $\vartheta(A) \leq \vartheta(B)$ for every $A, B \in E(X)$ with $A \subset B$.
 (iii) $\vartheta(\bigcup_{i \in \mathbb{N}} A_i) = \sum_{i \in \mathbb{N}} \vartheta(A_i)$ for every nonoverlapping collection sets $\{A_i\} \subset E(X)$ with $\bigcup_{i \in \mathbb{N}} A_i \in E(X)$.

In this paper X is a nondiscrete locally compact metric space.

If E is a cell and $\delta: E \rightarrow \mathbb{R}^+$, then a finite collection of cell-point pairs

$$P = \{(A_{x_i}, x_i), 1 \leq i \leq n\} = \{(A_x, x)\}$$

is called a **Perron δ -fine partition on E** if $x_i \in A_{x_i} \subset N(x_i, \delta(x_i))$ and $\{A_{x_i}\}$ is a partition on E .

If $E \subset X$ is a cell and $\delta: E \rightarrow \mathbb{R}^+$, then there exists a Perron δ -fine partition on E (Manuharawati, 2002: 18).

A real function $\|\cdot\|$ on a vector space Y (over field K) is called a norm on Y if for every $x, y \in Y$ and $\alpha \in K$ satisfies:

- (i) $\|x\| \geq 0$,
 (ii) $\|x\| = 0 \Rightarrow x = \mathbf{0}$,
 (iii) $\|\alpha x\| = |\alpha| \|x\|$,
 (iv) $\|x + y\| \leq \|x\| + \|y\|$.

A vector space Y together a norm is called a norm space. A Banach space is a complete norm space.

A function $g: X \rightarrow Y$ is said to be Henstock ϑ -integrable on a cell $E \subset X$ there exists a vector $a \in Y$ such that for every real number $\varepsilon > 0$ there exists a function $\delta: E \rightarrow \mathbb{R}^+$ such that for every Perron δ -fine partition

$$P = \{(A_{x_i}, x_i), 1 \leq i \leq n\} = \{(A_x, x)\}$$

on E we have

$$\|\sum_{i=1}^n g(x_i) \vartheta(A_{x_i}) - a\| = \|\sum g(x) \vartheta(A_x) - a\| < \varepsilon$$

If $A \in S$ with $\vartheta(A) = 0$, then integral- ϑ Henstock of a function $g: X \rightarrow Y$ on A is defined by

null vector $\mathbf{0}$. The set of all ϑ -Henstock integrable function on a cell E is denoted by $H(E, \vartheta)$ and a vector a is called ϑ -Henstock integral value of a function f on E and denoted by

$$(H - \vartheta) \int f = a.$$

(Manuharawati and Yunianti, 2013: 20-21).

Some properties of the ϑ -Henstock integrable function which will be used in discussion are as follow.

Theorem 1.2.2: Cauchy's Criterion (Manuharawati and Yunianti, 2013a: 71-75): *A function $g \in H(E, \vartheta)$ if and only if for any real number $\varepsilon > 0$ there exists a function $\delta: E \rightarrow \mathbb{R}^+$ such that for any Perron δ -fine partition*

$$P = \{(A_x, x)\} \text{ and } Q = \{(B_y, y)\}$$

on E we have

$$\|P \sum g(x) \vartheta(A_x) - Q \sum g(y) \vartheta(B_y)\| < \varepsilon.$$

Theorem 1.2.2: (Manuharawati and Yunianti, 2013: 24) *If $g \in H(E, \vartheta)$, Then for every a cell $F \subset E$, $g \in H(F, \vartheta)$.*

Theorem 1.2.3: (Manuharawati, 2013: 22-26) *A function $g: X \rightarrow Y$ is Henstock ϑ -integrable on a cell $E \subset X$ if only if there exists an additive function $G: I(E) \rightarrow Y$ with property for any real number $\varepsilon > 0$ there exists a function $\delta: E \rightarrow \mathbb{R}^+$ such that for any Perron δ -fine partition*

$$P = \{(A_{x_i}, x_i), 1 \leq i \leq n\} = \{(A_x, x)\}$$

on E ,

$$\|\sum_{i=1}^n g(x_i) \vartheta(A_{x_i}) - G(E)\| = \|P \sum g(x) \vartheta(A_x) - G(E)\| < \varepsilon.$$

If E is a cell and $I(E)$ is a collection of all subinterval in E , then an additive function $G: I(E) \rightarrow Y$ is called ϑ -Henstock primitive of f on E if $G(\emptyset) = \mathbf{0}$ (null vektor) and for every $A \in I(E)$, $A \neq \emptyset$, we have

$$G(A) = (H - \vartheta) \int_{cl(A)} g.$$

Theorem 1.2.4: Henstock's Lemma (Manuharawati, 2013: 22-26): *If $g: X \rightarrow Y$ ϑ -Henstock integrable on a cell $E \subset X$ with primitive G , then for any real number $\varepsilon > 0$ there exists a function $\delta: E \rightarrow \mathbb{R}^+$ such that for any Perron δ -fine partition $P = \{(A_x, x)\}$ and any $Q \subset P$ we have*

$$\|Q \sum [g(x)\vartheta(A_x) - G(A_x)]\| < 2\varepsilon.$$

2. RESULT AND DISCUSSION

Let X be a nondicret metric space, Y be Banach space, ϑ be a volume function on $E(X)$, and E be a cell in X . We start define an interval continuous function relative to volume function ϑ .

Definition 2.1: A function $G: I(E) \rightarrow Y$ is said to be v -continuous in an interval $A \in I(E)$ if for every real number $\varepsilon > 0$ there exist a real number $\gamma_A > 0$ such that for every $B, C \in I(E)$ with $B \subset A \subset C$, $v(A - B) < \gamma_A$, $v(C - A) < \gamma_A$ we have

$$\|G(A - B)\| < \varepsilon \text{ and } \|G(C - A)\| < \varepsilon.$$

A function G is said to be v -continuous on $I(E)$ if G is ϑ -continuous in every interval $A \in I(E)$.

Lemma 2.2: If an additive function $G: I(E) \rightarrow Y$ is v -continuous on $I(E)$ then for every real number $\varepsilon > 0$ there exist a real number $\gamma > 0$ such that for every interval $A \in I(E)$ with $v(A) < \gamma$ we have

$$\|G(A)\| < \varepsilon.$$

Proof: Let $\varepsilon > 0$ be given. Since G is v -continuous on $I(E)$, then G is v -continuous at $\{x\}$ for every $x \in E$. So for every $x \in E$ there exist a real number $\gamma_x > 0$ such that for every open interval $O_x \in I(E)$ with $x \in O_x$ and $\vartheta(O_x) < \gamma_x$ we

$$\|G(O_x)\| < \varepsilon.$$

If

$$\mathcal{L}_1 = \{O_x \in I(E): x \in \text{int}(E), x \in O_x, O_x \text{ is open, and } \vartheta(O_x) < \gamma_x\}$$

$$\mathcal{L}_2 = \{B_x \in S: x \in \partial(E), x \in B_x, B_x \text{ is open, and } \vartheta(B_x \cap E) < \gamma_x\},$$

then

$$\mathcal{L} = \mathcal{L}_1 \cup \mathcal{L}_2.$$

is a vitali cover of E . Since E is compact set, then we have $x_1, x_2, \dots, x_n \in E$ such that

$$\{O_{x_i} \in \mathcal{L}_1: 1 \leq i \leq l\} \cup \{O_{x_i} \in \mathcal{L}_1: 1 \leq i \leq l\} \{B_{x_i} \in \mathcal{L}_2: 1 \leq i \leq m\}$$

with $l + m = n$ is an open cover of E .

Further, put

$$\gamma = \min\{\gamma_{x_i}: 1 \leq i \leq n\}.$$

Since G is an additive function, then for every interval $A \in I(E)$ with $v(A) < \gamma$ we have

$$\|G(A)\| < \varepsilon. \quad \blacksquare$$

Definition 2.3: A function $G: I(E) \rightarrow Y$ is said to be strongly absolutely v -continuous on $F \subset E$ if for every real number $\varepsilon > 0$ there exist a real number $\gamma > 0$ such that for every partition $\{A_1, A_2, A_3, \dots, A_n\}$ in E with $\partial(A_i) \cap F \neq \emptyset$ and $\sum_{i=1}^n \vartheta(A_i) < \gamma$ we have

$$\sum_{i=1}^n \|G(A_i)\| < \varepsilon.$$

Theorem 2.4: Let a function $g: X \rightarrow Y$ be given. If $g \in H(E, \vartheta)$ with ϑ -Henstock primitive of g on E is G and g is bounded, then G is strongly absolutely v -continuous on E .

Proof: From boundedness of g on E , there exist a real number $M > 0$ such that for every $x \in E$ implies

$$\|g(x)\| \leq M.$$

Given any real number $\varepsilon > 0$. Based on density property on $\mathbb{R}$, there exists a real number $\eta > 0$ such that $\eta < \frac{\varepsilon}{2M}$. Take a collection $\{A_i, i = 1, 2, \dots, n\}$, $A_i \subset E$, with

$$\sum_{i=1}^n v(A_i) < \eta.$$

Since $g \in H(E, \vartheta)$, then by Henstock's Lemma and Theorem 1.2.3, there exist a function $\delta: E \rightarrow \mathbb{R}^+$ such that for every δ -fine partition $P_i\{(A_x, x)\}$ on A_i we have

$$\left\| P_i \sum [G(A_x - g(x)v(A_x))] \right\| < \frac{\varepsilon}{2^{i+1}}.$$

Since G is a v -Henstock primitive of g on E , then by Theorem 1.2.2 for every i we have

$$\begin{aligned} \|G(A_i)\| &= \|P_i \sum G(A_x)\| \\ &\leq \|P_i \sum G(A_x) - g(x)v(A_x)\| + \|P_i \sum g(x)v(A_x)\| \\ &< \frac{\varepsilon}{2^{i+1}} + M v(A_i). \end{aligned}$$

So we have

$$\begin{aligned} \sum_{i=1}^n \|G(A_i)\| &< \sum_{i=1}^n \left[\frac{\varepsilon}{2^{i+1}} + M v(A_i) \right] \\ &< \sum_{i=1}^n \frac{\varepsilon}{2^{i+1}} + M \sum_{i=1}^n v(A_i) \\ &< \frac{\varepsilon}{2} + M \frac{\varepsilon}{M} = \varepsilon. \end{aligned}$$

This mean that G is strongly absolutely ϑ -continuous on E . ■

Theorem 2.5: Let a function $g: X \rightarrow Y$ be given. If $g \in H(E, \vartheta)$ with ϑ -Henstock primitive of g on E is G , then G is ϑ -continuous on $I(E)$.

5

Proof: Let ε be areal positive number. If A is a cell in $I(E)$, then there exists open set O_A in X such that $A \subset O_A$ and $\vartheta(O_A - A) < \varepsilon$. If $\mathcal{L}$ be a collection of all O_A with $A \subset O_A$ for all $A \in I(E)$, then $\mathcal{L}$ is an open cover of E . Since E is a compact set, then there exists

$$\mathcal{M} = \{O_{A(i)}, i = 1, 2, \dots, n\} \subset \mathcal{L}$$

such that

$$E \subset \bigcup_{i=1}^n O_{A(i)}$$

If $F \in I(E)$, then by Lemma 2.2 and additivity of G , there exists areal number $\gamma_F > 0$ such that if $\vartheta(F) < \gamma_F$ we have

$$\|G(F)\| < \frac{\varepsilon}{n}.$$

Let

$$\gamma = \min\{\gamma_A : O_A \in \mathcal{M}\}$$

and $B, C, D \in I(E)$ be cells with

$$B \subset C \subset D, \vartheta(C - B) < \gamma \text{ and } \vartheta(D - C) < \gamma.$$

Then there exist two finite collection of nonoverlapping cells

$$\{C_i : i = 1, 2, \dots, l\} \subset I(E) \text{ and } \{D_i : i = 1, 2, \dots, s\} \subset I(E)$$

such that

$$cl(C - B) = \bigcup_{i=1}^l C_i \text{ and } cl(D - C) = \bigcup_{i=1}^s D_i.$$

Since $\vartheta(C - B) < \gamma$ and $\vartheta(D - C) < \gamma$, then for any i , $\vartheta(C_i) < \gamma$ and $\vartheta(D_i) < \gamma$. Note that, G is an additive function on $I(E)$. So, we have

$$\|G(C - B)\| = \sum_{i=1}^l \|G(C_i)\| < l \frac{\varepsilon}{n} \text{ and } \|G(D - C)\| = \sum_{i=1}^s \|G(D_i)\| < s \frac{\varepsilon}{n}.$$

So, G is ϑ -continuous on $I(E)$. ■

Theorem 2.6: Let a function $g: X \rightarrow Y$ be given. If $g \in H(E, \vartheta)$ with ϑ -Henstock primitive of g on E is G , then $D_v G(x)$ exists almost everywhere on E and $D_v G(x) = g(x)$ almost everywhere on E .

Proof: Given a real number $\varepsilon > 0$. Since $g \in H(E, \vartheta)$ with ϑ -Henstock primitive is G , then by Henstock's Lemma, there exist a function $\delta: E \rightarrow \mathbb{R}^+$ such that for every Perron δ -fine partition $P = \{(A_x, x)\}$ on E and every subpartition P_0 of P we have

$$P_0 \sum \|G(A_x) - g(x)v(A_x)\| < \varepsilon.$$

If

$$Z = \{x \in E : D_v G(x) \text{ exist or } D_v G(x) \text{ exist but } D_v G(x) \neq g(x)\},$$

then for every $x \in Z$ there exists a real number $\eta(x) > 0$ with property: for every real number $\gamma > 0$ there exists $B_x \in E(x, \gamma, \eta(x))$ such that

$$\|G(B_x) - g(x)v(B_x)\| > \eta(x)v(B_x).$$

For every $k \in \mathbb{N}$, put

$$Z_k = \{x \in Z : \eta(x) > \frac{1}{k}\}.$$

So we have

$$Z = \bigcup_{k \in \mathbb{N}} Z_k.$$

If

$$V = \{B_x : x \in Z_k, B_x \in E(x, \gamma, \eta(x)), \gamma < \delta(x)\},$$

then V is a Vitali cover for Z_k . So, there exists a nonoverlapping collection of cell

$$\{B_{x_i} : i = 1, 2, \dots, p\} \subset V$$

such that

$$\begin{aligned} v^*(Y_k) &\leq \sum_{i=1}^p v(B_{x_i}) + \varepsilon \\ &< k \sum_{i=1}^p \|G(B_{x_i}) - g(x)v(B_{x_i})\| + \varepsilon \\ &< k\varepsilon + \varepsilon. \end{aligned}$$

This mean that $v^*(Z_k) = 0$ and then $v(Z_k) = 0$ or $D_v G(x)$ exists almost everywhere on E . ■

If $A \in S$ then we define shape of A as follow

$$r(A) = \frac{\sup\{\vartheta(N(x, l)) : x \in A, N(x, l) \subset A\}}{\inf\{\vartheta(N(x, l)) : x \in A \subset N(x, l)\}}.$$

Further, if $E \in S, x \in E, \delta > 0, \rho > 0$ be given, a collection of all cell A with property $x \in A \subset N(x, \delta) \cap E$ and $r(A) > \rho$, denoted by $E(x, \delta, \rho)$.

Theorem 2.7: Given a function $g: X \rightarrow Y$. If $g \in H(E, \vartheta)$ with ϑ –Henstock primitive of g on E is G , then for every real number $\varepsilon > 0$ there exist a real number $\gamma > 0$ such that for every interval $A \in I(E)$ with $v(A) < \gamma$ we have

$$\|G(A)\| < \varepsilon.$$

Proof: Since G is a ϑ –Henstock primitive of g on E , then G is an additive function on $I(E)$. By Lemma 2.2 and Theorem 2.5, then for every real number $\varepsilon > 0$ there exist a real number $\gamma > 0$ such that for every interval $A \in I(E)$ with $v(A) < \gamma$ we have

$$\|G(A)\| < \varepsilon. \quad \blacksquare$$

3. CONCLUSION

If X is a locally compact metric space, Y is a Banach space, and G is a ϑ –Henstock primitive of a function $g: X \rightarrow Y$ on a cell E , then we have:

- 3.1 G is ϑ -continuous on $I(E)$,
- 3.2 G is strongly absolutely ϑ -continuous on $I(E)$ if g is a bounded on E ,
- 3.3 $D_\vartheta G(x)$ exists almost everywhere on E and $D_\vartheta G(x) = g(x)$ almost everywhere on E ,
- 3.4 for every real number $\varepsilon > 0$ there exist a real number $\gamma > 0$ such that for every interval $A \in I(E)$ with $v(A) < \gamma$ we have $\|G(A)\| < \varepsilon$.

REFERENCES

- Cao, S.S., 1992. The Henstock Integral for Banach Valued Function. *SEA Bull. Math.* 16,1: 35 – 40.
- Lee, P. Y., 1989, *Lanzhou Lectures on Hensock Integration*. World Scientific Publishing Co.Pte.Ltd. Singapore.
- Manuharawati. 2002. Integral Henstock di dalam Ruang Metrik Kompak Lokal. *Disertasi S3*. PPs-UGM, Yogyakarta.
- Manuharawati, 2013. Lemma Henstock Untuk Suatu Fungsi Bernilai Vektor di dalam Ruang Metrik Kompak Lokal. *Jurnal Sains & Matematika*. Vol. 2, No: 1. FMIPA Unesa Surabaya.
- Manuharawati dan Yunianti, D. N. 2013a. *Kriteria Cauchy dari suatu Fungsi Bernilai Vektor Pada Suatu Sel Di Dalam Ruang metrik Kompak lokal*. Prosiding Seminar Nasional Matematika dan Aplikasinya, 2013. Unair Surabaya.
- Manuharawati dan Yunianti, D. N. 2013. Integral ϑ –Henstock di dalam Ruang Metrik Kompak Lokal Dari Fungsi Bernilai Vektor. *Laporan Penelitian*. LPPM Unesa. Surabaya.



ORIGINALITY REPORT

7

%

SIMILARITY INDEX

4

%

INTERNET SOURCES

3

%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

f.library.uny.ac.id

Internet Source

4%

2

B BONGIORNO. "The Henstock-Kurzweil
Integral", Handbook of Measure Theory, 2002

Publication

1%

3

Nicolas Bourbaki. "Elements of Mathematics",
Springer Nature, 2004

Publication

1%

4

Iosif Petrakis. "Chapter 21 The Urysohn
Extension Theorem for Bishop Spaces",
Springer Science and Business Media LLC,
2016

Publication

<1%

5

Irving E. Segal, Ray A. Kunze. "Integrals and
Operators", Springer Nature, 1978

Publication

<1%

6

Veronique Fischer, Michael Ruzhansky.
"Quantization on Nilpotent Lie Groups", Springer

<1%

7

Bonotto, E.M.. "Limit sets and the Poincare-Bendixson Theorem in impulsive semidynamical systems", Journal of Differential Equations, 20080501

Publication

<1%

Exclude quotes

Off

Exclude matches

< 3 words

Exclude bibliography

On