

# How to Manipulate Axillary Bud Outgrowth in Tomato: Prospects for Practical Application?

*by*

---

**Submission date:** 17-Jun-2020 12:19PM (UTC+0700)

**Submission ID:** 1345220011

**File name:** 28\_How to manipulate Axillary bud outgrowth in tomato.pdf (217.34K)

**Word count:** 6634

**Character count:** 34959

---

**How to Manipulate Axillary Bud Outgrowth in Tomato:  
Prospects for Practical Application?**

Yuni Sri Rahayu  
Jurusan Biologi – FMIPA  
Universitas Negeri Surabaya

**Abstract**

Axillary bud outgrowth is controlled by apical dominance, which is regulated by the plant hormonal and plant nutritional status. In this study, we investigated perspectives for manipulation axillary bud growth by application of different N-forms to reduce manual work and thus production cost in greenhouse culture of tomato. Shoot growth was more affected than root growth when plants were supplied with  $\text{NH}_4^+$  as a sole nitrogen, leading to a decreased shoot/root ratio. Also axillary bud outgrowth could be significantly reduced under  $\text{NH}_4^+$  nutrition compared to  $\text{NO}_3^-$  nutrition. However, yield parameters such as number of flowers and trusses per plant, fruit per truss and total fruit yield were also decreased, and the incidence of blossom end rot in tomato fruit was increased by  $\text{NH}_4^+$  applications. However, the fruit size was not affected by the supply of different N forms.

**Key words:** ammonium, axillary bud, *Lycopersicon esculentum*, nitrate

### Introduction

In greenhouse tomato production, several manual manipulations are performed very intensively and constantly, such as removing the axillary shoots, “truss pruning”, removal of the oldest leaves, and keeping the main stem in optimal position. All these manual operations are necessary to maximise the quality and quantity of tomato fruits, but are also associated with high production costs. For tomato plants, a negative correlation between the weight of axillary shoots and yield has been demonstrated indicating that axillary shoots compete for nutrients and photosynthates with fruits. When de-shooting (removing of axillary shoots) was performed with low frequency, diameter and vigour of the stem were reduced, as well as the number of fruits per m<sup>2</sup> leading to a significantly lower yield (Navarrete and Jeannequin, 2000; Hartman, 1977). Thus, Navarrete and Jeannequin (2000) suggested an optimal de-shooting frequency between 7-14 days. However, the frequency of de-shooting also increases the mechanically induced stress (MIS), thus impairing yield (Biddington, 1986). Even simple shaking of tomato plants reduced the yield by 9% - 17% respectively when shaking was performed once or twice a day (Buitelaar, 1988). Also frequent mechanical measurements on leaves and fruits decreased height, leaf area and yield of sweet pepper plants (Kläring, 1999).

Lateral bud outgrowth is controlled by a process called apical dominance. In shoots, auxin derived from the shoot apex has been postulated to inhibit lateral bud growth either directly via entry into the lateral buds with subsequent repression of outgrowth or indirectly by involvement of i.e. activation of a second messenger, increased auxin-cytokinin ratio, secondary growth substances, nutrient diversion, auxin transport auto-inhibition, etc. (Bangerth *et al.*, 2000; Li and Bangerth, 1999; Tamas, 1995; Cline, 1994; Bangerth, 1994; Cline, 1991; Martin, 1987). In many species, removal of the shoot apex resulted in the

outgrowth of lateral branches (reviewed by Cline, 1991). Moreover, there is evidence that cytokinin is actively involved in the regulation of this process (Bangerth *et al.*, 2000; Chatfield *et al.*, 2000; Bangerth, 1994). Reduction of basipetal IAA transport leads to a strong increase in root derived cytokinin transport to the shoot. This increased delivery of cytokinin to the lateral buds promoted a strong increase in the production and export of IAA of lateral buds and their release from dormancy (Bangerth, 1994; Eklöf *et al.*, 1997; Kaminek *et al.*, 1997).

It is well known that nitrogen (N) deficiency leads to a drastic decrease of cytokinin levels in the plants (Wagner and Beck, 1993; Kuiper *et al.*, 1988). Moreover, there is evidence that the N form affects the endogenous levels of phytohormones such as cytokinin (Wang and Below, 1996; Gao *et al.*, 1992; Smiciklas and Below, 1992). Our previous studies with tobacco (Walch-Liu *et al.*, 2000) and tomato (Rahayu *et al.*, 2001) revealed that NO<sub>3</sub><sup>-</sup> supply compared to sole NH<sub>4</sub><sup>+</sup> nutrition increased the levels of zeatin+zeatin riboside (Z+ZR), the putative active forms of cytokinin, in the whole plant and at the same time particularly promoted shoot and leaf growth. As a consequence of these enhanced cytokinin concentrations, the release of lateral bud growth may be stimulated more in NO<sub>3</sub><sup>-</sup> than in NH<sub>4</sub><sup>+</sup> treated plants.

In this study, we focused on the effects of N form on axillary bud growth and yield quantity and quality of fruit in order to investigate options for minimization of cost-intensive manual removal of axillary shoots by manipulating plant growth and hormonal balances via fertilization management.

### Material and methods

#### *Plant cultivation*

Seeds of tomato (*Lycopersicon esculentum* L. cv. Moneymaker) were placed on filter paper which was soaked with 10 mM CaSO<sub>4</sub> solution and kept in the dark at a temperature of 25°C. After germination, the seedlings were transferred to a hydroponic culture system (4 plants per 3 l pots) (Walch-Liu *et al.*, 2000). After 4 weeks, each plant was transferred to 5 l pots, and after 8 weeks to 10 l pots until harvest time. Nitrogen was applied either as KNO<sub>3</sub> or (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> at a concentration of 2 mM N. For the NH<sub>4</sub><sup>+</sup> treatment, K<sub>2</sub>SO<sub>4</sub> was added to compensate for K applied at the KNO<sub>3</sub> variant. Depletion of N in the nutrient solution lower than 50% was avoided by checking the N concentration in the culture medium at least once a day using a RQflex reflectometer (Merck, Darmstadt, Germany) and by re-supply of appropriate N amounts. The nutrient solution was replaced completely every 2 d and pH was adjusted to 6.8 - 7.2 by addition of CaCO<sub>3</sub>. Plants were grown in a greenhouse with a temperature regime around 20-24°C at day time and 18-16°C at night time and additional light (300-400 μE m<sup>-2</sup> s<sup>-1</sup>) at a light/dark duration of 16/8 h.

The axillary shoot of the plants were removed by hand for all N treatments except the control plants every 3 days, after axillary shoot had reached about 2 cm length. Removal was performed for all axillary shoot including the ones located near the apex. The number of leaves at the main stem and the height of the main stem measured with a ruler was recorded at several defined times.

#### *Yield parameters*

After 14 weeks of growth, fresh weights of shoots, roots, axillary shoots of control plants and of single fruits were determined. The dry weight was obtained from oven-dried the

fresh material at 70°C until the weight reached a constant value. Yield parameters, such as the number of flowers, trusses per plant, and number of fruits per truss were obtained by manual counting. The incidence of blossom-end rot was determined by visual observation after harvest of the fruits.

***Analysis of Ca<sup>2+</sup> concentration in the fruits***

Fruits of the 1<sup>st</sup> – 4<sup>th</sup> fruit truss were separately analyzed, (3 fruits of different size per truss). Calcium concentration was determined according to Gericke and Kurmies (1952). About 250 mg dried fruit material was dry-ashed at 500°C for 5 hours and then dissolved in 1:30 diluted HCl solution. In the filtered solution Ca<sup>2+</sup> concentration was measured by flame photometry (ELEX 6361, Eppendorf, Germany).

***Analysis of organic acids in the fruits***

Oven-dried fruit material was homogenised with 5% (v/v) H<sub>3</sub>PO<sub>4</sub> (1ml acid per 10 mg dry weight). After centrifugation, the supernatant was diluted 10-fold with HPLC-elution buffer (18 mM KH<sub>2</sub>PO<sub>4</sub>; pH 2.1), and analysed by a reversed phase HPLC in the ion suppression mode. Separation was conducted on a 250 x 4 mm reversed phase column (GROM-SIL 120 ODS 5 ST, 5 µm particle size) equipped with a 20 x 4 mm Hypersil ODS guard column (Grom, Herrenberg, Germany). Sample solutions (20 µL) were injected onto the column and 18 mM KH<sub>2</sub>PO<sub>4</sub> adjusted to pH 2.1 with H<sub>3</sub>PO<sub>4</sub>, was used for isocratic elution with a flow rate of 0.5 mL min<sup>-1</sup> at 40°C and UV detection at 215 nm. Identification of organic acids was performed by comparing retention times and absorption spectra with those of known standards. The identity of citric, malic and oxalic acids was further confirmed by enzymatic tests (Boehringer, Mannheim, Germany) as described by Neumann and Römheld (1999).

---

**Results***Effect of N forms on vegetative plant growth*

Growth of tomato plants responded significantly to the supply of different N forms (Figs. 1A, C; 3). Shoot and root dry weight of  $\text{NH}_4^+$  supplied de-shooted plants decreased by 38% or 21%, respectively, compared to  $\text{NO}_3^-$  supplied plants. Control plants without de-shooting behaved similar with a reduction of shoot dry weight by 43% at  $\text{NH}_4^+$  supply compared to  $\text{NO}_3^-$  supplied plants (Fig. 1C). Although, root biomass was not influenced by the N form in control plants (Fig. 1C), the relation of shoot to root growth expressed as shoot/root dry weight ratio was significantly decreased at  $\text{NH}_4^+$  supply compared to  $\text{NO}_3^-$  fed plants (Figs. 1B, D). Interestingly, the supply with different N forms had no significant effect on leaf number of the main stem, although  $\text{NH}_4^+$ -supplied plants had a significantly smaller height (Fig. 2) and smaller leaf area than  $\text{NO}_3^-$  supplied plants.






Fig. 1. Effect of different N forms on dry weight of shoots (A, D), roots (B, E) shoot/root ratio (C, F) of de-shooted tomato plants and of control plants.



2 Fig. 2. Effect of different N forms on the number of leaves of the main stem (A) and height of tomato plants (B) at depending on plant age of de-shooted plants.

## How to Manipulate Axillary Bud Outgrowth in Tomato: Prospects for Practical Application?



1 Fig. 3. Effects of different N forms on shoot growth (A) and axillary bud growth (leaves on main stem removed) (B). Plants were 8 weeks after sowing.




Fig. 4. Effects of different N forms on total length and weight of axillary shoot of de-shooted plants (A, B) and control plants (C, D).

---

*Effect of N forms on axillary bud outgrowth*

The supply of different N forms affected axillary bud outgrowth. In control plants (without removing the axillary shoot) developed less axillary shoot under  $\text{NH}_4^+$  supply than  $\text{NO}_3^-$  treated plants (Figs. 3; 4A, B, C, D). Furthermore, compared to  $\text{NO}_3^-$  nutrition,  $\text{NH}_4^+$  supply decreased significantly the length and weight axillary shoots by 33% and 53%, respectively of de-shooted plants (Figs. 4 A, B), and by 53% and 72%, respectively of control plants (Figs. 4 C, D).

*Effects of N forms on yield*

Fruit setting processes such as number of flowers, number of fruits, number of trusses, total fruit weight and single fruit weight were affected by the N form (Fig. 5). Compared to  $\text{NO}_3^-$  nutrition,  $\text{NH}_4^+$  supply decreased the number of flowers, number of fruits and number of trusses by 40%, 21%, or 26%, respectively in de-shooted plants (Figs. 5 A, B, C), whereas mean fruit dry weight was not affected by the N form (Fig 5E). The final yield, particularly the total fruit weight was reduced by 48% in plants supplied with  $\text{NH}_4^+$  instead of  $\text{NO}_3^-$  (Fig. 5D). Some fruits of  $\text{NH}_4^+$ -fed plants exhibited symptoms of blossom end rot,  $\text{Ca}^{2+}$  related disorder, which was not observed in fruits of  $\text{NO}_3^-$  supplied plants (Tab. 1). A similar trend was also obtained for control plants, which were not de-shooted (Fig. 5). While number of flower and trusses were decreased by  $\text{NH}_4^+$  supply by 49% or 22%, respectively, (Figs. 5A, C), reduction of number of fruit was less pronounced (Figs. 5B). The total fruit yield and single fruit weight was not reduced by the  $\text{NH}_4^+$  supply (Fig. 5D, E).

**How to Manipulate Axillary Bud Outgrowth in Tomato:  
Prospects for Practical Application?**




Fig. 5. Effect of different N forms on number of flower (A, F), number of fruit (B, G), number of truss (C, H), total fruit weight (D, I), and single fruit weight (E, J) of de-shooted and control plants.

---

*Effect of N forms on Ca<sup>2+</sup> and organic acid concentrations in fruits*

---

Blossom end rot (BER), which was observed in fruits of NH<sub>4</sub><sup>+</sup> fed plants, is related to low Ca<sup>2+</sup> concentrations caused by the limited ability of the plant to regulate Ca<sup>2+</sup> distribution into growing regions and by the immobility of Ca<sup>2+</sup> in the phloem and in non-vascular tissues (Bangerth, 1973). Therefore, Ca<sup>2+</sup> concentrations were analysed in the fruits. However, the N form did not affect Ca<sup>2+</sup> concentrations in fruits, although the fruits of NH<sub>4</sub><sup>+</sup> supplied plants showed incidence of BER (Table 1). Another cause for Ca<sup>2+</sup> disorders could be its precipitation by chelators such as organic acids. However, NO<sub>3</sub><sup>-</sup> supplied plants had even higher organic acid concentrations in the fruits than NH<sub>4</sub><sup>+</sup> fed plants (Tab.1).

Table 1. Effect of different N forms on incidence of blossom end rot (BER) of fruits, calcium concentration and organic acid concentration of fruits of de-shooted plants. Plants were grown in nutrient solution containing 2 mM nitrogen either solely supplied as NO<sub>3</sub><sup>-</sup> or NH<sub>4</sub><sup>+</sup> and fruits were harvested 14 weeks after sowing.

Treatment	Ca conc. [mg (g DM) <sup>-1</sup> ]	Incidence of BER (%)	Organic acids [mg (g DW) <sup>-1</sup> ]					
			Malic	Shikimic	Citric	Fumaric	C-aconitic	aconitic
NO <sub>3</sub> <sup>-</sup>	1,40±0,05	0	20,5±1,1	46,8±1,4	22,7±1,2	17,8±2,4	6,4±1	13,7±1,5
NH <sub>4</sub> <sup>+</sup>	1,39±0,03	12,4±0,94	5,3±1,5	22,2±3,5	14,2±1,8	-	6,3±1,2	3,5±0,8

---

**Discussion***NH<sub>4</sub><sup>+</sup> supply decreased plant growth*

When plants were supplied with NH<sub>4</sub><sup>+</sup> compared to NO<sub>3</sub><sup>-</sup>, shoot biomass was drastically reduced. (Figs. 1; 3A). These results are consistent with our previous studies with *Nicotiana tabacum* L. (Walch-Liu *et al.*, 2000), and also with other reports of *Beta vulgaris* L., *Lycopersicon esculentum* Mill., *Phaseolus vulgaris* L., and *Zea mays* L. (Raab and Terry, 1994; Cramer and Lewis, 1993; Magalhães and Huber, 1989; Chaillou *et al.*, 1986). Reduced growth due to sole NH<sub>4</sub><sup>+</sup> supply was often related to decreased net photosynthesis caused by accumulation of NH<sub>4</sub><sup>+</sup> ions in the shoot (Claussen and Lenz, 1995; Goyal *et al.*, 1982; Puritch and Baker, 1967), acidification of the external media (Gerendas *et al.*, 1997; Magalhães and Wilcox, 1983), leading to NH<sub>4</sub><sup>+</sup> toxicity symptoms such as marginal necrosis and intervenal chlorosis on the leaves, wilting, stunted root growth or brownish roots (Goyal *et al.*, 1982; Maynard and Barker, 1969) or to a lack of osmolytes (i.e. nitrate, cations, and sucrose) involved in leaf expansion (Raab and Terry, 1995). However, previously we could demonstrate that these metabolic disorders related to NH<sub>4</sub><sup>+</sup> toxicity could be largely excluded by application of moderate NH<sub>4</sub><sup>+</sup> concentrations and by buffering the nutrient solution (Walch-Liu *et al.*, 2000). Ammonium-induced inhibition of leaf growth was associated with decreased zeatin+zeatin riboside (Z+ZR) concentrations in the xylem sap and in the plant tissue of tomato (Rahayu *et al.*, 2001) and of tobacco (Walch-Liu *et al.*, 2000). Cytokinins are involved in processes regulating leaf growth such as cell division and cell expansion (Francis and Sorrell, 2001; Ivanova and Rost, 1998).

Not only plant biomass, but also plant height was reduced by sole  $\text{NH}_4^+$  supply, whereas number of leaves on the main stem was not affected, indicating that the N form had no affect on general plant developmental (Fig. 2). Similar results were shown for tomato, tobacco, and sugar beets (Rahayu *et al.*, 2001; Walch-Liu *et al.*, 2000; Santamaria *et al.*, 1999; Raab and Terry, 1994).

***$\text{NH}_4^+$  supply repressed lateral bud outgrowth***

Ammonium nutrition significantly reduced axillary bud growth (Figs. 3, 4C, D) and also decrease the total length and weight of removed axillary shoot (Fig. 4 A, B). There is evidence that cytokinins are involved in the regulation of bud outgrowth. When cytokinins were applied to lateral buds, outgrowth was stimulated in intact plants (Mox, 1994). A study with *Arabidopsis* revealed that <sup>4</sup>basally applied cytokinin was found to release outgrowth of lateral shoot from inhibition by endogenous or exogenous apical auxin (Chatfield *et al.*, 2000). Our previous studies with tomato and tobacco demonstrated that  $\text{NH}_4^+$  supply obviously decreased cytokinin concentrations in the whole plant (Rahayu *et al.*, 2001; Walch-Liu *et al.*, 2000). Under these conditions low cytokinin levels may promote apical dominance leading to inhibition of lateral bud outgrowth.

However, the mechanisms controlling shoot branching and particularly the involvement of cytokinin are still obscure. Several models have been suggested to explain apical dominance involving an influence of auxin <sup>4</sup>on cytokinin export from or biosynthesis by the roots (Bangerth, 1994; <sup>4</sup>Eklof *et al.*, 1997; Kaminek *et al.*, 1997). According to <sup>4</sup>this studies, reduction of basipetal IAA transport leads to a strong increase in root derived cytokinin transport to the shoot. This increased delivery of cytokinin to the lateral buds promoted a strong increase in the production and export of IAA of lateral buds and their release from

dormancy. However, there are also evidence <sup>4</sup> that the endogenous levels of these two hormones are not always correlated with the degree of shoot branching, in rms mutants of pea e.g. suggests that additional signals could be involved in regulating apical dominance <sup>4</sup> (Beveridge *et al.*, 1997a; Beveridge *et al.*, 1997b; Beveridge *et al.*, 1994).

#### ***NH<sub>4</sub><sup>+</sup> supply reduced yield***

Ammonium nutrition decreased yield by reducing the number of flowers, the number of fruits and number of trusses (Figs. 5A, B, C, F, H) either due to increased flower and/or fruit abortion or due to low flower production. However, fruit size <sup>2</sup> was not affected by the N form (Fig. 5E, J). Thus, the N form influenced flowering and fruit setting processes but not fruit growth.

Reduced yield of tomato plants receiving NH<sub>4</sub><sup>+</sup>, instead of NO<sub>3</sub><sup>-</sup> was also reported by other workers (Siddiqi *et al.*, 2002; Sandoval-Villa *et al.*, 2001; 1999). It was reported for legume that abscission processes leading to flower abortion may begin very soon after anthesis (Dybing *et al.*, 1986; Brun and Betts, 1984; Huff and Dybing, 1980). This abortion may be caused by abscisic acid, a phytohormone which may be involved in promoting abscission of flowers and fruits (Taiz and Zeiger, 1998), since there is evidence that abscisic acid is increased by sole NH<sub>4</sub><sup>+</sup> nutrition (Peuke *et al.*, 1994). Moreover, there is also evidence that cytokinin application to selected flower of soybean increased flower production and reduced the rates of flower abortion indicating that cytokinin levels play a significant role in determining total yield in soybean (Nagel *et al.*, 2001; Reese *et al.*, 1995; Mosjidis *et al.*, 1993; Ghiasi *et al.*, 1987; Dybing *et al.*, 1986). Since in our previous studies with tomato (Rahayu *et al.*, 2001) and tobacco (Walch-Liu *et al.*, 2000) revealed

that  $\text{NH}_4^+$  supply decreased cytokinin levels and increased abscisic acid levels, it is likely that both flower initiation was decreased and flower abscission was increased.

Furthermore, there are reports that high cytokinin levels may increase fruit size during fruit development, because fruit size is a function of cell number rather than final cell size (Cowan *et al.*, 2001) and control of final fruit size also depends on an early and high rate of cell division (Higashi *et al.*, 1999; Bohner and Bangerth, 1988). However, in our study mean fruit size was not affected by lower cytokinin levels at  $\text{NH}_4^+$  supply (Figs. 6E, 6.7E).

When de-shooting of plants was omitted, flower initiation was decreased by  $\text{NH}_4^+$  supply similar to de-shooted plants. However the N form had no effect on number of fruit total yield and fruit size (Figs. 5G, I, J). Low frequency of de-shooting in tomato plants led to reduced vigour and diameter of the main stem, indicating a decreased vegetative growth compared to a de-shooting practiced every 7 days (Navarrete and Jeannequin, 2000; Navarrete *et al.*, 1997). As a consequence of this decreased vegetative growth, yield and in particular number of fruits was reduced. Similarly, Hartmann (1977) reported that yield of tomato plants was decreased with increasing weight of axillary shoots remaining at the plant. It was suggested that a part of the assimilates produced by the main stem is diverted away from the fruits to the axillary shoots.

***$\text{NH}_4^+$  supply reduced quality of fruit by increasing incident of blossom end rot (BER)***

Fruits of  $\text{NH}_4^+$ -fed plants were affected at a higher percentage by the incidence of blossom end rot (BER) than fruits of  $\text{NO}_3^-$  fed plants. There are many reports confirming that application of  $\text{NH}_4^+$  is associated with increased incidence of BER of tomato fruits (Siddiqi *et al.*, 2002; Vielemeyer and Weissert, 1990; DeKock *et al.*, 1982; Phil and

### How to Manipulate Axillary Bud Outgrowth in Tomato: Prospects for Practical Application?

Lambeth, 1980). Moreover, when supplying both  $\text{NO}_3^-$  and  $\text{NH}_4^+$ , incidence of blossom end rot increased when the percentage of  $\text{NH}_4^+$  reached 50% or more (Siddiqi *et al.*, 2002). Blossom end rot (BER) is associated with disintegration and increased ion permeability of cell membranes due to  $\text{Ca}^{2+}$  deficiency (Simon, 1978; Bangerth, 1973), as a consequence of the fundamental role of  $\text{Ca}^{2+}$  for membrane stability and cell integrity (Marschner, 1995). Blossom ends had always the lowest  $\text{Ca}^{2+}$  levels within the tomato fruit, and fruits affected by BER had lower  $\text{Ca}^{2+}$  levels than healthy fruits (Nonami *et al.*, 1995; Franco *et al.*, 1994; Adam and Ho, 1993; Ehret and Ho, 1986). However, in our study the N form did not affect  $\text{Ca}^{2+}$  concentrations in fruits (Tab. 1). These results are supported by our previous work with tobacco. Here we could not find a decrease of  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  concentration in expanding leaves of  $\text{NH}_4^+$ -fed plants compared to  $\text{NO}_3^-$  supply (Walch-Liu *et al.*, 2000). This is contradictions to many studies showing decreased cation concentrations ( $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ) in plant supplied with  $\text{NH}_4^+$  as nitrogen source (Siddiqi *et al.*, 2002; Jungk, 1977; Wilcox *et al.*, 1973; Barker *et al.*, 1967). Furthermore, in our study the incidence of blossom end rot could not be related to  $\text{Ca}^{2+}$  levels (Tab. 1). Similar results are reported (Nonami *et al.*, 1995; Barker and Ready, 1994; Pill and Lambeth, 1980) that this missing relationship of BER and  $\text{Ca}^{2+}$  concentrations may be explained by a rather local  $\text{Ca}^{2+}$  deficiency, which is not reflected in  $\text{Ca}^{2+}$  concentrations of the whole fruit, since the  $\text{Ca}^{2+}$  concentration in the distal placental tissue of the fruit during the rapid expansion period is critical for the induction of BER in tomato, rather than the  $\text{Ca}^{2+}$  concentration of the whole fruit at maturity (Adams and Ho, 1992; Ehret and Ho, 1986).

For distribution of  $\text{Ca}^{2+}$  in fruits and thus incidence of BER, the development of xylem vessels and network of vascular bundles in the distal fruit tissue during fruit development may play an important role (Belda *et al.*, 1996; Belda and Ho, 1993). There is evidence

that different levels of auxin or rates of auxin in transport together with other hormonal molecules such as cytokinins are important for vascular cell differentiation (Sachs, 2000; Aloni, 1993). Since in our previous studies  $\text{NH}_4^+$  supply lead to decreased cytokinin levels compared to  $\text{NO}_3^-$  nutrition, (Rahayu, *et al.*, 2001; Walch-Liu *et al.*, 2000), it is likely that vascular cell differentiation and xylem formation in  $\text{NH}_4^+$  fed plants is less pronounced than in  $\text{NO}_3^-$  supplied plants.

To explain the missing relationship of BER and  $\text{Ca}^{2+}$  concentrations it is also possibly that a part of  $\text{Ca}^{2+}$  was precipitated in the fruits by complexation with organic acids, e.g. citric acid. However, in our study  $\text{NH}_4^+$ -fed plants had less organic acids in the fruits than plants supplied with  $\text{NO}_3^-$  (Tab. 1), indicating that  $\text{Ca}^{2+}$  precipitation by organic acids might not be the cause for increased BER at  $\text{NH}_4^+$  supply. Organic acid concentrations were also decreased in roots by  $\text{NH}_4^+$  nutrition compared to  $\text{NO}_3^-$  supply (Marschner, 1995). The low levels of organic acid in  $\text{NH}_4^+$  nutrition plants should be also considered, since the taste and aroma of tomato fruits are affected by organic acids concentration in the fruits (Stevens, 1986).

### **Conclusion**

Our results showed that the specific application of different N forms can be used to manipulate the axillary bud outgrowth. Axillary shoot growth can be repressed by  $\text{NH}_4^+$  nutrition, however, yield of  $\text{NH}_4^+$ -fed plants was reduced and fruits showed higher incidence of BER. Thus, an optimum yield should be gained by switching the supply of  $\text{NH}_4^+$  to  $\text{NO}_3^-$  after initiation of the flowering process to decrease flower abscission and incident of BER or modify the  $\text{NH}_4^+/\text{NO}_3^-$  concerning the flowering process.

This type of using particular N-forms is valid or can be practised only under rather controlled growing conditions, e.g. if Nutrient Film Technique (NFT) is used. Further experiments should explore if altering  $\text{NH}_4^+/\text{NO}_3^-$  supply can reduce lateral shoot growth and possibly BER. With NFT control of the pH and N supply should be of no problem and could even be automated.

### References

- Adams P, Ho LC (1992) The susceptibility of modern tomato cultivars to blossom-end rot in relation to salinity. *Journal of Horticultural Science* 67: 827-839.
- Adams P, Ho LC (1993) Effects of environment on the uptake and distribution of calcium in tomato and on the incidence of blossom-end rot. *Plant and Soil* 154: 127-132.
- Aloni R (1993) The role of cytokinin in organised differentiation of vascular tissues. *Australian Journal of Plant Physiology* 20: 601-608.
- Bangerth F (1973) Investigation upon Ca related physiological disorders. *Phytopathologie Zeitschrift* 77: 20-37.
- Bangert F (1994) Response of cytokinin concentration in xylem exudates of bean (*Phaseolus vulgaris* L.) plants to decapitation and auxin treatment, and relationship to apical dominance. *Planta* 194: 439-442.
- Bangerth F, Li C, Gruber J (2000) Mutual interaction of auxin and cytokinins in regulating correlative dominance. *Plant Growth Regulation* 32: 205-217.
- Barker AV, Maynard DN, Lachmann WH (1967) Induction of tomato stem and leaf lesion and potassium deficiency by excessive ammonium nutrition. *Soil Science* 103: 319-327.
- Barker AV, Ready KM (1994) Ethylene evolution by tomatoes stressed by ammonium nutrition. *Journal of American Society Horticultural Science* 119: 706-710.
- Belda RM, Fenlon JS, Ho LC (1996) Salinity effects on the xylem vessels in tomato fruits among cultivars with different susceptibilities to blossom-end rot. *Journal of Horticultural Science* 71: 173-179.
- Belda RM, Ho LC (1993) Salinity effects on the network of vascular bundles during tomato fruit development. *Journal of Horticultural Science* 68: 557-564.
- Beveridge CA, Murfet IC, Kerhoas L, Sotta B, Miginiac E, Rameau C (1997a) The shoot controls zeatin riboside export from pea roots: evidence from the branching mutant rms4. *Plant Journal* 11: 339-345.
- Beveridge CA, Ross JJ, Murfet IC (1994) Branching mutant rms-2 in *Pisum sativum*. Grafting studies and endogenous indole-3-acetic acid levels. *Plant Physiology* 104: 953-959.
- Beveridge CA, Symons GM, Murfet IC, Ross JJ, Rameau C (1997b) The rms1 mutant of pea has elevated indole-3-acetic acid levels and reduced root-sap zeatin riboside content but increased branching controlled by graft-transmissible signals. *Plant Physiology* 115: 1251-1258.
- Biddington, NL (1986) The effects of mechanically-induced stress in plants – a review. *Plant Growth Regulation* 4: 103-123.

- Bohner J, Bangerth F (1988) Cell number, cell size and hormone levels in semi-isogenic mutant of *Lycopersicon pimpinellifolium* differing in fruit size. *Physiologia Plantarum* 72: 316-320.
- Brun WA, Betts KJ (1984) Source/Sink relations of abscising and non-abscising soybean flower. *Plant Physiology* 84: 187-191.
- Buitelaar K (1988) Are your tomatoes a little shaky? *Grower*, February 4, pp. 32-33.
- Chaillou S, Morot-Gaudry JF, Salsac L, Lesaint C, Jolivet E (1986) Compared effects of  $\text{NO}_3^-$  and  $\text{NH}_4^+$  on growth and metabolism of french bean. *Physiologie Vegetale* 24: 679-687.
- Chatfield SP, Stirnberg P, Forde BG, Leyser O (2000) The hormonal regulation of axillary bud growth in *Arabidopsis*. *The Plant Journal* 24(2): 159-169.
- Claussen W, Lenz F (1995) Effect of ammonium and nitrate on net photosynthesis, flower formation, growth and yield of eggplants (*Solanum melongena* L.). *Plant and Soil* 171: 267-274.
- Cline MG (1991) Apical dominance. *The Botanical Review* 57: 318-358.
- Cline MG (1994) The role of hormones in apical dominance. New approaches to an old problem in plant development. *Physiologia Plantarum* 90: 230-237.
- Cowan AK, Cripps RF, Richings EW, Taylor NJ (2001) Fruit size: Toward an understanding of the metabolic control of fruit growth using avocado as a model system. *Physiologia Plantarum* 111: 127-136.
- Cramer MD, Lewis OAM (1993) The influence of  $\text{NO}_3^-$  and  $\text{NH}_4^+$  nutrition on the carbon and nitrogen partitioning characteristics of wheat (*Triticum aestivum* L.) and maize (*Zea mays* L.) plants. *Plant and Soil* 154: 289-300.
- DeKock PC, Hall A, Boggie R, Inkson RHE (1982) The effect of water stress and form of nitrogen on the incidence of blossom-end rot in tomato. *Journal of the Science of Food and Agriculture* 33: 509-515.
- Dybing CD, Ghiasi H, Paech C (1986) Biochemical characterization of soybean ovary growth from anthesis to abscission of aborting ovaries. *Plant Physiology* 81: 1069-1074.
- Ehret DL, Ho LC (1986) Translocation of calcium in relation to tomato fruit growth. *Annals of Botany* 58: 679-688.
- Eklof S, Astot C, Blackwell J, Moritz T, Olsson O, Sandberg G (1997) Auxin-cytokinin interactions in wild-type and transgenic tobacco. *Plant and Cell Physiology* 38: 225-235.
- Francis D, Sorrell DA (2001) The interface between the cell cycle and plant growth regulators: a mini review. *Plant Growth Regulation* 33: 1-12.
- Franco JA, Banon S, Madrid R (1994) Effects of protein hydrolysate applied by fertigation on the effectiveness of calcium as a corrector of blossom-end rot in tomato cultivated under saline conditions. *Scientia Horticulturae* 57: 283-292.
- Gao YP, Motosugi H, Sugiura A (1992) Rootstock effects on growth and flowering in young apple trees grown with ammonium and nitrate nitrogen. *Journal of American Society Horticultural Science* 117: 446-452.
- Gerendas J, Zhu Z, Bendixen R, Ratcliffe RG, Sattelmacher B (1997) Physiological and biochemical processes related to ammonium toxicity in higher plants. *Zeitschrift für Pflanzenernährung und Bodenkunde* 160: 239-251.
- Gericke VS, Kurmies B (1952) Die Kolorimetrische Phosphorsäurebestimmung mit Ammonium-Vanadat-Molybdat und ihre Anwendung in der Pflanzenanalyse. *Zeitschrift für Pflanzenernährung und Bodenkunde* 59: 235-247.

**How to Manipulate Axillary Bud Outgrowth in Tomato:  
Prospects for Practical Application?**

- Ghiasi H, Dybing CD, Paech C (1987) Free amino acid content and metabolic activities of setting and aborting soybean ovaries. *Plant Physiology* 81: 1069-1074.
- Goyal SS, Huffaker RC, Lorenz OA (1982) Inhibitory effects of ammoniacal nitrogen on growth of radish plants. II. Investigation on the possible causes of ammonium toxicity to radish plants and its reversal by nitrate. *Journal of American Society of Horticultural Science* 107: 130-135.
- Hartmann HD (1977) Einfluß der Seitensprosse auf Wuchs- und Ertragsleistung von Tomatensorten. *Gartenbauwissenschaft* 42: 178-184.
- Higashi K, Hosoya K, Ezura H (1999) Histological analysis of fruit development between two melon (*Cucumis melo* L. *reticulatus*) genotypes setting a different size of fruit. *Journal of Experimental Botany* 50: 1593-1597.
- Huff A, Dybing CD (1980) Factors affecting shedding of flowers in soybean (*Glycine max* L.) Merril). *Journal of Experimental Botany* 31: 51-62.
- Ivanova M, Rost TL (1998) cytokinins and the plant cell cycle: Problems and pitfalls of proving their function. In: JA Bryant and D Chiatante (eds) *Plant Cell Proliferation and its Regulation in Growth and development*. John Wiley&Sons, New York, pp. 45-57.
- Jungk A (1977) Wirkung von Ammonium- und Nitrat-Stickstoff auf das Wachstum und die Zusammensetzung von Pflanzen. *Landwirtschaftliche Forschung*, Kongreßband, Teil II, 18-26.
- Kaminek M, Motyka V, Vankova R (1997) Regulation of cytokinin content in plant cells. *Physiology Plantarum* 101: 689-700.
- Kläring HP (1999) Effect of non-destructive mechanical measurements on plant growth: a study with sweet pepper (*Capsicum annum* L.). *Scientia Horticulturae* 81: 369-375.
- Kuiper D, Kuiper PJC, Lambert H, Schuit J, Staal M (1989) Cytokinin concentration in relation to mineral nutrition and benzyladenine treatment in *Plantago major* ssp. *Pleiosperma*. *Physiologia Plantarum* 75: 511-517.
- Li CJ, Bangerth F (1999) Autoinhibition of indoleacetic acid transport in the shoots of two-branched pea (*Pisum sativum*) plants and its relationship to correlative dominance. *Physiologia Plantarum* 106: 415-420.
- Magelhäs JR, Huber DM (1989) Ammonium assimilation in different plant species as affected by nitrogen form and pH control in solution culture. *Fertilizer Research* 21: 1-6.
- Magelhäs JR, Wilcox GF (1983) Tomato growth and nutrient uptake pattern as influenced by nitrogen form and light intensity. *Journal of Plant Nutrition* 6: 941-956.
- Martin, GC (1987) Apical dominance. *Horticulturae Science* 22: 824-833.
- Marschner H (1995a) Mineral nutrition of higher plants. 2<sup>nd</sup> edn. London: Academic Press.
- Maynard DN, Barker AV (1969) Studies on the tolerance of plants to ammonium nutrition. *Journal of American Society of Horticultural Science* 94: 235-239.
- Mosjidis CO'H, Peterson CM, Truelove B, Dute RR (1993) Stimulation of pod and ovule growth of soybean, *Glycine max* (L.) Merr., by 6-benzylaminopurine. *Annals of Botany* 71: 193-199.
- Mok MC (1994) Cytokinin and plant development-An overview. In: DWS Mok, and MC Mok (eds) *Cytokinins: Chemistry, Activity, and Function*. CRC Press, Boca Raton, FL., pp 155-166.
- Navarrete M, Jeannequin B (2000) Effect of frequency of axillary bud pruning on vegetative growth and fruit yield in greenhouse tomato crops. *Scientia Horticulturae* 86: 197-210.

- Navarete M, Jeannequin B, Sebillotte M (1997) Vigour of greenhouse tomato plants (*Lycopersicon esculentum* Mill.): Analysis of the criteria used by grower and search for objective criteria. *Journal of Horticultural Science* 72: 821-829.
- Neumann G, Römheld V (1999) Root excretion of carboxylic acids and protons in phosphorus-deficient plants. *Plant and Soil* 211: 121-130.
- Nonami H, Fukuyawa T, Yamamoto M, Yang L, Hashimoto Y (1995) Blossom-end rot of tomato plants may not be directly caused by calcium deficiency. *Acta Horticulturae* 396: 107-114.
- Peuke AD, Jeschke WD, Hartung W (1994) The uptake and flow of C, N and ions between roots and shoots in *Ricinus communis* L. III. Long-distance transport of abscisic acid depending on nitrogen nutrition and salt stress. *Journal of Experimental Botany* 45: 741-747.
- Phil WG, Lambeth VN (1980) Effects of soil water regime and nitrogen form on blossom-end rot, yield, water relations, and elemental composition of tomato. *Journal of America Society Horticultural Science* 105: 730-734.
- Puritch GS, Barker AV (1967) Structure and function of tomato leaf chloroplast during ammonium toxicity. *Plant Physiology* 42: 1229-1238.
- Raab TK, Terry N (1994) Nitrogen source regulation of growth and photosynthesis in *Beta vulgaris* L. *Plant Physiology* 105: 1159-1166.
- Raab TK, Terry N (1995) Carbon, nitrogen, and nutrient interaction in *Beta vulgaris* L. as influenced by nitrogen source,  $\text{NO}_3^-$  versus  $\text{NH}_4^+$ . *Plant Physiology* 107: 575-584.
- Rahayu YS, Walch-Liu P, Neumann G, von Wiren N, Römheld V, Bangerth F (2001) Effect of long-term and short-term supply of  $\text{NO}_3^-$  or  $\text{NH}_4^+$  on cytokinin levels and leaf expansion rate in tomato (*Lycopersicon esculentum* L. cv. MoneyMaker). In: Horst WJ ed. *Plant Nutrition, Food security and sustainability of agro-ecosystems through basic and applied research*. Kluwer Academic Publishers, Dordrecht, The Netherlands: 134-135.
- Reese RN, Dybing CD, White CA, Page SM, Larson JE (1995) Expression of vegetative storage protein in soybean raceme tissues in response to flower set. *Journal of Experimental Botany* 46: 957-964.
- Sachs T (2000) Integrating cellular and organismic aspects of vascular differentiation. *Plant and Cell Physiology* 41: 649-656.
- Sandoval-Villa M, Guertal EA, Wood CW (2001) Greenhouse tomato response to low ammonium-nitrogen concentrations and duration of ammonium-nitrogen supply. *Journal of Plant Nutrition* 24: 1787-1798.
- Sandoval-Villa M, Wood CW, Guertal EA (1999) Effects of nitrogen form, nighttime nutrient solution strength, and cultivar on greenhouse tomato production. *Journal of Plant Nutrition* 22: 1931-1945.
- Santamaria P, Elia A, Serio F, Gonnella M, Parente A (1999) Comparison between nitrate and ammonium nutrition in Fennel, Celery, and Swiss Chard. *Journal of Plant Nutrition* 22(7): 1091-1106.
- Siddiqi MY, Malhotra B, Min X, Glass ADM (2002) Effect of ammonium and inorganic carbon enrichment on growth and yield of a hydroponics tomato crop. *Journal of Plant Nutrition Soil Science* 165: 191-197.
- Simon EW (1978) The symptoms of calcium deficiency in plants. *New Phytology* 80: 1-15.
- Smiciklas KD, Below FE (1992) Role of cytokinin in enhanced productivity of maize supplied with  $\text{NH}_4^+$  and  $\text{NO}_3^-$ . *Plant and Soil* 142: 307-313.
- Stevens MA (1986) Inheritance of tomato fruit quality components. *Plant Breeding Reviews* Vol 4: 273-311.

**How to Manipulate Axillary Bud Outgrowth in Tomato:  
Prospects for Practical Application?**

- 
- Tamas IA (1995) Hormonal regulation of apical dominance. IN: Davies PJ, ed. Plant hormones, physiology, biochemistry and molecular biology. 2<sup>nd</sup> edn. Boston: Kluwer Academic Publishers, 572-597.
- Taiz L, Zeiger E (1998) Plant Physiology. Sunderland: Sinauer Associates, Inc.
- Vielemeyer H-P, Weissert P (1990) Diagnose des Ca-Ernährungszustandes und der Blütenendfäule bei Buschtomate. *Gartenbauwissenschaft* 55: 209-212.
- Wagner BM, Beck E (1993) Cytokinins in the perennial herb *Urtica dioica* L. as influenced by its nitrogen status. *Planta* 190: 511-518.
- Walch-Liu P, Neumann G, Bangerth F, Engels C (2000) Rapid effects of nitrogen form on leaf morfogenesis in tobacco. *Journal of Experimental Botany* 51: 227-237.
- Wang X, Below FE (1996) Cytokinins in enhanced growth and tillering of wheat induced by mixed nitrogen source. *Crop Science* 36: 121-126.
- Wilcox GE, Hoff JE, Jones CE (1973) Ammonium reduction of calcium and magnesium concentration of tomato and sweet corn leaf tissue and influence on the incidence of blossom end rot of tomato fruit. *Journal of American Society of Horticultural Science* 98: 86-89.

# How to Manipulate Axillary Bud Outgrowth in Tomato: Prospects for Practical Application?

## ORIGINALITY REPORT

13%

SIMILARITY INDEX

6%

INTERNET SOURCES

15%

PUBLICATIONS

4%

STUDENT PAPERS

## PRIMARY SOURCES

- 1** "Plant Nutrition", Springer Science and Business Media LLC, 2001 **4%**  
Publication
- 2** P. Walch-Liu. "Rapid effects of nitrogen form on leaf morphogenesis in tobacco", Journal of Experimental Botany, 02/02/2000 **3%**  
Publication
- 3** J. Shen, H. Li, G. Neumann, F. Zhang. "Nutrient uptake, cluster root formation and exudation of protons and citrate in *Lupinus albus* as affected by localized supply of phosphorus in a split-root system", Plant Science, 2005 **2%**  
Publication
- 4** [onlinelibrary.wiley.com](http://onlinelibrary.wiley.com) **1%**  
Internet Source
- 5** Y. S. Rahayu. "Root-derived cytokinins as long-distance signals for NO<sub>3</sub><sup>-</sup>-induced stimulation of leaf growth", Journal of Experimental Botany, 02/21/2005 **1%**  
Publication

---

6

M.C. Saure. "Blossom-end rot of tomato (Lycopersicon esculentum Mill.) — a calcium- or a stress-related disorder?", *Scientia Horticulturae*, 2001

Publication

1%

---

7

M Navarrete, B Jeannequin. "Effect of frequency of axillary bud pruning on vegetative growth and fruit yield in greenhouse tomato crops", *Scientia Horticulturae*, 2000

Publication

1%

---

---

Exclude quotes      On

Exclude bibliography      On

Exclude matches      < 1%