

# 31. Determination of Chromium Content i Various Foodstuffs.pdf

*by*

---

**Submission date:** 06-Aug-2021 02:12PM (UTC+0700)

**Submission ID:** 1628351020

**File name:** 31. Determination of Chromium Content i Various Foodstuffs.pdf (284.13K)

**Word count:** 2774

**Character count:** 14977

## Determination of chromium content in various foodstuffs

Rudiana Agustini and I Gusti Made Sanjaya  
Department of Chemistry, Faculty of Mathematics and science, Surabaya State  
University, Indonesia  
E-mail is [rudianaagustini@unesa.ac.id](mailto:rudianaagustini@unesa.ac.id)

**Abstract.** Research has been conducted to identify the content of chromium in foodstuffs, including broccoli, celery, snake grass, green mustard greens, tomatoes, carrots, beans, green beans, cauliflower, chicory, cassava, potatoes, black rice, white rice, brown rice, black sticky rice, yeast (bakery yeast) and yeast extract commercial. The determination of chromium was carried out using voltammetry method, which the results were processed using OriginPro program. The data were then analyzed descriptively. The results showed that the foodstuffs containing the highest  $\text{Cr}^{3+}$  is cauliflower that is  $280 \times 10^{-4}\%$ , but the  $\text{Cr}^{6+}$  content is  $83 \times 10^{-4}\%$ . The foodstuffs containing  $\text{Cr}^{6+}$  in small quantities are cassava, nuts and commercial yeast, which is about  $8 \times 10^{-4}\%$  to  $9 \times 10^{-4}\%$ , but  $\text{Cr}^{3+}$  content is also not very high that is about  $18 \times 10^{-4}\%$  to  $21 \times 10^{-4}\%$ .

*Keywords:* Chromium content, Foodstuffs, Chromium in foodstuffs, chromium determination

### 12 1. INTRODUCTION

Chromium is one of the seven most abundant elements in the Earth's crust and at low concentrations is an essential element for living organisms naturally. Chromium is present in various forms of compounds. In addition to being Cr metal, the chromium is found as  $\text{Cr}^{2+}$ ,  $\text{Cr}^{3+}$  (trivalent chromium), and  $\text{Cr}^{6+}$  or chromium hexavalent.  $\text{Cr}^{3+}$  is known to be non-toxic, while  $\text{Cr}^{6+}$  is toxic to the human body. Cr chemicals are persistent, bio-accumulative, toxic and not readily degradable in the environment, thus accumulating in the human body through the food chain. Cr from for example the soil environment is entering the food chain through the plant. According to the World Health Organization,  $\text{Cr}^{6+}$  concentrations in soil should not be more than 0.05 mg/L or 50 ppb [1].  $\text{Cr}^{6+}$  compounds also have genotoxic, mutagenic, and carcinogenic properties [2]. The toxicity brought by this metal can harm vital organs such as liver, kidneys, cause lung cancer, acute poisoning, chronic, irritation to the respiratory system, and irritation to the human skin [3].

Various natural ingredients including foodstuffs are known to contain chromium, both  $\text{Cr}^{3+}$  and  $\text{Cr}^{6+}$  ions. Several studies have revealed that consumption of chromium can decrease type 2 diabetes mellitus (DM). Chromium is an essential mineral that the body needs for carbohydrate and fat metabolism [4]. Diabetes mellitus type 2 (T2DM) is the most common form of diabetes found worldwide [5] [6] [7]. This is characterized by abnormalities in pancreatic insulin secretion or actions that cause hyperglycemia due to impaired metabolism of carbohydrates, fats and lipids [5][7]. T2DM prevalence worldwide is increasing and more than 366 million people are expected to be affected by 2030 [5][7]. T2DM is continuing to be a public health concern, and many people are using alternative medicine using chromium. Chromium is a common supplement used by many T2DM patients for the purpose of improving glucose regulation and in 2002 sales of chromium supplements were estimated at \$85 million [8].

According to the National Institute of Health: Dietary Supplement Office, adequate intake of Cr for men and women is 35 and 25  $\mu\text{g}/\text{day}$ , respectively [5][9][10]. Chromium chloride is a natural

trivalent chromium variety found in common foodstuffs sources such as: whole grains, broccoli, mushrooms and green beans. In contrast, Cr picolinate is the synthetic family of Cr chloride. Additional forms of Cr supplements can also come from Cr's yeast and brewer's yeast. Chromium is an important micronutrient associated with the regulation of many processes in the human body including glucose homeostasis. Chromium helps regulate glucose homeostasis by activating insulin receptors through chromodulin oligopeptide thus increasing insulin signal transduction and sensitivity. Cr deficiency can lead to glucose intolerance, high circulating insulin, hyperglycemia at the time of fasting, and even disruption of growth [5][11].

Chromium can be obtained from foodstuffs and is available in very small quantities (1-2 micrograms or less). Chromium in the foodstuffs has a form of chromium (III) (indicating the amount of oxidation). Trivalent chromium or chromium (III) or  $\text{Cr}^{3+}$  is the most stable chromium and most secure, including one of the least toxic.  $\text{Cr}^{3+}$  is relatively harmless and has a role in the body's metabolism, while  $\text{Cr}^{6+}$  has a potential of 100-1000 times more toxic than  $\text{Cr}^{3+}$  because it has a high oxidation potential.

The following is a list of foods rich in chromium: broccoli (18.55 mcg/1 cup), barley (8.16 mcg/0.33 cup), oats (5.38 mcg/0.25 cup), beans (2.04 mcg/1 cup), tomatoes (1.26 mcg/1 cup). Chromium has proven its involvement in the mechanism of treatment of type 2 diabetes in several ways, including by increasing glucose tolerance. Chromium supplements increase glucose tolerance in humans with type 2 of diabetes [12]. Cr supplementation with brewer's yeast may provide a marginal benefit in lowering blood glucose levels in patients with T2DM compared with placebo but no effect on glycated hemoglobin [5].

It is generally accepted that chromium is an essential element for humans. Chromium deficiency has been described in both humans and animals, but a clear quantitative definition of the daily requirement of chromium in human nutrition has not been arrived at estimates that the daily minimum population mean intake likely to meet normal requirements for chromium might be approximately 33  $\mu\text{g}$ /person [1]. Results of panel related to the Food Supplement and Nutrition Source,  $\text{Cr}^{3+}$  added to food and food aimed at the general population (including dietary supplements), concluded that after oral administration, trivalent chromium is poorly absorbed. The result of in vitro bacterial mutagenic test consistently is negative. The panel concluded that in very large quantities, certain trivalent chromium compounds have been shown to be cytotoxic and cause chromosomal damage. The panel also evaluated long-term toxicity and carcinogenicity data for  $\text{Cr}^{3+}$ . Based on the facts, it is known: 1) a maximum intake level up to 250  $\mu\text{g}$ /day for additional intake, 2) that in vitro, at high levels of concentration,  $\text{Cr}^{3+}$  can cause DNA damage, 3) that DNA damage not reflected in the in vivo genotoxicity test, 4) that  $\text{Cr}^{3+}$  is not carcinogenic, 5) that it is safe for a daily intake of 250  $\mu\text{g}$ /day, equivalent to 4.1  $\mu\text{g}/\text{kg}$  body weight/day for people 60 kg [13]. Although chromium is an essential trace element for humans because it helps us to use glucose. However, it is poisonous in excess.

Daily chromium intake according to US dietary guidelines is 50-200 mg for adults, 30-35 mg for adult males and 20-25 mg for adult women [14][1]. About 2% of  $\text{Cr}^{3+}$  or trivalent chromium can be absorbed and the rest is excreted in the stool. Amino acids, vitamin C and niacin can increase the absorption of chromium by channels of the intestine. These minerals further accumulate in the liver, bone, and spleen. Trivalent chromium is found in a variety of foodstuffs, including wheat products, processed meats, cereals, coffee, beans, green beans, broccoli, spices, and some brands of wine and beer. Most fruits and vegetables and dairy products contain only low amounts.

## 2. METHODOLOGY

### 2.1 Preparation of materials

Foodstuffs identified about their chromium content are broccoli, celery, snake grass, green mustard, lettuce, tomatoes, carrots, beans, green beans, cauliflower, chicory, cassava, and potatoes, black sticky rice, white rice, brown rice, black rice, yeast (bakery yeast) and yeast commercial extract. The foodstuffs cut into small pieces, then weighed each 5 grams. The next step of each

foodstuff is heated in a furnace at a temperature of 700°C for 7 hours to obtain ash. The obtained ash is dissolved in 1 ml of the concentrated HNO<sub>3</sub> and 1 ml of the concentrated HCl. Each solution is ready to determine its chromium content.

## 2.2 Determination of chromium content

Determination of chromium content is done by using voltammetry method. Instrument used is Voltammeter. The chromium to be determined is Cr<sup>3+</sup> and Cr<sup>6+</sup>. The first step taken on the determination of this chromium content is to make a standard solution. The standard solution used was prepared by dissolving K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> and CrCl<sub>3</sub>·6H<sub>2</sub>O in variations of 5, 10, 20, 40 and 80 ppm. Measurements were made using Silica Carbon Electrode with scan rate of 50mV/sec [15]. Measurement of standard solution using voltammeter produces voltammogram, which then processed using Origin-Pro program and made standard curve between concentration versus current, so obtained linear regression equation  $y = a + bx$ . Then measured each sample solution or foodstuff, the result is incorporated into the equation.

## 3. RESULT AND DISCUSSION

Figures-1 and -2 show the standard curves of Cr<sup>3+</sup> and Cr<sup>6+</sup> in variations of concentrations of 5, 10, 20, 40, and 80 ppm. The regression equation obtained from the standard measurement of Cr<sup>3+</sup> is  $Y = 3.370798x + 0.004378$  and the regression equation obtained from the standard measurement of Cr<sup>6+</sup> is  $Y = 3.139306x + 0.008648$ .




Figure-1 Standard Curve of Cr<sup>3+</sup>




Figure-2 Standard Curve of Cr<sup>6+</sup>

By entering the data from the current strength obtained from each sample or foodstuff tested using voltammeter into the equation  $Y = 3.370798x + 0.004378$  for the determination of Cr<sup>3+</sup> and  $Y = 3.139306x + 0.008648$  for the determination of Cr<sup>6+</sup> then the data obtained as shown in Table-1 and -2.

Table 1 Content of Cr<sup>3+</sup> Determined from Various Foodstuffs.

Groups	Name of foodstuffs	Cr <sup>3+</sup> (10 <sup>-4</sup> %)	
Vegetable group	Broccoli	Brassica oleraceavaritalica	116
	Celery	Apiumgraveolens L	96
	Snake grass	<i>Clinacanthusnutanslindau</i>	82
	green mustard	Brassica rapavarparachinensis	96
	lettuce	Lactuca sativa	136
	Cauliflower	Brassica oleraceavarBotrytis	280
	Chinese cabbage	Brassica juncea	176
Group of fruit	Tomatoes	Licopersicumesculentum	141
	Carrots	Daucuscarota	110
	Long beans	Vignasinensis	20
	Chili	Capsicum frutescens	70
Group of rice /cereals	black rice	Oryza sativa L	20
	Black sticky rice	Oryza sativa varglutinosa	21
	Brown rice	Oryzapunctata	19
	Green beans	Oryza sativa	18
	White rice	Phaseolusaureus	20
Group of tubers	Red beans	Vignaangularis	18
	Potatoes	Solanumtuberosum L	83
	Cassava	Manihotutilissima	41
Group of yeast	Yeast (bakery's yeast)	Saccharomyces cerevisiae	16
	Yeast extract commercial sample	Saccharomyces cerevisiae	18

Table 2 Content of Cr<sup>6+</sup> Determined from Various Foodstuffs.

Groups	Name of foodstuffs	Cr <sup>6+</sup> (10 <sup>-4</sup> %)	
Vegetable group	Broccoli	Brassica oleraceavaritalica	54
	Celery	Apiumgraveolens L	45
	Snake grass	<i>Clinacanthusnutanslindau</i>	39
	green mustard	Brassica rapavarparachinensis	45
	lettuce	Lactuca sativa	64
	Cauliflower	Brassica oleraceavarBotrytis	83
	Chinese cabbage	Brassica juncea	39
Group of fruit	Tomatoes	Licopersicumesculentum	66
	Carrots	Daucuscarota	132
	Long beans	Vignasinensis	52
	Chili	Capsicum frutescens	33
Group of rice /cereals	black rice	Oryza sativa L	9
	Black sticky rice	Oryza sativa varglutinosa	10
	Brown rice	Oryzapunctata	9
	Green beans	Oryza sativa	9

	White rice	Phaseolusaureus	9
	Red beans	Vignaangularis	9
Group of tubers	Potatoes	Solanumtuberosum L	19
	Cassava	Manihotutilissima	8
Group of yeast	Yeast (bakery's yeast)	Saccharomyces cerevisiae	8
	Yeast extract commercial sample	Saccharomyces cerevisiae	9

Based on the data in Tables-1, and Table-2, it appears that for the vegetable group  $Cr^{3+}$  content ranges from  $82 \times 10^{-4}\%$  to  $280 \times 10^{-4}\%$ . Cauliflower has the highest  $Cr^{3+}$  content, which is  $280 \times 10^{-4}\%$ . While snake grass, has the smallest content, which is  $82 \times 10^{-4}\%$ . Snake grass is a local crop that is believed to be useful for treating diabetes mellitus. Groups of fruits contain  $Cr^{3+}$  ranging from  $20 \times 10^{-4}\%$  to  $141 \times 10^{-4}\%$ . Tomato fruit has the highest  $Cr^{3+}$  content, which is  $141 \times 10^{-4}\%$ . While the long bean, has the smallest content, which is  $20 \times 10^{-4}\%$ . The black rice has the highest  $Cr^{3+}$  content, which is  $21 \times 10^{-4}\%$ . While green beans and red beans, has the smallest content, which is  $18 \times 10^{-4}\%$ . Tuber groups contain  $Cr^{3+}$  ranging from  $41 \times 10^{-4}\%$  to  $83 \times 10^{-4}\%$ . Potatoes have the highest  $Cr^{3+}$  content, which is  $83 \times 10^{-4}\%$ . While cassava has the smallest content, which amount to  $41 \times 10^{-4}\%$ . The yeast group has  $Cr^{3+}$  content ranging from  $16 \times 10^{-4}\%$  to  $18 \times 10^{-4}\%$ . The commercial yeast extract which is thought to be made of brewery yeast has the highest  $Cr^{3+}$  content, which is  $18 \times 10^{-4}\%$ . While yeast (bakery yeast/yeast bread), has the smallest content, which is  $18 \times 10^{-4}\%$ . Based on the findings in this study, it can be concluded that the vegetable group has the highest  $Cr^{3+}$  levels among the foodstuffs tested, especially cauliflower.

Based on the data in Table -2, it can be seen that for the vegetable group  $Cr^{6+}$  content ranges from  $39 \times 10^{-4}\%$  to  $83 \times 10^{-4}\%$ . Cauliflower has the highest  $Cr^{6+}$  content, which is  $83 \times 10^{-4}\%$  while the chicory and snake grass have the smallest content, which is  $39 \times 10^{-4}\%$ . The fruit group contained  $Cr^{6+}$  ranging from  $33 \times 10^{-4}\%$  to  $132 \times 10^{-4}\%$ . Carrot fruit has the highest  $Cr^{6+}$  content, which is  $132 \times 10^{-4}\%$  while chili has the smallest content, which is equal to  $33 \times 10^{-4}\%$ . The rice/cereals group contained  $Cr^{6+}$  ranging from  $9 \times 10^{-4}\%$  to  $10 \times 10^{-4}\%$ . Black rice has the highest  $Cr^{6+}$  content, which is  $10 \times 10^{-4}\%$  whereas white rice, brown rice, black sticky rice, green beans and red beans contain  $9 \times 10^{-4}\%$ . Tuber groups contain  $Cr^{6+}$  ranging from  $8 \times 10^{-4}\%$  to  $19 \times 10^{-4}\%$ . Potatoes have the highest  $Cr^{6+}$  content, which is  $8 \times 10^{-4}\%$  while cassava has a content of  $19 \times 10^{-4}\%$ . The yeast group has a  $Cr^{6+}$  content ranging from  $8 \times 10^{-4}\%$  to  $9 \times 10^{-4}\%$ . Commercial yeast extract has the highest  $Cr^{6+}$  content, which is  $9 \times 10^{-4}\%$  while yeast (bakery yeast/baker yeast) is  $8 \times 10^{-4}\%$ . Based on the findings in this study, it can be concluded that carrots have the highest  $Cr^{6+}$  content among the foodstuffs tested.

#### 4. CONCLUSION

The results showed that foodstuffs containing the highest  $Cr^{3+}$  were cauliflower having  $280 \times 10^{-4}\%$ , but  $Cr^{6+}$  was  $83 \times 10^{-4}\%$ . Foodstuffs that have a small  $Cr^{6+}$  content are cassava, nuts and commercial yeast, which range from  $8 \times 10^{-4}\%$  to  $9 \times 10^{-4}\%$  but  $Cr^{3+}$  content is not very high, ranging from  $18 \times 10^{-4}\%$  to  $20 \times 10^{-4}\%$ .

#### 5. REFERENCES

- [1] World Health Organization, Trace elements in human nutrition and health. Geneva, 1996..
- [2] Ertürün KEH, Yilmaz M, Kilic E., Construction of an anion selective electrode: Dicromate-selective electrode. Sensors and Actuators B, 127, 2007, pp. 497-504
- [3] Slamet et al.. Pengolahan limbah organik (fenol) dan logamberat (Cr VI atau Pt VI) secara simultan dengan fotokatalis TiO<sub>2</sub>, ZNO-TiO<sub>2</sub>, dan CDS-TiO<sub>2</sub>. Makasra Teknol, 9, 2005, pp. 66-71.
- [4] Wilson BE, Gondy A., Effects of chromium supplementation on fasting insulin levels and

11

lipid parameters in healthy, non-obese young subjects. *Diabetes Research and Clinical Practice*, 1995.

- [5] Yin, R.V. and Phung, O.J., Effect of chromium supplementation on glycated hemoglobin and fasting plasma glucose in patients with diabetes mellitus. *Nutrition Journal* 14, 14, 2015.
- [6] Yeghiazaryan K, Schild HH, Golubnitschaja O., Chromium-picolinate therapy in diabetes care: individual outcomes require new guidelines and navigation by predictive diagnostics. *Infect Disord Drug Targets* vol.12, issue 5, 2012, pp. 332–9.
- [7] Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 27, 2004, pp. 1047–53.
- [8] *Nutrition Business Journal*, NBJ's Supplement Business Report. San Diego, CA: Penton Media Inc., 2003.
- [9] Rabinovitz H, Friedensohn A, Leibovitz A, Gabay G, Rocas C, Habet B., Effect of chromium supplementation on blood glucose and lipid levels in type 2 diabetes mellitus elderly patients. *Int J VitamNutr Res.*, Vol. 74, Issue 3, 2004, pp. 178–82.
- [10] National Institute of Health, Chromium: Dietary supplement fact sheet, 2013 .
- [11] Vincent J.B., Elucidating a biological role for chromium at a molecular level, *AccChem Res.*, Vol. 33, Issue 7, 2000, pp. 503–10.
- [12] Evans GW., The effect of chromium picolinate on insulin controlled parameters in humans. *Int J Biosocial Med Res.*, 11, 1989, pp. 163-80.
- [13] EFSA Panel, Scientific Opinion on the safety of trivalent chromium as a nutrient added for nutritional purposes to foodstuffs for particular nutritional uses and foods intended for the general population (including food supplements. EFSA Panel on Food Additives and Nutrient Sources, DOI: 10.2903/j.efsa.2010.1882, 2010.
- [14] Trumbo, P. et al., Dietary reference intakes : Vitamin A, Vitamin K, Arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium and zinc. *Journal of the American Dietetic Association*, 2001.
- [15] Yulianto, E. and Setiarso, P., Making of Chitosan Modified Carbon Paste Electrode for Cr (VI) Metallic Analysis With Ion Interfering Fe (II) and Zn (II) By Cyclic Stripping Voltammetry. Surabaya: Chemistry , Surabaya State University, 2014.

# 31. Determination of Chromium Content i Various Foodstuffs.pdf

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

9%

PUBLICATIONS

8%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://efsa.onlinelibrary.wiley.com">efsa.onlinelibrary.wiley.com</a> Internet Source	4%
2	Submitted to Udayana University Student Paper	3%
3	<a href="http://www.euro.who.int">www.euro.who.int</a> Internet Source	2%
4	<a href="https://en.wikipedia.org">en.wikipedia.org</a> Internet Source	2%
5	Xiao, T.. "Environmental concerns related to high thallium levels in soils and thallium uptake by plants in southwest Guizhou, China", Science of the Total Environment, The, 20040105 Publication	1%
6	Submitted to K12 Incorporated Student Paper	1%
7	R Agustini, I G M Sanjaya, Lupita. "Characterization of yeast hydrolysate enzymatic (yhe) from yeast fermented in the	1%

variation of rice flour", Journal of Physics:  
Conference Series, 2019

Publication

---

8	Submitted to Endeavour College of Natural Health Student Paper	1 %
9	costco.aisle7.net Internet Source	1 %
10	Submitted to Catholic High School Student Paper	1 %
11	onlinelibrary.wiley.com Internet Source	1 %
12	S ZAFRASTONE. "Benefits of chromium(III) complexes in animal and human health", The Nutritional Biochemistry of Chromium (III), 2007 Publication	1 %
13	suttermedicalcenter.org Internet Source	<1 %
14	Submitted to Edison State College Student Paper	<1 %
15	Nelly Wahyuni, Imelda H. Silalahi, Dina Angelina. "ISOTERM ADSORPSI FENOL OLEH LEMPUNG ALAM", Jurnal Teknologi Lingkungan Lahan Basah, 2019 Publication	<1 %

---

16

Putu Wijaya Sunu, I Made Rasta, Daud Simon Anakottapary, I Made Suarta, I D M Cipta Santosa. "Capillary Tube and Thermostatic Expansion Valve Comparative Analysis in Water Chiller Air Conditioning", Journal of Physics: Conference Series, 2018

Publication

<1 %

17

docplayer.net

Internet Source

<1 %

18

Amir Khodavirdipour, Fatemeh Haddadi, Shiva Keshavarzi. "Chromium Supplementation; Negotiation with Diabetes Mellitus, Hyperlipidemia and Depression", Journal of Diabetes & Metabolic Disorders, 2020

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On