

The role of laserpuncture exposure on gonad maturation mechanism of catfish (*Clarias* sp.) through Ca^{2+} , PKC and GABA neurotransmitter

by Dyah Hariani

Submission date: 26-Jul-2021 08:43AM (UTC+0700)

Submission ID: 1623997646

File name: The_role_of_laserpuncture_exposure-Pungky-2.pdf (266.45K)

Word count: 3266

Character count: 17814

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Egyptian Journal of Aquatic Research

journal homepage: www.sciencedirect.com/locate/ejar

Full length article

The role of laserpuncture exposure on gonad maturation mechanism of catfish (*Clarias* sp.) through Ca^{2+} , PKC and GABA neurotransmitter

Pungky Slamet Wisnu Kusuma*, Dyah Hariani

Faculty of Science and Mathematics, University of PGRI Adi Buana Surabaya, Surabaya, Indonesia



19

ARTICLE INFO

Article history:

Received 14 July 2016

Revised 19 September 2017

Accepted 18 October 2017

Available online 11 December 2017

Keywords:

GABA

Gonad maturation

Spawning

Laserpuncture

Calcium

PKC

GABA

ABSTRACT

Laser puncture exposure at reproduction acupoint is proven to increase cellular activity like Ca^{2+} in the skin tissues. The aim of the study is to determine the role of laserpuncture exposure on gonad maturation by evaluating Ca^{2+} stimulation and PKC activity in skin tissue and the release of GABA from GABAergic neurons of the brain tissue of catfish (*Clarias* sp.). A total of 36 females and 36 males of 8–9-month old of F1 catfish broodstock *Sangkuriang* (female) and *Paiton* (male). This study used Completely Randomized Design (CDR) experimental method. Expression analysis was conducted using immunohistochemical staining with a streptavidinbiotin method with calcineurin kit, PKC kit, and GABA kit. The results showed that laserpuncture can stimulate calcineurin and PKC expression in skin tissue, and GABA expression in the brain tissue on the condition pre-spawn, spawn, and post-spawn ($P < .05$). It can be concluded that laserpuncture stimulates gonad maturation through Ca^{2+} , PKC, and GABA neurotransmitter.

© 2017 National Institute of Oceanography and Fisheries. Hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Laser is a short-wave light that can inhibit and stimulate biological tissues (Chaves, 2014). Low power laser can provide such biological stimulation improve cellular activity by affecting the membrane permeability to sodium, potassium, and calcium. This mechanism increases the activity of the enzyme (Hunt et al., 2009; Hashmi et al., 2011; Ross et al., 2015). Low-power laser with power 5 mW and 632.8 nm wavelength has been studied and applied to the Black varieties GIFT Tilapia (Tilapia Farmer Genetic Improvement) newly females spawned the first time. That laser inducted at the reproduction acupoint for 6 s on 2/6 ventral part of the body with the firing frequency of once a week can increase Gonado Somatic Index (GSI), proves that cellular activity occurs at the reproductive acupoint that is induced by laserpuncture (Pryor, 2011; Yao et al., 2014).

Cellular activity at the reproduction acupoint caused by laserpuncture exposure will affect to the release and storage process of Ca^{2+} from Endoplasmic Reticulum (ER). The laserpuncture plays role electromagnetic wave energy that can cause nerve cell membrane receptors such as G-protein subunit α would

undergo phosphorylation to activate phospholipase C PLC (Kusuma and Mahendra, 2010). PLC will further hydrolyze phosphatidylinositol bisphosphate (PIP_2) into inositol triphosphate (IP_3) and diacylglycerol (DAG). In the ER Membrane, IP_3 bind to inositol triphosphate receptor (insP3R) to stimulate the release of RE Ca^{2+} . This pathway of cellular activities is known as a metabotropic path.

The release of Ca^{2+} from the ER will increase the levels of Ca^{2+} in the cytoplasm then Ca^{2+} will activate calcineurin. So calcineurin can be an indicator for Ca^{2+} stimulation. Ca^{2+} with DAG activates protein kinase C (PKC). PKC activity can increase the transcription of specific genes. Ca^{2+} and PKC play a role in physiological processes in the cell such as the synthesis of neurotransmitters. However, metabotropic path is not the main pathway. The induction could activate ionotropic pathway, starting from laserpuncture exposure on reproductive acupoint energy of the electromagnetic wave laser beam will be converted into an electrical signal. The electrical signal will cause nerve cell membrane depolarization. Membrane depolarization causes action potential of nerve cell membranes which responds with the opening of extracellular Ca^{2+} channels. Extracellular Ca^{2+} enters through the Calcium Sensing Receptor (CaSR) or via Voltage-Gated Calcium Channels (VGCC) (Clapham, 2007; Chow et al., 2011). Due to the influx of extracellular Ca^{2+} , Ca^{2+} meets with synaptic vesicles and membrane vesicles and opens to release neurotransmitters by exocytosis into the cleft at synapses. Neurotransmitter binds to

Peer review under responsibility of National Institute of Oceanography and Fisheries.

* Corresponding author.

E-mail address: pungky@unipasby.ac.id (P.S.W. Kusuma).

<https://doi.org/10.1016/j.ejar.2017.10.006>

1687-4285/© 2017 National Institute of Oceanography and Fisheries. Hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

specific receptors in postsynaptic can result in a positive or negative effect on the postsynaptic cell membrane. If the bond with neurotransmitter receptors in the postsynaptic matches, the impulse will continue towards the brain where causes a series of physiological reactions in activating the enzyme *Glutamic Acid Decarboxylase* (GAD-65) which will further stimulate the GABAergic neurons to synthesize GABA in brain tissue. The release of GABA can be seen from the expression of GABA-PKC and GABA-Calcineurin in brain tissue.

γ -Amino butyric acid (GABA) is an immunoreactive group of amino acids that are found in the main neuroendocrine area (Mueller et al., 2008; Lado et al., 2014; Watanabe et al., 2014). Telencephalon and hypothalamus have the most prominent stimulatory acts to increase the presence of GnRH release and inhibit dopamine (DA) in the hypothalamus-pituitary complex (Popesku et al., 2008). In principle, DA may reduce the influx of extracellular Ca^{2+} and inhibit the release mechanism of the GnRH release. Thus DA is the control center in the release of GtH-I and GtH-II of the pituitary (Chang et al., 2009; Popesku et al., 2010; Kim et al., 2011). GtH-I plays a role in growth, steroidogenesis and vitellogenesis process while GtH-II has a role in end of gonad maturation, ovulation and spawning (Vijayan et al., 2007; Pham et al., 2010; Yousefian and Mousavi, 2011; Pandey, 2013; Swanson, 2014; Shi et al., 2015). Gonad development and spawning regulated by the hypothalamic-pituitary axis regulates the gonads and liver (Kah and Dufour, 2011; Urbatzka et al., 2011).

However, basic information about laserpuncture effect on Ca^{2+} and PKC expression in skin tissue and the release of GABA is important. Therefore, we examined the role of laserpuncture exposure on gonad maturation by evaluating Ca^{2+} stimulation and PKC activity in skin tissue and the GABA release in the brain tissue of catfish (*Clarias* sp.).

16 Materials and methods

This research was conducted in Freshwater Aquaculture Management Unit (FAMU) Kepanjen, Indonesia. Samples used prospective parent catfish (*Clarias* sp.) gonads that have not been spawned. F1 catfish aged 9 months old were collected from broodstock Sangkuriang (female) and Paiton (male). A total of 36 females and males with weight of 1010–1690 g and 1140–1750 g were used as samples. The female catfish were treated for 15 s with laserpuncture (soft laser Helium-Neon (He-Ne); 5 mW and wavelength 632.8 nm) at the reproductive acupoint (2/3 of the ventral body) until gonads matured (Hashmi et al., 2011; Yao et al., 2014). Completely randomized design was used as an experimental method. The treatments consisted of 6 levels with six repetitions. The treatments were pre-spawn, spawn and post-spawn.

Calcineurin, PKC and GABA expression analysis

The experiment was performed to determine the histology scores calcineurin (expression (Ca^{2+} indicator) and PKC in skin tissue, and GABA expression in the brain tissue by isolating the

skin and brain tissues after exposure to by laserpuncture. Five micrograms of skin thickness and brain tissue was prepared and stained by immunohistochemical staining. The activity of calcineurin, PKC and GABA detected by cell presentation and colour intensity that arise as a result of enzymatic reactions that occur between the HRP enzyme with its substrate. Cell expression indicated calcineurin was red, PKC indicated brown and GABA expression showed dark blue colour under a light microscope with 1000x magnification. The results of calculations performed 10 fields of view to obtain an average of 1 time the field of view. The obtained data were compared between control and treated.

14 Statistical analysis

Data was analyzed using SPSS software (one-way ANOVA). Significant different used $p < .05$ and LSD test at 95% used for the confidence level.

Result

Expression of calcineurin-PKC in the skin tissue of *Clarias* sp. on pre-spawn, spawn and post-spawn after treatment showed high expression respectively by 117%, 43.8%, and 33.3% compared with the control group. The expression of calcineurin-GABA after laserpuncture exposure on the pre-spawn, spawn and post-spawn were higher and precisely at 300%, 46.4% and 97% respectively than the control group. The expression of PKC-GABA after laserpuncture exposure on the pre-spawn, spawn and post-spawn were also higher and recorded as 188.3%, 83.5% and 86.7% respectively than the control. The statistical analysis showed calcineurin, PKC and GABA expression on the pre-spawn, spawn and post-spawn were significantly different ($P < .05$) than the control (Table 1).

Discussion

The calcineurin and PKC expression in the skin tissue of *Clarias* sp. on pre-spawn, spawn and post-spawn were higher after exposure to laserpuncture. The laserpuncture properties was adopted from Kusuma and Mahendra (2010) that has lowpower (soft laser) Helium-Neon (He-Ne) with a wavelength of 632.8 nm, the power of 5 mW/cm² and 0.2 cm² wide light outputs which 15 s are equivalent to an energy of 0.375 Joules/cm²/reproduction acupoint. This energy can induce ER for releasing Ca^{2+} . It happens because when laserpuncture expose the reproductive acupoint, the energy of the electromagnetic wave laser can penetrate the skin tissue to the dermis and epidermis which is suspected to have on peripheral nerve endings. Also, IP₃ headed into the cytosol and binds to specific receptors on the ER and associated with Ca^{2+} channels trigger the release of Ca^{2+} from ER to the cytosol, thereby it may increase the intracellular levels of Ca^{2+} . The receptor activation of phospholipase pathway gains some second messengers such: DAG, IP₃, and Ca^{2+} . DAG and Ca^{2+} plays a role in activating PKC. PKC activity depends on Ca^{2+} . PKC plays a role in the synthesis of specific genes. Ca^{2+} activity intracellular can be seen by calcineurin expression.

Table 1
Calcineurin and PKC expression in the skin tissue compare with calcineurin and GABA, PKC-GABA expression in the brain tissue of catfish (*Clarias* sp.).

Group		N	Calcineurin-PKC	Calcineurin-GABA	PKC-GABA
Control	Pre-spawn	6	4.7 ± 1.9 ^a	3.2 ± 2.4 ^a	6.0 ± 3.6 ^a
	Spawn	6	13.0 ± 4.0 ^b	13.8 ± 3.7 ^{cd}	15.8 ± 2.9 ^c
	Post-spawn	6	9.0 ± 3.0 ^b	6.5 ± 2.9 ^b	9.8 ± 2.3 ^b
Laserpuncture exposure	Pre-spawn	6	10.2 ± 1.5 ^d	12.8 ± 3.3 ^c	17.3 ± 3.4 ^{cd}
	Spawn	6	18.7 ± 1.8 ^e	20.2 ± 2.0 ^e	29.0 ± 1.4 ^e
	Post-spawn	6	12.0 ± 4.0 ^f	12.8 ± 3.5 ^c	18.3 ± 4.9 ^d

Notes: different letter means significant different.

Ca²⁺ along with PKC play a part in the physiological activity of neurotransmitter release. Electromagnetic wave energy of laserpuncture exposure may be converted to electrical signals. This depolarization effect will cause an action potential in nerve cell membranes and the membranes of nerve cells will respond by opening Ca²⁺ channel allowing extracellular Ca²⁺ entry through the Calcium Sensing Receptor (CaSR) or via VoltageGated Calcium Channels (VGCC) (Clapham, 2007; Chow et al., 2011). Due to the influx of extracellular Ca²⁺, intracellular Ca²⁺ will increase. Increased intracellular Ca²⁺ will stimulate the release of neurotransmitters, which will activate a GAD-65 that is responsible for GABA synthesis (Deidda et al., 2014).

GABA stimulates neurons in the hypothalamus and then released GnRH (Maffucci and Gore, 2010; Shi et al., 2015). So GABA can trigger gonadotropin hormone release (GtH-I and GtH-II) (Popescu et al., 2008; Dedden, 2011; Bencic et al., 2013; Karigo and Oka, 2013; Ogawa and Parhar, 2014). GtH-I has a role in stimulating growth, gametogenesis and gonad steroidogenesis fish. While GtH-II has a role in spawning and gamete maturation (Pandolfi et al., 2009; Martyniuk et al., 2009).

This study demonstrates that exposing laserpuncture at reproductive acupoint of *Clarias* sp. could activate enzyme GAD-65 in the brain tissue which will further stimulate the GABAergic neurons to synthesize GABA. In the control group, there was also an increase in GABA expression in the brain tissue on pre-spawn, spawn, and post-spawn. But GABA expression in the brain tissue was still low when compared with the group exposed to laserpuncture. Improvement can occur due to external stimuli such as chemicals (pheromones) released by the opposite sex in a pool. The previous study reported other factors that can affect the spawning as photoperiod, a substrate for spawning, the addition of new water and pheromones that come from the opposite sex in a pool (Reed and Jennings, 2011).

Conclusion

Laserpuncture stimulates gonad maturation through Ca²⁺, PKC, and GABA neurotransmitter. Its exposure at skin tissue of brain activates enzyme GAD-65 in the brain tissue for stimulating GABAergic neurons to synthesize GABA.

References

- Bencic, D.C., Villeneuve, D.L., Biales, A.D., Blake, L., Durhan, E.J., Jensen, K.M., Kahl, M. D., Makynen, E.A., Martinović-Weigelt, D., Ankley, G.T., 2013. Effect of the insecticide fipronil on reproductive endocrinology in the fathead minnow. *Environ. Toxicol. Chem.* 32 (8), 1828–1834.
- Chang, J.P., Johnson, J.D., Sawisky, G.R., Grey, C.L., Mitchell, G., Booth, M., Volk, M.M., Parks, S.K., Thompson, E., Goss, G.G., Klausen, C., Habibi, H.R., 2009. Signal transduction in multifactorial neuroendocrine control of gonadotropin secretion and synthesis in teleosts- studies on the goldfish models. *Gen. Comp. Endocrinol.* 161, 42–52.
- Chaves, M.E.A., 2014. Effect of low-power light therapy on wound healing: LASER x LED. *An. Bras. Dermatol.* 89 (4), 616–623.
- Clapham, D.E., 2007. Calcium signaling. *Cell* 131, 1047–1058.
- Chow, J.Y.C., Estrema, C., Orneles, T., Dong, X., Barret, K.E., Dong, H., 2011. Calcium-sensing receptor modulates extracellular Ca²⁺ entry via TRPC-encoded receptor-operated channels in human aortic smooth muscle cells. *Am. J. Physiol. Cell Physiol.* 301 (2), C461–C468.
- Dedden, I., 2011. The Effect of Gonadotropin Releasing Hormone on Opsin Gene Expression and Spectral Sensitivity in Zebra Cichlid Fish (*Metricula zebra*). Thesis. Queen's University, Canada.
- Deidda, G., Bozart, I.F., Cacedda, L., 2014. Modulation of GABAergic transmission in development and neurodevelopmental disorders: investigating physiology and pathology to gain therapeutic perspectives. *Front. Cell. Neurosci.* 8, 119.
- Hashmi, J.T., Huang, Y., Sharma, S.K., Kurup, D.B., Taboada, L.D., Carroll, J.D., Hamblin, M.R., 2011. Effect of pulsing in low-level light therapy. *Lasers Surg. Med.* 42 (6), 450–466.
- Hunt, R.W., Zavalin, A., Bhatnagar, A., Chinnasamy, S., Das, K.C., 2009. Electromagnetic biostimulation of living cultures for biotechnology, biofuel and bioenergy applications. *Int. J. Mol. Sci.* 10 (10), 4515–4558.
- Kah, O., Dufour, A., 2011. Conserved and divergent features of reproductive neuroendocrinology in teleost fishes. *Hormones and Reproduction of Vertebrates*, 15–42.
- Karigo, T., Oka, Y., 2013. Neurobiological study of fish brains gives insight into the nature of gonadotropin-releasing hormone 1–3 neurons. *Front. Endocrinol. (Lausanne)* 4, 177.
- Kim, D.J., Suzuki, Y., Aida, K., 2011. The control mechanism of gonadotropin-releasing hormone and dopamine on gonadotropin release from cultured pituitary cells of rainbow trout *Oncorhynchus mykiss* at different reproductive stages. *Fish. Aquat. Sci.* 14 (4), 379388.
- Kusuma, P.S.W., Mahendra, A.P.W., Aulanni'am, Marsoedi, 2010. Calcineurin expression against the protein kinase C in catfish (*Clarias* sp.) skin tissue following laserpuncture exposure at reproduction acupoints. *IJET* 12 (5), 97–100.
- Lado, W.E., Spanswick, D.C., Lewis, J.E., Trudeau, V.L., 2014. Electrophysiological characterization of male goldfish (*Carassius auratus*) ventral preoptic area neurons receiving olfactory inputs. *Front. Neurosci.* 8, 185.
- Maffucci, J.A., Gore, A.C., 2010. Hypothalamic neural systems controlling the female reproductive life cycle: Gonadotropin-releasing hormone, glutamate and GABA. *Int. Rev. Cell. Mol. Biol.* 274, 69.
- Martyniuk, C.J., Kroll, K.J., Porak, W.F., Steward, C., Grier, H.J., Denslow, N.D., 2009. Seasonal relationship between gonadotropin, growth hormone, and estrogen receptor mRNA expression in the pituitary gland of largemouth bass. *Gen. Comp. Endocrinol.* 163 (3), 306–317.
- Mueller, T., Wullmann, M.F., Guo, S., 2008. Early teleostean basal ganglia development visualized by zebrafish *Dlx2a*, *Lhx6*, *Lhx7*, *Tbr2* (*eomesa*), and *GAD67* gene expression. *J. Comp. Neurol.* 507, 1245.
- Ogawa, S., Parhar, I.S., 2014. Structural and functional divergence of gonadotropin-inhibitory hormone from jawless fish to mammals. *Front. Endocrinol.* 5 (177), 1–17.
- Pandey, A.K., 2013. Dietary and hormonal manipulations for gonadal maturation and seed production of Indian major carps and catfish. *J. Exp. Zool. India* 16 (1), 19–37.
- Pandolfi, M., Canepa, M.M., Meijide, F.J., Alonso, F., Vazquez, G.R., Maggese, C., Vissio, P.G., 2009. Studies on the reproductive and developmental biology of *Cichlasoma dimerus* (Perciformes, Cichlidae). *Biocell* 33 (1).
- Pham, H.Q., Nguyen, A.T., Nguyen, M.D., Arukwe, A., 2010. Sex steroid levels, oocyte maturation and spawning performance in Waigieu sea perch (*Psammoperca waigiensis*) exposed to thyroxin, human chorionic gonadotropin, luteinizing hormone releasing hormone and carp pituitary extract. *Comp. Biochem. Physiol. Part A* 155, 223–230.
- Popescu, J.T., Martyniuk, C.J., Mennigen, J., Xiong, H., Zhang, D., Xia, X., Cossins, A.R., Trudeau, V.L., 2008. The goldfish (*Carassius auratus*) as a model for neuroendocrine signaling. *Mol. Cell. Endocrinol.* 293, 43–56.
- Popescu, J.T., Martyniuk, C.J., Denslow, N.D., Trudeau, V.L., 2010. Rapid dopaminergic modulation of the fish hypothalamic transcriptome and proteome. *PLoS One* 5 (8), e12338.
- Pryor, A.B., 2011. Advances in laser therapy for the treatment of work related injuries. *Current Perspectives in Clinical Treatment & Management Workers Compensation Cases*. 191201.
- Reed, B., Jennings, M., 2011. Guidance on the housing and care of Zebrafish (*Danio rerio*). Research Animals Department. RSPCA, UK.
- Ross, C.L., Siriwardane, M., Porada, G.A., Porada, C.D., Brink, P., Crist, G.J., Harrison, B. S., 2015. The effect of low-frequency electromagnetic field on human bone marrow stem/progenitor differentiation. *Stem Cell Res.* 15 (1), 96–108.
- Shi, B., Liu, X., Xu, Y., Wang, A., 2015. Molecular characterization of three gonadotropin subunit and their expression patterns during ovarian maturation in *Cynoglossus semilaevis*. *Int. J. Mol. Sci.* 16 (2), 2767–2793.
- Swanson, P., 2014. Environmental and endocrine control of reproduction in cultured salmonids. Coastal Zone and Estuarine Studies Division. Northwest Fisheries Science Center, Washington.
- Urbatzka, R., Rocha, M.J., Rocha, E., 2011. Regulation of ovarian development and function in teleosts. In: Norris, D.O., Lopez, K.H. (Eds.), *Hormones and Reproduction of Vertebrates*. Academic Press, pp. 65–82.
- Vijayan, K.K., Jayasankar, P., Vijayagopal, P., 2007. Indian Fisheries: A Progressive Outlook. Central Marine Fisheries Research Institute, India.
- Watanabe, M., Fukuda, A., Nabekura, J., 2014. The role of GABA in the regulation of GnRH neurons. *Front. Neurosci.* 8, 387.
- Yao, W., Yang, H., Yin, N., Ding, G., 2014. Mast cell-nerve cell interaction at acupoint: Modelling mechanotransduction pathway induced by acupuncture. *Int. J. Biol. Sci.* 10 (5), 511–519.
- Yousefian, M., Mousavi, S.E., 2011. The mechanism of reproduction and hormonal function in finfish species: a review. *Sci. Res. Essays* 6 (17), 3561–3570.

Further reading

- Taylor, C.W., Tovey, S.C., 2010. IP3 receptor: Toward understanding their activation. *Cold Spring Harb. Perspect. Biol.* 2 (12), a004010.

The role of laserpuncture exposure on gonad maturation mechanism of catfish (Clarias sp.) through Ca²⁺, PKC and GABA neurotransmitter

ORIGINALITY REPORT

16%

SIMILARITY INDEX

13%

INTERNET SOURCES

11%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1	profiles.uonbi.ac.ke Internet Source	2%
2	www.laser-therapy.us Internet Source	2%
3	ijens.org Internet Source	1%
4	karyailmiah.unipasby.ac.id Internet Source	1%
5	journals.physiology.org Internet Source	1%
6	macrothink.org Internet Source	1%
7	Seyed Ehsan Mousavi. "Effects of exogenous hormones on plasma cortisol, sex steroid hormone and glucose levels in male and female grass carp, <i>Ctenopharyngodon idellus</i> , during the spawning induction", AFRICAN JOURNAL OF BIOTECHNOLOGY, 2012	1%

8	jurnal.unipasby.ac.id Internet Source	1 %
9	Milton Levin, Brenda Morsey, Sylvain De Guise. "Non-coplanar PCBs Induce Calcium Mobilization in Bottlenose Dolphin and Beluga Whale, but not in Mouse Leukocytes", Journal of Toxicology and Environmental Health, Part A, 2007 Publication	<1 %
10	departments.agri.huji.ac.il Internet Source	<1 %
11	mts.intechopen.com Internet Source	<1 %
12	Niladri Basu, Chieu Anh Ta, Andrew Waye, Jinqin Mao, Mark Hewitt, John T. Arnason, Vance L. Trudeau. "Pulp and Paper Mill Effluents Contain Neuroactive Substances That Potentially Disrupt Neuroendocrine Control of Fish Reproduction", Environmental Science & Technology, 2009 Publication	<1 %
13	Toshihide Tabata, Masanobu Kano. "GABAB Receptor-Mediated Modulation of Metabotropic Glutamate Signaling and Synaptic Plasticity in Central Neurons", Elsevier BV, 2010	<1 %

14

www.utoledo.edu

Internet Source

<1 %

15

Houguo Xu, Bo Sun, Zhangbin Liao, Elena Pribytkova, Qinggong Zhang, Yuliang Wei, Mengqing Liang. " Possible involvement of PKC/MAPK pathway in the regulation of GnRH by dietary arachidonic acid in the brain of male tongue sole ", Aquaculture Research, 2019

Publication

<1 %

16

eprints.uthm.edu.my

Internet Source

<1 %

17

journals.sagepub.com

Internet Source

<1 %

18

media.neliti.com

Internet Source

<1 %

19

theses.ucalgary.ca

Internet Source

<1 %

20

www.scribd.com

Internet Source

<1 %

21

Nicholls, John G.. "From Neuron to Brain", Oxford University Press

Publication

<1 %

22

Thomas Mueller, Mario F. Wullimann, Su Guo. "Early teleostean basal ganglia development

<1 %

visualized by ZebrafishDlx2a,Lhx6,Lhx7,Tbr2 (eomesa), andGAD67 gene expression", The Journal of Comparative Neurology, 2008

Publication

23

Sethu Selvaraj, Hajime Kitano, Hirofumi Ohga, Akihiko Yamaguchi, Michiya Matsuyama.

"Expression changes of mRNAs encoding kisspeptins and their receptors and gonadotropin-releasing hormones during early development and gonadal sex differentiation periods in the brain of chub mackerel (*Scomber japonicus*)", General and Comparative Endocrinology, 2015

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

The role of laserpuncture exposure on gonad maturation mechanism of catfish (Clarias sp.) through Ca²⁺, PKC and GABA neurotransmitter

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
