

# Phytochemical Screening, Total Flavonoid Content, and Total Phenolic Content of Ethanol Extract of the Indonesian Fern *Selaginella Plana*

*by I G M Sanjaya*

---

**Submission date:** 14-Mar-2022 06:09AM (UTC+0700)

**Submission ID:** 1783410987

**File name:** 125966487.pdf (564.41K)

**Word count:** 3467

**Character count:** 18845

# Phytochemical Screening, Total Flavonoid Content, and Total Phenolic Content of Ethanol Extract of the Indonesian Fern *Selaginella Plana*

Suyatno Sutoyo<sup>1,\*</sup> Amaria<sup>1</sup> I G. M. Sanjaya<sup>1</sup> Rusly Hidayah<sup>1</sup>

Devy Puspita Sari<sup>1</sup> Nurrulhidayah A. Fadzlillah<sup>2</sup>

<sup>1</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Negeri Surabaya, Indonesia

<sup>2</sup>International Institute for Halal Research and Training (INHART), International Islamic University Malaysia (IIUM), Gombak 53100, Kuala Lumpur

\*Corresponding author E-mail: [suyatno@unesa.ac.id](mailto:suyatno@unesa.ac.id).

## ABSTRACT

*Selaginella plana* (Cakar ayam) is one of the ferns that grow widely in Indonesia and has been used by the community as food, ornamental plants and traditional medicines to treat wounds, respiratory infections, liver disorders, urinary tract infections, fractures, rheumatism, cancer and increase the body immunity. Research on the chemical constituents of this plant has not been widely reported. This study aimed to describe secondary metabolites from ethanol extract of *S. plana* and determine total flavonoid and total phenolic content. Secondary metabolites content was determined using phytochemical tests, total phenolic and total flavonoid content were determined using UV-Vis spectroscopy. The results showed that the ethanol extract of *S. plana* contained secondary metabolites of steroids, saponins, phenolics, flavonoids, alkaloids, and tannins. Based on the results of determination using the UV-Vis spectroscopy, it was known that the ethanol extract of *S. plana* has total flavonoid and a total phenolic content of  $35.725 \pm 0.110$  mg QE/ g sample mass and  $36.363 \pm 0.143$  mg GAE/ g dried extract, respectively.

**Keywords:** *Selaginella plana*, Phytochemical screening, Total phenolic content, Total flavonoid content.

## 1. INTRODUCTION

*Selaginella plana* (cakar ayam) is one of the ferns which becomes Indonesia's biological wealth [1]. This plant is spread in Southeast Asia, in Java it grows to a height of approximately 750 above sea level, generally on the moist and protected riverbank and on the slopes of ravines. *S. plana* has been known and used by society as food (vegetables), ornamental plants, and traditional medicines. It has been used for postpartum care, treating wounds, treating bleeding, menstrual disorders, increasing the body immunity, as well as a postnatal tonic [2-4]. Traditionally, *S. plana* has been used in the treatment of several diseases such as cancer, respiratory infections, liver disorders, urinary tract infections, fractures and rheumatism [5].

Chikmawati, et al. reported that *Selaginella* species on the island of Java was found to have

secondary metabolites of the flavonoid group, with the highest content of biflavonoid compounds [6]. Biflavonoid compounds from *Selaginella* genus have various bioactive properties, including antioxidants, anticancer, antibacterial, antiviral, antifungal, antimalarial, and anti-inflammatory properties [7-10].

As a phenolic compound, biflavonoid had been extracted from several fern species in Selaginellaceae family. Amentoflavone was extracted from *S. tamariscina* and *S. bryopteris*, [11,12]. From *S. lepydophylla*, robustataflavone, ginkgetin, kayaflavone and podocarpus flavone A had been extracted [13]. Meanwhile, hinokiflavone had been reported from *S. cauculosa* and *S. bryopteris* [12][14]. Research on the content of phenolic compounds and flavonoid content in the fern *S. plana* has never been reported. This paper will report on results of the phytochemical screening

18 ethanol extract from *S. plana* and determining total flavonoid and total phenolic content.

## 2. METHODS

### 2.1. Material

The dried powder of *S. plana* aerial parts, ethanol (80%, p.a.), methanol, acetic anhydride, concentrated sulfuric acid, hydrochloric acid (1 M, 2 M, concentrated), magnesium ribbon, FeCl<sub>3</sub> (1%, 5%), Mayer reagent, Dragendorff reagent, Wagner reagent, gallic acid (Merck), Folin-Ciocalteu reagent (Merck), Na<sub>2</sub>CO<sub>3</sub> 7.5%, quercetin (Sigma), AlCl<sub>3</sub> 10%, ammonia, sodium acetate 1 M.

### 2.2. Instruments

Spectrophotometer UV-Vis (Shimadzu UV-1800), magnetic stirrer (Heidolph), rotary vacuum evaporator (Buchi R-300), oven (Heraeus ST-5042), analytical balance (Advanturer Ohaus), vacuum pump (Gast DOA-P-504-BN), water bath (Memmert), freeze dryer (Martin Christ Alpha 1-2 LDplus), vortex mixer, volumetric flask, beaker glass, Buchner funnel, micropipette, test tube, drip plate, spatula, maceration vessel.

### 2.3. Research procedures

#### 2.3.1. Collection and preparation of sample

Samples of *S. plana* were collected from the Kletak forest, Nongkojajar, Pasuruan, East Java. Before the further investigation, the sample was identified at LIPI Botanical Garden, Purwodadi 23<sup>st</sup> Java. Sample was cleaned of attached dirt, then dried at room temperature for 12 days. The dried sample was grinded into a fine powder ready for extraction [15].

#### 2.3.2. Extraction

Dried powder of *S. plana* aerial part (1000 g) was macerated with ethanol 80% for 11<sup>h</sup> 24 hours. Then filtered using a Buchner funnel. The ethanol extract obtained was evaporated in vacuum using rotary vacuum evaporator, resulted the concentrated extract. It was dried in freeze dryer to produce dried extract [15].

#### 2.3.3. Phytochemical screening

##### 2.3.3.1. Triterpenoid and steroid test (Liebermann-Burchard test)

About 1 mg of ethanolic extract was placed on a drip plate, added 3 drops of acetic anhydride and stirred with a spatula until dissolved. Then one drop

38 of concentrated sulfuric acid was added to the mixture [16].

##### 2.3.3.2. Phenolic test

About 1 mg ethanolic extract was placed on a drip plate, added 5 drops of methanol, then stirred with spatula until dissolved. Then 3 drops of 5% FeCl<sub>3</sub> solution in ethanol was added into mixture [16].

##### 2.3.3.3. Flavonoid test (Shinoda test)

About 1 mg of ethanolic extract was placed on a drip plate, added 5 drops of methanol, then 3-4 small pieces of Mg ribbon and 2 drops of concentrated HCl were added to the mixture [17].

##### 2.3.3.4. Alkaloid test

2 A total of 4 mg of ethanolic extract was dissolved with 3 mL of methanol and 5 mL of ammonia to 28<sup>h</sup> 8-9, then the mixture was filtered. The filtrate was added with 2 mL of 2 M HCl solution. After the mixture was shaken, 5 drops of the top layer were added into 4 test tubes. The solution in tube 1 was used as a blank, solution in tube 2, 3, and 4 was added one drop of Mayer, Wagner, and Dragendorff reagents, respectively [16].

##### 2.3.3.5. Saponin test

17 About 1 mg ethanolic extract was put into a test tube, then added 5 mL of distilled water, shaken for 1 minute. If foam appeared, 2 drops of 1 M HCl solution was added. If foam with height of 1-3 cm did not disappear or lasted for 10 minutes, sample contained saponins [16].

##### 2.3.3.6. Tannin test

About 1 mg of ethanolic extract was placed on a drip plate, added 5 drops of methanol, 12<sup>h</sup> stirred with a spatula until dissolved. Then 2-3 drops of 1% FeCl<sub>3</sub> solution were added. Appearance of blue, green, or brownish green color indicated the presence of tannins [17].

##### 2.3.4. Determination of total phenolic contents

9 A total of 1.0 mL of the extract solution was pipetted from the stock solution (1000 ppm) was put in a test tube, then added 1.5 mL of 36<sup>h</sup> in Ciocalteu reagent (10% = 1:10). After being allowed to stand for 3 minutes, 1.2 mL of 7.5% Na<sub>2</sub>CO<sub>3</sub> was added, mixed for 3 seconds using a vortex mixer. The mixture was allowed to stand for 30 minutes, then its absorbance was measured by UV-Vis spectroscopy

at a maximum absorption wavelength of 7615 nm. The concentration of phenolic compounds in the extract was determined using a standard gallic acid curve with various concentrations of 10, 12, 14, 16, 18, and 20 ppm. The content of phenolic compounds was expressed as milligrams of gallic acid equivalent (GAE)/gram dried extract [18].

### 2.3.5. Determination of total flavonoid contents

A total of 0.5 mL of 250 ppm extract solution was put in a test tube, then 0.1 mL of 10% aluminum (III) chloride, 0.1 mL of 1 M sodium acetate and 2.8 mL of distilled water were added. After being incubated for 30 minutes. The absorbance was measured using a UV-Vis spectrophotometer at a maximum wavelength of 433 nm. The total flavonoid content of the ethanol extract of the fern *S. plana* was determined based on the standard curve of quercetin with various concentrations of 20, 30, 40, 50, and 60 ppm. The total flavonoid content was expressed as milligrams of quercetin equivalent (QE)/gram of dried extract [19].

## 3. RESULTS AND DISCUSSION

### 3.1. Extraction of *S. plana* aerial part

The dried powder of *S. plana* aerial part (1000 g) was macerated using ethanol 80% as a solvent for 24 hours and repeated three times. Evaporation of the solvent in vacuo, followed by drying at freeze drier for 16 hours, yielded the dark green solid (49.14 g).

### 3.2. Phytochemical screening

Phytochemical screening was carried out to determine the group of secondary metabolites contained in the ethanol extract of the fern *S. plana*. The tests carried out included steroids-triterpenoids test, phenolic test, flavonoids test, alkaloid test, saponin test, and tannin test.

Steroid-triterpenoid test was carried out using Liebermann-Burchard reagent. Ethanol extract of *S. plana* after being dissolved with acetic anhydride and dripped with concentrated sulfuric acid produced a bluish-green solution. The appearance of blue color in the test with the Liebermann-Burchard reaction indicated that the sample contained steroid compounds [16].

Phenolic test was carried out using ferric chloride reagent with a concentration of 5% in ethanol. The ethanol extract of *S. plana* after being dissolved in methanol, then dripped with 5% ferric chloride solution turned out to be a purple solution. The appearance of green, blue, or purple color indicates that the sample contains phenolic compounds. These colors occur as result of formation of colored complex ions between ferric ions ( $Fe^{3+}$ ) and phenolic ions in phenolic compounds [16,20].

In this study, flavonoid test was conducted using Shinoda test. After being dissolved in methanol, 3-4 pieces of magnesium ribbon were added to the ethanol extract of *S. plana*. After adding 1 M hydrochloric acid solution, a yellow, orange, blue, or red colored solution was formed indicating the presence of flavonoid compounds in the ethanol extract of *S. plana*. The color occurred was result of the formation of colored complex ions between magnesium ions ( $Mg^{2+}$ ) and phenolic ions in phenolic compounds [17].

The alkaloid test was carried out using Mayer, Wagner and Dragendorff reagents. The ethanol extract of *S. plana* after being tested using the three reagents showed positive results because it showed white, brown and orange precipitates after being added with Mayer, Wagner and Dragendorff reagents, respectively [16,20].

The test for the presence of saponin compounds in the ethanol extract was conducted using Forth test. A number of *S. plana* ethanolic extract was dissolved in water and then heated. It turns out that a stable foam was formed. These results indicated that the extract contained secondary metabolites of the saponin group [16,20].

In this study, the tannin test was carried out with 1% ferric chloride reagent in ethanol. Ethanol extract solution of *S. plana* after being reacted with 1% ferric chloride solution formed greenish-blue solution. According to Edeoga, *et al* the formation of blue, green, or brownish green color indicated the presence of tannin compounds in the sample. Thus, the ethanol extract of *S. plana* contained compounds belonging to the tannin group. The color was produced due to the formation of a complex between ferric ions and phenolic ions from tannin compounds [17]. Results of the phytochemical tests above are briefly presented in the Table 1.


**Table 1.** Phytochemical test of ethanol extract of *S. plana*

No.	Phytochemicals	Test methods	Results (Indicating positive test)
1	Steroids and triterpenoids	Liebermann-Burchard test	A bluish green color + (Steroids)
2	Phenolic	Ferric chloride test	A purple color +
3	Flavonoids	Shinoda test	A orange color +
4	Alkaloid	- Mayer test - Dragendorff test - Wagner test	- A white precipitate - A orange precipitate - A chocolate precipitate +
5	Saponins	Forth test	Persistent foam +
6	Tannins	Ferric chloride test	A greenish blue color +

Phytochemical test showed that the ethanol extract of *S. plana* contained secondary metabolites of steroids, phenolics, flavonoids, alkaloids, saponins, and tannins. The presence of phenolics and flavonoids compounds supports the results of previous studies which stated that the ferns of the genus *Selaginella* from Java contain flavonoid compounds, especially biflavonoids [6]. Beside that it also supports the potential use of *S. plana* as an Indonesian medicinal plant.

**3.3. Total phenolic content**

The total phenolic content in the ethanol extract of the fern *S. plana* was determined by the spectroscopic method using Folin-Ciocalteu reagent. The reagent will be reduced by phenolic compounds in alkaline condition to produce a blue phosphotungstate-phosphomolybdate complexes. Alkaline condition is needed to convert the phenolic hydroxyl group into phenolate ions due to proton dissociation [18]. Based on measurement of the absorbance of the standard solution of gallic acid, it was obtained the standard curve presented in Figure 1.




**Figure 1** Standard curve of gallic acid solution.

The total phenolic content was determined based on the standard gallic acid curve determined by UV-Vis spectroscopy at a wavelength of 763 nm. From the standard curve (Fig 1), it was obtained the

regression equation  $Y = 0.462X - 0.13$  with  $R^2 = 0.9768$ . The results of determining the total phenolic content of the ethanol extract of the fern *S. plana* 1000 ppm using the spectroscopic method from three measurements obtained an absorbance value of 0.558; 0.562 and 0.557. Based on the gallic acid standard curve and taking into account the dilution factor, the total phenolic content in the ethanol extract of the fern *S. plana* was  $36.310 + 36.526 + 36.255$ , respectively with an average value of  $36.364 \pm 0.143$  mg gallic acid equivalent (GAE)/gram dried extract.

**3.4. Total flavonoid content**

Total phenolic content in ethanol extract of fern *S. plana* was determined by an aluminium chloride reagent. This reagent will form yellow complexes with the carbonyl group of flavonoids at the C-4 with an adjacent hydroxyl group at the C-3 (flavonols) or C-5 (flavonols and flavone) and provide specific absorption in UV-Vis spectroscopy [19,21]. The total flavonoid content was determined based on the standard curve of quercetin as measured by UV-Vis spectroscopy at a wavelength of 433 nm. Based on the measurement results of the absorbance of the standard solution of quercetin, the standard curve was obtained which was presented in Figure 2.



**Figure 2** Standard curve of quercetin solution.

The linear regression equation obtained from the standard curve was  $Y = 0.0091X + 0.0298$  with  $R^2 = 0.9649$ . The results of the determination of the total flavonoid content of the ethanol extract of the fern *S. plana* 1000 ppm using the spectroscopic method from three measurements obtained an absorbance value of 0.678; 0.680 and 0.682. Based on the standard curve of quercetin, the total phenolic content in the ethanol extract of the fern *S. plana* were 35.615; 35.726; 35.835, respectively with an average value of  $35.725 \pm 0.110$  mg of quercetin equivalent (QE)/gram dried extract. The total phenolic content and total flavonoid content of *S. plana* correspond with the results of phytochemical tests that it contains phenolics, flavonoids and tannins. The plant containing phenolics and flavonoids compounds such as *S. plana* has the potential to be used as antioxidants [18-20].

#### 4. CONCLUSION

The study concludes that the ethanol extract of *S. plana* contains secondary metabolites of steroids, saponins, phenolics, flavonoids, alkaloids, and tannins. It was found to have total flavonoid a total phenolic content of  $35.725 \pm 0.110$  mg QE/ g dried extract and  $36.363 \pm 0.143$  mg GAE/ g dried extract, respectively.

#### AUTHORS' CONTRIBUTION

Suyatno Sutoyo: conceptualizing the research and writing the manuscript. Amaria, I Gede Made Sanjaya and Nurulhidayah Ahmad Fadzilillah: developing research methods and analyzing data. Rusly Hidayah and Devy Puspita Sari: presenting data and doing data analysis.

#### ACKNOWLEDGMENT

We thank the Universitas Negeri Surabaya for financial support through the foreign collaboration

research grant 2021 and Mr. Sugiono from LIPI Purwodadi Botanical Garden for help in collecting and identifying the fern *Selaginella plana*.

#### REFERENCES

- [1] K. Heyne, *Tumbuhan Berguna Indonesia* Jilid 1, Departemen Kehutanan, 1987.
- [2] A. Uluk, M. Sudana, E. Wollenberg, *Ketergantungan Masyarakat Dayak terhadap Hutan di Sekitar Taman Nasional Kayan Mentarang*, Cifor, 2001.
- [3] K. Harada, M. Rahayu, A. Muzakkir, *Medicinal Plants of Gunung Halimun National Park, West Java, Indonesia*, Biodiversity Conservation Project -JICA, PHPA and LIPI, 2002.
- [4] A. D. Setyawan, Review: Recent Status of *Selaginella* (Selaginellaceae) Research in Nusantara, *Bobierrites Journal of Biological Diversity*, 12(2) (2011) 112-124. DOI: <https://doi.org/10.13057/biodiv/d120209>
- [5] D. Bensky, S. Clavey, E. Stöger, *Chinese Herbal Medicine Materia Medica*, Eastland Press, 2004.
- [6] T. Chikmawati, A. D. Setyawan, Miftahudin, *Phytochemical Composition of Selaginella spp from Java Island Indonesia*, *Makara Journal of Science*, 16(2) (2012) 129-133. DOI: <https://doi.org/10.7454/mss.v16i2.1408>
- [7] M. Sannomiya, C. R. P. Cardoso, M. E. Figueiredo, C. M. Rodrigues, L. C. dos Santos, F. V. dos Santos, J. M. Serpeloni, L. M. S. Colus, W. Vilegas, Eliana, A. Varanda, *Mutagenic Evaluation and Chemical Investigation of Byrsonima Intermedia A. Juss. Leaf Extracts*, *Journal Ethnopharmacol*, 112(2) (2007) 319-326. DOI: <https://doi.org/10.1016/j.jep.2007.03.014>
- [8] Y. M. Lin, *Beta-Fluoroethyl Thiourea Compounds and Use*, United States Patent 6677350, 2004.
- [9] J. J. Chen, C. Y. Duh, J. F. Chen, *New Cytotoxic Biflavonoids from Selaginella delicatula*, *Planta Med.* 71(7) (2005) 659-665. DOI: <https://doi.org/10.1055/s-2005-871273>
- [10] A. R. Ahmad, Juwita, S. A. D. Ratulangi, A. Malik, *Penetapan Kadar Fenolik dan Flavonoid Total Ekstrak Metanol Buah dan Daun Patikala (Etlingera elatior (Jack) R.M.SM)*, *Pharmaceutical Sciences and Research*, 2(1)

- (2015) 1-10. DOI: <http://dx.doi.org/10.7454/psr.v2i1.3481>
- [11] J. W. Yang, Y. R. Pokharel, M. R. Kim, E. R. Woo, H. K. Choi, K. W. Kang, Inhibition of Inducible Nitric Oxide Synthase by Sumaflavone Isolated from Selaginella tamariscina, *Journal Ethnopharmacol*, 105(1-2) (2006) 107-113. DOI: <http://dx.doi.org/10.1016/j.jep.2005.10.001>
- [12] N. K. Sah, S. N. P. Singh, S. Sahdev, S. Banerji, V. Jha, Z. Khan, S. E. Hasnain, Indian Herb 'Sanjeevani' (Selaginella bryopteris) can Promote Growth and Protect Against Heat Shock and Apoptotic Activities of Ultra Violet and Oxidative Stress, *Journal of Biosciences*, 30(4) (2005) 499-505. DOI: <http://dx.doi.org/10.1007/BF02703724>
- [13] Y. Su, C. M. Sun, H. H. Chuang, P. T. Chang, Studies on the cytotoxic mechanisms of ginkgetin in a human ovarian adenocarcinoma cell line, *Naunyn-Schmiedeberg's Archives of Pharmacology*, 362 (2000) 82-90. DOI: <https://doi.org/10.1007/s002100000240>
- [14] R. C. Swamy, Kunert, O. W. Schuhly, F. Bucar, D. Ferreira, V. S. Rani, B. R. Kumar, A. V. N. A. Rao, Structurally Unique Biflavonoids from Selaginella chrysocaulos and Selaginella bryopteris, *Chem Biodiversity*, 3(4) (2006) 405-413. DOI: <https://doi.org/10.1002/cbdv.2006900442006>
- [15] S. Sutoyo, Tukiran, S. Khotijah, Antioxydant Activity of the Silver Nanoparticles (AgNPs) Synthesized Using Nephrolepisradicans Extract as Bioreductor, in *Proceedings of Mathematics, Informatics, Science, and Education International Conference (MISEIC)*, *Journal of Physics: Conference Series*, 2021, pp. 1-6. DOI: <https://doi.org/10.1088/1742-6596/1747/1/012038>
- [16] P. Tiwari, B. Kumar, M. Kaur, G. Kaur, H. Kaur, Phytochemical Screening and Extraction: A Review, *Internationale Pharmaceutica Scientia*, 1 (2011) 98-106. Retrieved from: [https://www.scirp.org/\(S\(1z5mqp453edsnp55rrgjct55\)\)/reference/referencespapers.aspx?referenceid=1949793](https://www.scirp.org/(S(1z5mqp453edsnp55rrgjct55))/reference/referencespapers.aspx?referenceid=1949793)
- [17] H. O. Edeoga, D. E. Okwu, B. O. Mbaebie, Phytochemical constituents of some Nigerian medicinal plants, *African Journal of Biotechnology*, 4(7) (2005) 685-688. DOI: <https://doi.org/10.5897/AJB2005.000-3127>
- [18] A. Kusbandari, H. Susanti, Similarity Check Determination of Total Phenolic Content and Antioxidant Activity of Methanol Extract of Maranta arundinacea L Fresh Leaf and Tuber, in *International Pharmacy Conference UAD (IPCUAD)*, *IOP Conference Series Materials Science and Engineering*, 2017, pp. 1-4. DOI: <https://doi.org/10.1088/1757-899X/259/1/012010>
- [19] A. R. Surana, M. R. Kumbare, R. D. Wagh, Estimation of Total Phenolic and Total Flavonoid Content and Assessment of in vitro Antioxidant Activity of Extracts of Hamelia patens Jacq. Stems, *Research Journal of Phytochemistry*, 10 (2016) 67-74 DOI: <https://dx.doi.org/10.3923/rjphyto.2016.67.74>
- [20] J. R. Shaikh, M. K. Patil, Qualitative tests for preliminary phytochemical screening: An overview, *International Journal of Chemical Studies*, 8(2) (2020) 603-608. DOI: <https://doi.org/10.22271/chemi.2020.v8.i2i.8834>
- [21] E. N. Sembiring, B. Elya, R. Sauriasari, Phytochemical Screening, Total Flavonoid and Total Phenolic Content and Antioxidant Activity of Different Parts of Caesalpinia bonduc (L.) Roxb, *Pharmacognosy Journal*, 10(1) (2018) 123-127. DOI: <http://dx.doi.org/10.5530/pj.2018.1.22>

# Phytochemical Screening, Total Flavonoid Content, and Total Phenolic Content of Ethanol Extract of the Indonesian Fern *Selaginella Plana*

## ORIGINALITY REPORT

24%

SIMILARITY INDEX

18%

INTERNET SOURCES

19%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://irep.iium.edu.my">irep.iium.edu.my</a> Internet Source	3%
2	<a href="http://www.myfoodresearch.com">www.myfoodresearch.com</a> Internet Source	2%
3	<a href="http://mzuir.inflibnet.ac.in:8080">mzuir.inflibnet.ac.in:8080</a> Internet Source	2%
4	Submitted to Universitas Pendidikan Ganesha Student Paper	1%
5	<a href="http://bmccomplementmedtherapies.biomedcentral.com">bmccomplementmedtherapies.biomedcentral.com</a> Internet Source	1%
6	Submitted to Universitas Negeri Surabaya The State University of Surabaya Student Paper	1%
7	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	1%
8	Submitted to Universitas Tadulako Student Paper	1%

9

Ni Luh Putu Mita Rismayanti, Amir Husni.  
"Antioxidant activity of methanolic extract of  
Eucheuma spinosum extracted using a  
microwave", IOP Conference Series: Earth and  
Environmental Science, 2021

Publication

---

1 %

10

Chutima Limmatvapirat, Chutipap  
Nateesathittarn, Kamol Dechasathian, Tharit  
Moohummad, Pana Chinajitphan, Sontaya  
Limmatvapirat. "Phytochemical analysis of  
baby corn silk extracts", Journal of Ayurveda  
and Integrative Medicine, 2020

Publication

---

1 %

11

Dian Anriani A. Makuasa, Purnama Ningsih.  
"The Analysis of Total Flavonoid Levels In  
Young Leaves and Old Soursop Leaves  
(Annona muricata L.) Using UV-Vis  
Spectrophotometry Methods", Journal of  
Applied Science, Engineering, Technology, and  
Education, 2020

Publication

---

1 %

12

Taous Khan, Sayyad Ali, Rahila Qayyum, Izhar  
Hussain, Fazli Wahid, Abdul Jabbar Shah.  
"Intestinal and vascular smooth muscle  
relaxant effect of Viscum album explains its  
medicinal use in hyperactive gut disorders  
and hypertension", BMC Complementary and  
Alternative Medicine, 2016

1 %

---

13	<a href="http://ijpsr.com">ijpsr.com</a> Internet Source	1 %
14	<a href="http://insightsociety.org">insightsociety.org</a> Internet Source	1 %
15	<a href="http://www.hindawi.com">www.hindawi.com</a> Internet Source	1 %
16	<a href="http://www.science.gov">www.science.gov</a> Internet Source	1 %
17	Sri Wahyuni Nasution, Stepfany Monica Helmin, Jessyca Sembiring S, Daniel Arswendo, Novia Fransiska Br S, Ronel Polin Sihombing. "Phytochemical and Antibacterial Analysis of Mangkokan Leaf Extract Against Salmonella typhimurium Bacteria", 2021 IEEE International Conference on Health, Instrumentation & Measurement, and Natural Sciences (InHeNce), 2021 Publication	<1 %
18	Assefa, Awraris Derbie, Eun Young Ko, So Hyun Moon, and Young-Soo Keum. "Antioxidant and antiplatelet activities of flavonoid-rich fractions of three citrus fruits from Korea", 3 Biotech, 2016. Publication	<1 %
19	Submitted to Universiti Sains Malaysia Student Paper	<1 %

---

20 Patrícia de Araújo Rodrigues, Selene Maia de Morais, Liza Araújo Aguiar, Nadja Soares Vila-Nova, Stephen Rathinaraj Benjamin. "Effect of Byrsonima sericea DC. leaf extracts on mice gastrointestinal tract", Toxicology Reports, 2019  
Publication

---

21 [euacademic.org](http://euacademic.org)  
Internet Source

---

22 [www.thinkswap.com](http://www.thinkswap.com)  
Internet Source

---

23 G.M Saibu, D.R Katerere, D.J.G Rees, M Meyer. "In vitro cytotoxic and pro-apoptotic effects of water extracts of Tulbaghia violacea leaves and bulbs", Journal of Ethnopharmacology, 2015  
Publication

---

24 [repository-tnmgrmu.ac.in](http://repository-tnmgrmu.ac.in)  
Internet Source

---

25 [scholarbank.nus.edu.sg](http://scholarbank.nus.edu.sg)  
Internet Source

---

26 [tjs.udsm.ac.tz](http://tjs.udsm.ac.tz)  
Internet Source

---

27 [www.derpharmachemica.com](http://www.derpharmachemica.com)  
Internet Source

---

[www.jddtonline.info](http://www.jddtonline.info)

28

Internet Source

&lt;1 %

29

Russo, Daniela, Patrícia Valentão, Paula Andrade, Eloy Fernandez, and Luigi Milella. "Evaluation of Antioxidant, Antidiabetic and Anticholinesterase Activities of *Smallanthus sonchifolius* Landraces and Correlation with Their Phytochemical Profiles", *International Journal of Molecular Sciences*, 2015.

Publication

&lt;1 %

30

[jurnal.farmasi.umi.ac.id](http://jurnal.farmasi.umi.ac.id)

Internet Source

&lt;1 %

31

[jurnal.radenfatah.ac.id](http://jurnal.radenfatah.ac.id)

Internet Source

&lt;1 %

32

[macrothink.org](http://macrothink.org)

Internet Source

&lt;1 %

33

Maisuthisakul, P.. "Assessment of phenolic content and free radical-scavenging capacity of some Thai indigenous plants", *Food Chemistry*, 2007

Publication

&lt;1 %

34

Muhammad Bilal Azmi, Shamim A. Qureshi. "Methanolic Root Extract of Benth Improves the Glycemic, Antiatherogenic, and Cardioprotective Indices in Alloxan-Induced Diabetic Mice ", *Advances in Pharmacological Sciences*, 2012

&lt;1 %

35

Raushanara Akter, Sharmind Neelotpol, Md. Tanvir Kabir. "Effect of *Allium sativum* methanol extract in amelioration of arsenic-induced toxicity in Swiss albino mice", *Phytomedicine Plus*, 2022

Publication

---

<1 %

36

A Kusbandari, H Susanti. " Determination of total phenolic content and antioxidant activity of methanol extract of L fresh leaf and tuber ", *IOP Conference Series: Materials Science and Engineering*, 2017

Publication

---

<1 %

37

Hajji, M.. "GC/MS and LC/MS analysis, and antioxidant and antimicrobial activities of various solvent extracts from *Mirabilis jalapa* tubers", *Process Biochemistry*, 201009

Publication

---

<1 %

38

Sobha Kota, Vayunandana Rao Govada, Ratna Kumari Anantha, Mahendra Kumar Verma. "An Investigation into phytochemical constituents, antioxidant, antibacterial and anti-cataract activity of *Alternanthera sessilis* , a predominant wild leafy vegetable of South India", *Biocatalysis and Agricultural Biotechnology*, 2017

Publication

---

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On