

# Bird diversity of resort Ranu Darungan

*by* Reni Ambarwati

---

**Submission date:** 16-Jan-2023 09:17AM (UTC+0700)

**Submission ID:** 1993292869

**File name:** Birds\_of\_Ranu\_Darungan.pdf (373.33K)

**Word count:** 3013

**Character count:** 16982

# Bird diversity of resort Ranu Darungan, Bromo Tengger Semeru National Park, Indonesia

Boni Herdiawan<sup>1</sup>, Reni Ambarwati<sup>12</sup> and Toni Artaka<sup>2</sup>

<sup>12</sup>Department of Biology, Universitas Negeri Surabaya, Kampus Ketintang Unesa, Jalan Ketintang, Surabaya, Indonesia

<sup>2</sup>Bromo Tengger Semeru National Park, Blimbing-Malang, Indonesia

9  
(Received 27 September, 2019; Accepted 10 January, 2020)

## ABSTRACT

6  
Bromo Tengger Semeru National Parks is one of four national parks located in East Java Indonesia, which has unique ecosystem. Resort Ranu Darungan, which is part of this are, has potential bird diversity. This research is aimed to identify species of birds, analyze diversity and abundance of birds in Resort Ranu Darungan Bromo Tengger Semeru National Park. This research used birdwatching approach. The sampling of birds used Index Ponctualle de'Abondance (IPA) method. The identification of birds was conducted based on morphological character, meanwhile diversity and abundance analyzed using diversity and abundance index. The research result showed that there were 67 species of birds in Resort Ranu Darungan, the diversity index was 3.23. Five species have the highest abundance. As conservation area, Resort Ranu Darungan can support birds conservation because many species of birds were found in different habitat.

**Key words:** Resort Ranu Darungan, Bromo Tengger Semeru National Park, *Pernis ptilorhyncus*, *Accipiter soloensis*

## Introduction

1  
Bromo Tengger Semeru National Park is an important site for bird conservation belong to Important Bird Area (International, 2018). There are locations in Bromo Tengger Semeru National Park that have potential bird diversity, they are Resort Coban Trisula, Blok Ireng-Ireng, and Resort Ranu Darungan (Prasetya *et al.*, 2018). Based on previous research, there are 43 species of bird in Resort Coban Trisula with the aggregate spatial distribution and there are 30 species of bird in Blok Ireng-Ireng (Herdiawan *et al.*, 2019; Iqbal *et al.*, 2015). But, there is no publication found about the diversity of birds in Resort Ranu Darungan.

Based on the survey, Resort Ranu Darungan located in elevation 800 masl to 3360 masl. There's some habitat type in Resort Ranu Darungan like the primary forest, secondary forest, and a lake that

supports birds life. Resort Ranu Darungan was the habitat for protected and unprotected birds (Prasetya *et al.*, 2018) such as green peafowl that has observed with a population of less than ten (Balen and Prawiradilaga, 1995).

As a conservation area, diversity data of birds in all habitat types was very important for a reference in determining conservation steps to be taken in future. This research is aimed to identify species of birds, analyze diversity and abundance of birds in Resort Ranu Darungan Bromo Tengger Semeru National Park.

## 10 Materials and Methods

### Study Area

The sampling was located in Resort Ranu Darungan, Bromo Tengger Semeru National Park




Fig. 1. Map of Resort Ranu Darungan for the research. 1. Ranu transect, 2. Loji transect, 3. Tulungagung transect, 4. Nepenthes transect, 5. Supiturang1 transect, 6. Supiturang2 transect.

with coordinate  $8^{\circ}11'26''\text{S} - 112^{\circ}55'37''\text{E}$  until  $8^{\circ}09'38''\text{S} - 112^{\circ}58'23''\text{E}$ . Resort Ranu Darungan has The sampling was conducted in February 2019 until April 2019. Resort Ranu Darungan has altitude between 851 meter a.s.l to 1256 meter a.s.l. There're six transects as an observation path which as representative of habitat in Resort Ranu Darungan, namely primary forest, secondary forest, and a lake.

#### Data Collection and Analysis

This research used a birdwatching approach. The sampling was conducted by using Index Ponctualle de'Abundance (IPA) method (Fachrul, 2012; Syafrudin, 2011). The identification of birds was conducted based on morphological characters and used a guide book for birds identification (MacKinnon *et al.*, 2010). The research used equipments, namely binoculars, cameras, observation papers, a GPS, a thermometer, a hygrometer, a clinometer, and a meter

The sampling was done at 6 am until 11 am (Fachrul, 2012). A transect has one km length which divided into five points observation. Distance between points was 200 meters; observation radius was 25 meters adjust the condition of the observation point. The observation was made for twenty minutes at each point that divided into two times, five minutes early was made for normalizing the condition after movement, after that for fifteen minutes observations were made (Bibby *et al.*, 1992). The research was recorded species, the number of species, behavior, and vegetation samples includes trees species and trees height, and environmental

factors namely temperature, humidity, altitude, light intensity, and vegetation.

Bird species in the Resort Ranu Darungan area identified using a guide book for identification birds (MacKinnon *et al.*, 2010). Diversity birds analyze using Shannon-Winner diversity index (1) (Magurran, 2004), abundance birds analyze using the index of abundance (2) (Syafrudin, 2011). The diversity and abundance of birds analyze strengthened by other indexes namely index of evenness (3) (Magurran, 2004) and index of similarity (4) (Fachrul, 2012).

$$H' = - \sum p_i \ln p_i \quad (1)$$

$$Di = ni/N \times 100\% \quad (2)$$

$$E = H'/\ln S \quad (3)$$

$$SI = C/(a+b+c) \quad (4)$$

#### Results and Discussion

##### Birds of Resort Ranu Darungan, Bromo Tengger Semeru National Park

Based on observation, we found 67 species of birds, divided into 34 families and 14 orders in Resort Ranu Darungan. Passeriformes was order which has most families, the number is 20 families. Meanwhile, the family which has most species was Muscipidae, the number is 14 species. Birds that observed in this research cover one-third species of birds found in all of Bromo Tengger Semeru National Park area (Prasetya *et al.*, 2018).

There were 14 species from 10 families that are protected by law in accordance with the Peraturan Menteri Lingkungan Hidup dan Kehutanan No.106/MENLHK/SETJEN/Kum.1/2018 in Resort Ranu Darungan. Meanwhile in Resort Coban Trisula and Blok Ireng-Ireng Bromo Tengger Semeru National Park, there were 16 species birds that have been protected by law (Herdiawan *et al.*, 2019). Based on that, Resort Ranu Darungan has important value for protected birds.

There was a new species record for Bromo Tengger Semeru National Park. It was *Rhipidura javanica*. *Rhipidura javanica* found in 960 m a.s.l., habitat type was open forest and close to the edge of the national park area. *Rhipidura javanica* was known to be found in range 0 m a.s.l. to 1500 m a.s.l. (MacKinnon *et al.*, 2010). In general, *Rhipidura javanica* lives in low lands (Harmoko and Dewantara, 2018; Rusmendo *et al.*, 2009; Sukara *et al.*, 2014; Sulistyadi, 2017).

**Table 1.** Birds of Resort Ranu Darungan

No	Common Names	Ordo	Family	Scientific Name	Abundance (%)
1	Chinese sparrowhawk	Accipitriformes	Accipitridae	<i>Accipiter soloensis</i>	12.85
2	Black Eagle			<i>Ictinaetus malayensis</i>	1.65
3	Crested serpent eagle			<i>Spilornis cheela</i>	1.15
4	Oriental honey-buzzard			<i>Pernis ptilorhynchus</i>	20.43
5	Banded kingfisher	Coraciiformes	Alcedinidae	<i>Lacedo pulchella</i>	0.16
6	Javan kingfisher			<i>Halcyon cyanoventris</i>	0.66
7	Blue-eared kingfisher			<i>Alcedo meninting</i>	0.66
8	Cave swiftlet	Apodiformes	Apodidae	<i>Collocalia linchi</i>	6.43
9	Pied triller	Passeriformes	Campephagidae	<i>Lalage nigra</i>	0.16
10	Sunda minivet			<i>Pericorotus miniatus</i>	2.64
11	Scarlet minivet			<i>Pericocotus flammeus</i>	3.95
12	Javan leafbird	Passeriformes	Chloropsidae	<i>Chloropsis cochinchinensis</i>	0.33
13	Olive-backed tailorbird	Passeriformes	Cisticolidae	<i>Orthotomus sepium</i>	1.98
14	Plain prinia		Cisticolidae	<i>Prinia inornata</i>	0.33
15	Little Cuckoo-dove	Columbiformes	Columbidae	<i>Macropygia ruficeps</i>	1.65
16	Black-naped fruit-dove			<i>Ptilinopus melanospilus</i>	0.16
17	Grey-capped emerald dove			<i>Chalcophaps indica</i>	0.16
18	Slender-billed crow	Passeriformes	Corvidae	<i>Corvus enca</i>	1.81
19	Greater coucal	Cuculiformes	Cuculidae	<i>Centropus sinensis</i>	0.16
20	Chestnut-breasted malhoka			<i>Phaenicophaeus curvirostris</i>	0.33
21	Red-billed malhoka			<i>Phaenicophaeus javanicus</i>	0.16
22	Blood-breasted flowerpecker	Passeriformes	Dicaeidae	<i>Dicaeum sanguinolentum</i>	0.66
23	Scarlet-headed flowerpecker			<i>Dicaeum trochileum</i>	0.49
24	Crimson-breasted flowerpecker			<i>Prionochilus percussus</i>	0.49
25	Greater racket-tailed drongo	Passeriformes	Dicruridae	<i>Dicrurus paradiseus</i>	0.16
26	Black drongo			<i>Dicrurus macrocercus</i>	0.33
27	Ashy drongo			<i>Dicrurus leucophaeus</i>	3.79
28	White-headed munia	Passeriformes	Estrildidae	<i>Lonchura maja</i>	0.16
29	Black-thighed falconet	Falconiformes	Falconidae	<i>Microhierax fringillarius</i>	1.48
30	Yellow-eared barbet	Piciformes	Megalaimidae	<i>Psilopogon australis</i>	0.82
31	Black-banded barbet			<i>Psilopogon javensis</i>	1.15
32	White-crowned fork-tail	Passeriformes	Muscicapidae	<i>Enicurus leschenaulti</i>	0.16
33	Blue-and-white flycatcher			<i>Cyanoptila cyanomelana</i>	0.33
34	Snowy-browed flycatcher			<i>Ficedula hyperythra</i>	0.16
35	Yellow-rumped flycatcher			<i>Ficedula zanthopygia</i>	0.16
36	Narcissus flycatcher			<i>Ficedula narcissina</i>	0.16
37	Indigo flycatcher			<i>Eumyias indigo</i>	0.82
38	Blue whistling-thrush			<i>Myophonus caeruleus</i>	0.16
39	White-flanked sunbird	Passeriformes	Nectariniidae	<i>Aethopyga eximia</i>	0.16
40	Javan sunbird			<i>Aethopyga mystacalis</i>	1.32
41	Olive-backed sunbird			<i>Cinnyris jugularis</i>	0.49
42	Streaky-breasted spiderhunter			<i>Arachnothera affinis</i>	0.82
43	Golden whistler	Passeriformes	Pachycephalidae	<i>Pachycephala pectoralis</i>	2.64
44	Horsfield's Babbler	Passeriformes	Pellorneidae	<i>Malacocincla sepiaria</i>	5.27
45	Green junglefowl	Galliformes	Phasianidae	<i>Gallus varius</i>	0.33
46	Mountain warbler	Passeriformes	Phylloscopidae	<i>Phylloscopus trivirgatus</i>	0.16
47	Fulvous-breasted woodpecker	Piciformes	Picidae	<i>Dendrocopos macei</i>	0.16
48	Crimson-winged woodpecker			<i>Picus puniceus</i>	0.16
49	Javan banded pitta	Passeriformes	Pittidae	<i>Hydroornis guajanus</i>	0.16
50	Yellow-throated hanging-parrot	Psittaciformes	Psittacidae	<i>Loriculus pusillus</i>	1.81
51	Javan bulbul	Passeriformes	Pycnonotidae	<i>Ixos virens</i>	1.15
52	Black-headed bulbul			<i>Pycnonotus atriceps</i>	0.16



No	Common Names	Ordo	Family	Scientific Name	Abundance (%)
53	Sooty-headed bulbul	14	Rallidae	<i>Pycnonotus aurigaster</i>	2.97
54	Yellow-vented ulbul			<i>Pycnonotus goiavier</i>	5.27
55	White-breasted waterhen			<i>Amaurornis phoenicurus</i>	0.16
56	Sunda Pied Fantail			<i>Rhipidura javanica</i>	0.33
57	White-bellied fantail	Strigiformes	Rhipiduridae	<i>Rhipidura euryura</i>	0.99
58	Brown wood owl			<i>Strix leptogrammica</i>	0.16
59	Asian glossy starling			<i>Aplonis panayensis</i>	0.33
60	Grey-cheeked tit-babbler			<i>Mixornis flavicollis</i>	0.82
61	Chestnut-capped babbler	Passeriformes	Timaliidae	<i>Timalia pileata</i>	0.66
62	Crescent-chested babbler			<i>Cyanoderma melanothorax</i>	0.66
63	Orange-breasted trogon			<i>Harpactes oreskios</i>	0.33
64	Barred buttonquail			<i>Turnix suscitator</i>	0.16
65	Black-winged flycatcher-shrike	Passeriformes	Vangidae	<i>Hemipus hirundinaceus</i>	1.32
66	Pied shrike-babbler			<i>Pteruthius flaviscapris</i>	0.16
67	Javan grey-throated white-eye			<i>Heleia javanica</i>	1.48

### Bird Diversity and Abundance in Resort Ranu Darungan Bromo Tengger Semeru National Park

Based on observation and calculation, Birds of Resort Ranu Darungan has index diversity 3.23 which interpretative as not high diversity. Based on the index diversity value, diversity in every transect was different.

The highest diversity index was Loji transect (2.95). (Fig. 2), Ranu transect observed highest number of species 45 species. Based on diversity index, the diversity index of birds in Resort Ranu Darungan smaller than Gunung Ciremai National Park that has similar habitat (Dewi *et al.*, 2007).

There were some factors that have influence on diversity index, namely number of species and abundance (McNaughton and Wolf, 1992). If there were not balance between number of species and abundance, it can make different index diversity (Magurran, 2004). Ranu transect has highest number of species due to various habitat namely primary forest, secondary forest, and a lake (Krebs, 1989). Supiturang 2 transect has lowest number of species due to habitat condition which not stable after fire two years ago (Krebs, 1989).

Based on the observation, there were species birds that have dominant abundance. They were *Pernis ptilorhyncus* (20.42%), *Accipiter soloensis* (12.85%), *Collocalia linchi* (6.43%), *Turdinus sepiarius* (5.27%), and *Pycnonotus goiavier* (5.27%). Meanwhile, the index of evenness area was 0.5. It can be interpreted that there was unevenness on abundance of birds' species in Resort Ranu Darungan. If explained about the index of evenness in research

transect, Ranu transect was lowest (0.42) and Tulungagung transect was highest (0.88) (Fig 3).



Fig 2. The diversity index of each transect

Ranu transect has the lowest index of evenness due to there were some species that have highest abundance, meanwhile Tulungagung transect have highest index of evenness as all of observed species have similarity abundance. *Pernis ptilorhyncus* and *Accipiter soloensis* were species that have highest abundance in Ranu Transect. *Pernis ptilorhyncus* and *Accipiter soloensis* have highest abundance due to migration in large quantities. Bromo Tengger Semeru National Park is on the migrant raptor trajectory (Prasetya *et al.*, 2018). In addition, research time coincides with reverse flow of migrant birds (Nijman, 2001). The presence of migratory birds in Resort Ranu Darungan strengthened this area was important for birds conservation (Herdiawan *et al.*, 2019).

Checklist of birds observed in each transect has similarity. Based on analyzing, Index of similarity highest number was Nepenthes transect-Supiturang 1 transect (0.37037) meanwhile the lowest number was Ranu transect-Loji transect (0.076923) (Table 2).

**Table 2.** Index of similarity each transect

Transect		Total			SI
		a	b	c	
Ranu	Loji	40	20	5	0.076923**
Ranu	Tulungagungan	38	15	7	0.116667
Ranu	Nepenthes	36	7	9	0.173077
Ranu	Supiturang1	32	8	13	0.245283
Ranu	Supiturang2	34	0	11	0.244444
Loji	Tulungagungan	18	15	7	0.175
Loji	Nepenthes	18	9	7	0.205882
Loji	Supiturang1	20	16	5	0.121951
Loji	Supiturang2	20	6	5	0.16129
Tulungagungan	Nepenthes	16	10	6	0.1875
Tulungagungan	Supiturang1	13	12	9	0.264706
Tulungagungan	Supiturang2	17	6	5	0.178571
Nepenthes	Supiturang1	6	11	10	0.37037*
Nepenthes	Supiturang2	11	6	5	0.227273
Supiturang1	Supiturang2	16	6	5	0.185185

a = the number of species found only at transect a

b = the number of species found only at transect b

c = the number of species found in both transect a – b

\* = highest value

\*\* = lowest value

A comparison will have full similarity if index of similarity value equals one. There were different values of index similarity due to different habitat types in each transect, other than that bird cruising rate and adaptability influence to birds encounter in each transect (Soendjoto *et al.*, 2015; Adelina *et al.*, 2016; Wulandari and Kuntjoro, 2019).

**Fig 3.** Index evenness of each transect

## Conclusion

Based on research diversity of birds in Resort Ranu Darungan Bromo Tengger Semeru National Park, it can be concluded that there were 67 species of birds belong to 34 families and 14 orders in Resort Ranu Darungan Bromo Tengger Semeru National Park, Indonesia. Bird diversity in Resort Ranu Darungan

based on Shanon-Wiener index diversity relatively not high (3.23). Some species of birds have a high abundance (>5%), they were *Pernis ptilorhynchus* (20.42%), *Accipiter soloensis* (12.85%), *Collocalia linchi* (6.43%), *Turdinus sepiarius* (5.27%), and *Pycnonotus goiavier* (5.27%).

## References

- Adelina, M., Harianto, S.P. and Nurcahyani, N. 2016. Keanekaragaman Jenis Burung di Hutan Rakyat Pekon Kelung Kecamatan Kotaagung Kabupaten Tanggamus. *J. Hutan Lestari*. 4 : 51–60. <https://doi.org/10.23960/jsl2451-60>
- Balen, S. Van and Prawiradilaga, D. 1995. The distribution and status of green peafowl *Pavo muticus* in Java. *Biol. Conserv.* 71 : 289–297.
- Bibby, C.J., Burgess, N.D. and Hill, D.A. 1992. Bird Census Techniques, The Wilson Bulletin. Academic Press INC, San Diego. [https://doi.org/10.1676/0043-5643\(2001\)113\[0468:\]2.0.co;2](https://doi.org/10.1676/0043-5643(2001)113[0468:]2.0.co;2)
- Dewi, R.S., Mulyani, Y. and Santosa, Y. 2007. Keanekaragaman Jenis Burung Di Beberapa Tipe Habitat Taman Nasional Gunung Ciremai. *Media Konservasi*. 12 : 114–118.
- Fachrul, M.F. 2012. *Metode Sampling Bioekologi*. Bumi Aksara, Jakarta.
- Harmoko, E.P. and Dewantara, I. 2018. Studi Keanekaragaman Jenis Burung Diurnal Di Seksi Pengelolaan Taman Nasional (SPTN) Wilayah III

- Selimbau Taman Nasional Danau Sentarum Kabupaten Kapuas Hulu Provinsi Kalimantan Barat (Study on Diversity of Diurnal Birds in National Park Management Sectio. *J. Hutan Lestari*. 6: 293-298.
- Herdiawan, B., Mubarak, M.M., Ambarwati, R. and Nugra Prasetya, K. 2019. Status Konservasi Burung Pada Resort Coban Trisula Dan Blok Ireng-Ireng, Taman Nasional Bromo Tengger Semeru Conservation Status of Birds in Coban Trisula Resort and Ireng-Ireng Block, Bromo Tengger Semeru National Park. *SIMBIOSA* 8: 70-84. <https://doi.org/10.33373/sim-bio.v8i1.1871>
- International, B. 2018. Important Bird Areas Factsheet: Gunung Bromo Tengger-Semeru [WWW Document]. URL <http://www.birdlife.org> (accessed 6.15.18).
- Iqbal, K., Gama, Z.P. and Yanuwidi, B. 2015. Spatial Distribution of Birds in Coban Trisula, Bromo Tengger Semeru National Park, in: 6th ICGRC. pp. 26-30.
- Krebs, C.J. 1989. *Ecological Methodology*. Harper & Row Publisher, New York.
- MacKinnon, J., Phillips, K. and van Balen, B. 2010. Seri Panduan Lapangan Burung-Burung di Sumatera, Jawa, Bali, and Kalimantan. *Birdlife International-Pusat Penelitian and Pengembangan Biologi LIPI, Bogor*.
- Magurran, A.E. 2004. *Measuring Biological Diversity*. Blackwell Publishing, Malden.
- McNaughton, S.J. and Wolf, L.L. 1992. *Ekologi Umum*. Gajah Mada University Press, Yogyakarta.
- Nijman, V. 2001. Spatial and temporal variation in migrant raptors on Java, Indonesia. *Emu*. 101 : 259-263. <https://doi.org/10.1071/MU00022>
- Prasetya, K.N., Widodo, R., Rahmadana, M., Winedar, D.W. and Siswoyo, A. 2018. Burung-Burung Taman Nasional Bromo Tengger Semeru. Balai Besar Taman Nasional Bromo Tengger Semeru, Malang.
- Rusmendo, H., Ruskomalasari, A., Prayoga, H.B. and Apriyanti, L. 2009. Keberadaan jenis burung pada lima stasiun pengamatan di sepanjang Daerah Aliran Sungai (DAS) Ciliwung, Depok-Jakarta. *Vis Vitalis*. 2 : 50-64.
- Soendjoto, M.A., Riefani, M.K. and Zen, M. 2015. Penggunaan Tipe Habitat oleh Avifauna di Lingkungan PT Arutmin Indonesia - NPLCT, Kabupaten Kotabaru, Kalimantan Selatan. *J. Sains Mat*. 3 : 19-25.
- Sukara, G., Mulyani, Y. and Muntasib, E.K.S. 2014. Potensi untuk pengembangan wisata birdwatching di pusat konservasi tumbuhan Kebun Raya Bogor. *Bul. Kebun Raya*. 17 : 44-56.
- Sulistiyadi, E. 2017. Kemampuan Kawasan Nir-Konservasi dalam Melindungi Kelestarian Burung Endemik Dataran Rendah Pulau Jawa. Studi Kasus di Kabupaten Kebumen. *J. Biol. Indones*. 6 : 237-253.
- Syafrudin, D. 2011. Keanekaragaman Jenis Burung Pada Beberapa Tipe Habitat Di Tambling Wildlife Nature Conservation (TWNC), Taman Nasional Bukit Barisan Selatan Lampung. IPB (Bogor Agricultural University).
- Widodo, W. 2013. Kajian Fauna Burung Sebagai Indikator Lingkungan Di Hutan Gunung Sawal, Kabupaten Ciamis, Jawa Barat, In: *Proceeding Biology Education Conference: Biology, Science, Environmental, and Learning*. Universitas Negeri Sebelas Maret, Surakarta
- Wulandari, Y.E. and Kuntjoro, S. Keanekaragaman and Kelimpahan Jenis Burung di Kawasan Cagar Alam Besowo Gadungan dan sekitarnya, Kabupaten Kediri Jawa Timur. *Jurnal Riset Biologi and Aplikasinya*. 1 (1) : 18-25.

# Bird diversity of resort Ranu Darungan

## ORIGINALITY REPORT

8%

SIMILARITY INDEX

3%

INTERNET SOURCES

4%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1

[digilib.uinsby.ac.id](http://digilib.uinsby.ac.id)

Internet Source

2%

2

Hablana Rizka Achmadi, Adelia Ramadhani Savitri, Adhe Puspawari Hardhany, Ayu Rini Tirtasari Haryono, Muh. Danny Wirawan. "Illegal Hunting Prevention by Indigenous People in Bromo Tengger Semeru National Park", Jurnal Hukum Novelty, 2019

Publication

1%

3

Susan E. Little. "Myiasis in Wild Birds", Parasitic Diseases of Wild Birds, 12/19/2008

Publication

1%

4

Submitted to Universitas Airlangga

Student Paper

1%

5

Submitted to Universitas Brawijaya

Student Paper

<1%

6

Submitted to International University of Japan

Student Paper

<1%

7

Jerry W. McAtee, D. Lynn Drawe. "Human impact on beach and foredune vegetation of

<1%



# North Padre Island, Texas", Environmental Management, 1980

Publication

8

"Nature-Based Tourism in Asia's Mountainous Protected Areas", Springer Science and Business Media LLC, 2021

Publication

<1 %

9

dl.uctm.edu

Internet Source

<1 %

10

smujo.id

Internet Source

<1 %

11

F Prabandari, F Handoyo. "'ZERO TO HERO" Landscape Utilization of Volcanic Landscapes During and After Eruption", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

12

Reny Rochmawati, Ari Widayanti. "Evaluasi Produktivitas Bongkar Muat Petikemas pada PT. Berlian Jasa Terminal Indonesia (BJTI Port) di Tanjung Perak-Surabaya", Proceedings Series on Physical & Formal Sciences, 2021

Publication

<1 %

13

sig.id

Internet Source

<1 %

14

Lars Schmitz. "Quantitative estimates of visual performance features in fossil birds", Journal of Morphology, 2009

<1 %

15

F S Latumahina, G Mardiatmoko, J  
Sahusilawane. "Bird diversity on small islands  
in Maluku", IOP Conference Series: Earth and  
Environmental Science, 2020

<1 %

Publication

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On