

OF_STRATEGIES_FOR_PHYSIOL OGICAL_RECOVERY_COACH_AN D_PROVINCE.pdf

by

Submission date: 02-Apr-2023 10:32AM (UTC+0700)

Submission ID: 2053186687

File name: OF_STRATEGIES_FOR_PHYSIOLOGICAL_RECOVERY_COACH_AND_PROVINCE.pdf (309.73K)

Word count: 5173

Character count: 28014

THE COMPARISON OF STRATEGIES FOR PHYSIOLOGICAL RECOVERY COACH AND PROVINCE LEVEL FENCING ATHLETE DURING THE PANDEMIC COVID-19

Anisa Nur Fadilah¹, Nor Syafikah², Rini Ismalasari³, Azizati Rochmania⁴, Abdul Hafidz⁵, Kunjung Ashadi⁶

anisa.17060474015@mhs.unesa.ac.id

^{1,2,3,4,5,6}Departement of Sport Coaching Education, Faculty of Sport Science , Universitas Negeri Surabaya, 60213, Indonesia

Abstract. The importance of maintaining physical condition in the midst of a pandemic, especially for sports players must be balanced with the selection of the right *physiological recovery*. Poor choice of recovery can cause physical impairment. This study aims to compare the *physiological recovery* strategies applied by coaches and provincial fencing athletes during the Covid-19 pandemic when carrying out PNN (Regional Training Center *New Normal*) in welcoming PON (National Sports Week) 2021 during the pandemic Covid-19. The method applied in this research is quantitative and presented descriptively. Data were collected through an *online questionnaire* via a google form filled in by 20 samples and a structured oral interview as a complement. Three questions that have met the validity value include *recovery* (0.547), *sleep* (0.526), and *hydration* (0.585) and a reliability value of 0.919. The data analysis technique uses the *mean*, percentage, standard deviation, homogeneity test, and difference test using One Way Anova on normally distributed data. The results showed that the application of *physiological recovery* for coaches and athletes had a value 0,781 ($p > 0.05$). The conclusion shows that there is no significant difference between the application of *physiological recovery* between coach and athlete.

Keyword : *physiological recovery, coach, athlete, fencing, covid-19*

INTRODUCTION

Currently the global pandemic has changed the life order of the world community. Diseases that have a broad impact when compared to SARS and MERS have a high transmission rate. The virus that started in December 2019 in the city of Wuhan and continues to spread throughout the universe to date has been reported from many other countries [1].

A novel infectious disease of the coronavirus family is identified in wuhan, China in late December 2019, later called covid-19 (Chen et al, 2020). On January 30, who declared a global public health emergency. In February, outbreaks began in Iran, Italy, and other countries around the world. Then, the epidemic turned into a pandemic and by the end of March half the world's population was in some form of lockdown. Based on WHO data on April 16, 2020, the total number of COVID-19 cases exceeded 2.1 million cases worldwide, without a total death toll of more than 135,000 [15].

The pandemic managed to spread to parts of the world in just a short time, one of which is Indonesia. Various losses are caused by this virus, especially in the world of sports which postpones sports competitions. The financial crisis experienced by the regions that host one of them is Papua, which is appointed as the organizer of PON 2020. A carefully designed period has to be adapted again so that it gives losses to both the coach and the athlete. The good side is the development of a web-based training program that makes it easier for organizations / clubs to sports players [34].

Maintaining physical fitness is very important in times of a pandemic like this, especially for an athlete. In addition to increasing immunity in the body, physical fitness also keeps the athlete's physical capacity from decreasing, one of which is fencing [25]. Emphasis and dexterity with weapons are characteristic of the sport of fencing. Therefore, this sport requires players to have agility, especially in the hands [33]. The important role of the hands and feet is focused in terms of attacking to defending. There are basic movements in playing fencing, namely the mastery of basic technique movements that are trained first without neglecting other techniques. This is because the majority of the frequency is done in competing and practicing [27].

Various kinds of efforts were made in order to get a good physical quality. Through a systematic and targeted training program, it will have a positive impact on the excellent physical quality of athletes so that it can have a good impact on mental health which can support athlete's technique [5]. Apart from that the physical guidance and aggressive nature of fencing causes considerable muscle damage. In professional fencing, athletes only have a limited period of time to recover from origin after training sessions and competitions [35].

Fatigue that is not immediately followed up through the right procedures can result in damage to cells and tissues in the body, especially during its growth period. Acute injuries also haunt future athletes (early retirement). This has an impact on the decline in athlete's performance to the region or country being defended. Therefore, the province which is the forerunner of the sport in East Java will become an obstacle before reaching the golden peak of achievement [17].

Poor recovery that is done over and over again will cause physical fitness problems. Various efforts are made before the competition so that athletes can perform optimally, including evaluation of techniques, tactics, athlete's nutrition, mastery of strategy and recovery. The basic foundation that must be possessed by coaches and athletes is knowledge of *recovery*. Correct recovery supported by knowledge can certainly support athlete performance [23].

Getting the right recovery after strenuous exercise can minimize injury to reduced performance. Therefore, the role of the coach in choosing a *strategy physiological recovery* must be adjusted to the athlete's motor development. The coach is expected to be more transparent to athletes by adding strategies physiological recovery in the training program along with the types of training that the athletes will apply. The mandatory thing that must be understood and known by coaches and athletes is timing the right so that the type of recovery is right on target [17].

As is well known, body mass index is one of the important factors that can help improve performance. The higher the body mass index can inhibit athlete's performance, especially fencing which requires agility in the game. Therefore, monitoring body weight needs to be considered because the weight gain of 1 kg within 24 hours shows that the recovery is going well. If there is a weight loss of 1 kg, it shows that there is overtraining and the body needs a longer recovery. Conversely, increasing body weight exceeding 1 kg shows a light training load [32].

There is a problem, namely the limitations of the research literature that shows the facts of strategies physiological recovery during the covid-19 pandemic by analyzing the comparison of the application of recovery coaches and fencing athletes in East Java Province during the Covid-19 pandemic as a basis for further discussion. Therefore, it is important to do this research to determine the strategy physiological recovery. The renewal in this study lies in the research subject, namely the Provincial fencing team during the PNN (Regional Training Center for the scheme *New Normal*) during the pandemic covid-19 as a material for further discussion. This research is important to do to determine the comparison of the effectiveness of the implementation of strategies physiological recovery carried out by coaches and fencing athletes of East Java Province during the pandemic covid-19 in order to avoid decreased performance and avoid psychological disorders so that it can become a reference for clubs and the sports community so that they can maintain their strength hold the body well during the pandemic covid-19.

METHODS

The method used in this research is quantitative with a descriptive approach. Target in the research of trainers and fencing athletes in East Java Province. Participation of respondents in this study amounted to 20 people consisting of 7 coaches and 13 athletes. The sample criteria are as follows:

1. Status as an active trainer and athlete of the Puslatda.
2. Following the PNN (Regional Training Center for scheme *New Normal*) at Surabaya State University

The questionnaire used in this study is a questionnaire that has passed the validity test using Microsoft Excel version 365. Consists of 30 questions with indicators including recovery, sleep, hydration with a recovery value (0.547), sleep (0.526), hydration (0.585) and a reliability value of 0.919. Questions about recovery include the use of compressed garments, thermotherapy, cryotherapy, massage, and recovery activated during the Covid-19 pandemic. While questions about sleep include sleep duration, consumption of milk before bedtime, use of light sleepers, conditions after waking up. and questions about hydration including checking hydration, taking supplements, consuming electrolyte and caffeine drinks before exercise, to sleep reminder apps. Data analysis techniques used in this study include maximum, value minimum, average, standard deviation, percentage, data normality test using the Kolmogorov-Sminornov Test. The homogeneity test data for data differences uses One Way Anove. The flow of the research procedure is as follows, distributing the link to questionnaire the research sample via WhatsApp is a data collection method. Control and follow-up during data entry and structured oral interviews were the follow-up until data were collected. Data collection was carried out with a one shot case study in September 2020.

Daily activity during PNN (New Normal Regional Training Center) is a topic in a structured interview. The results in the structured interview show that sleep patterns and eating patterns are more regular because they have been scheduled while in the dormitory, activities carried out outside of training are carried out in the dormitory such as relaxing, playing gadgets or watching dramas, this is due to limited access to in and out during PNN (concentration Area Exercises *New Normal*).

Furthermore, the data were analyzed using a percentage. Data analysis used SPSS version 23. The determination of the criteria for the level of application regarding the application of physiological recovery is presented in the following table:

Table 1. Scale application assessment of physiological recovery Category interval

Intervals	Category
M-0.5 SD X M-1.5 SD	High
M-0.5 SD X <M-0.5 SD	Moderate
M-1.5 SD X <M-0.5 SD	Less

Sriundy, 2015.

Which one:

M = mean count

Elementary school is standard deviation

X = score obtained

RESULTS AND DISCUSSION

After conducting research through online questionnaire regarding the comparison of strategies for physiological recovery coaches and provincial level fencing athletes who are shared with GoogleForm, the following discussion is carried out. The results of the calculation of the characteristics of the research subjects include age, weight, height and body mass index as shown in table 2.

Table 2. Characteristics of research subjects

Subject	Mean and SD		characteristics
	Trainers	Athlete	
Age	40	20 years	
	±10.24	±1.73	

Height	166.46 cm	169.28 cm
	± 6.39	± 8.15
Weight	74.42 kg	62.53 kg
	± 10.29	± 9.99
BMI of	25.75 kg / m ²	22.46 kg / m ²
	± 4.14	± 2.59

Based on the profile data of the research sample shown in table two, the average age of the subjects is 20 to 40 years. The first subject had an average body weight of 74.42 kg with a height of 166.46 cm, while the second subject had an average body weight of 62.53 kg with a height of 169.28 cm. It can be seen that the first subject has an average body mass index of 25.75 kg / which is included in the overweight category, while the second subject has a body mass index of 22.54 kg / which is in the normal category.

Table 3. Description of Respondents Recovery, Sleep, Hydration of

Variable	Subjects	Min	Max	Mean &SD
Recovery	Trainers Athlete	2	6	4.7 \pm 1.48
		1	13	5.7 \pm 3.57
Sleep	Trainers Athlete	3	7	4.8 \pm 1.24
		4	11	7.7 \pm 2.83
Hydration	Coach	1	7	3.8 \pm 2.03
	Athlete	2	13	7.5 \pm 4.6

The data shown in table three shows that the average ratio of the application of recovery, sleep, and hydration in athletes is more dominant than that of coaches, as evidenced by the data that the average athlete who applies recovery is 5.7, while the coach has an average recovery of 4. , 7.`` The facts are reinforced by data on athletes who apply sleep and hydration with an average of 7.7 and 7.5, while athletes have an average of 4.8 and 3.8.

Table 4. Percentage of Recovery Application

Interval	Category	Trainer	Percent	Athlete	Percent
M-0.5 SD X <M-1.5 SD	Very High	1	14%	3	23%
M-0.5 SD X <M-0.5 SD	High	4	57%	9	69%
M -0.5 SD X M-0.5 SD	Moderate	2	29%	1	8%

Total	7	100%	13	100%
--------------	---	------	----	------

Based on Table four, it shows that the implementation of recovery by the majority coach is high with a percentage of 57% while the majority of athletes are high with a percentage of 69%.

Table 5.Percentage of Sleep Application

Interval	Category	Trainer	Percent	Athlete	Percent
M-0.5 SD X <M-1.5 SD	Very High	4	57%	4	31%
M-0.5 SD X <M-0.5 SD	High	2	29%	6	46%
M-0.5 SD X M-0.5 SD	Moderate	1	14%	3	23%
Total		7	100%	13	100%

Based on the data in table five, it shows that the implementation of sleep recovery in the majority of coaches is very high with a percentage of 57% while the majority of athletes are high with a percentage of 46%.

Table 6. Percentage of Application Hydration

Interval	Category	Trainer	Percent	Athlete	Percent
M-0.5 SD X <M-1.5 SD	Very High	1	14%	3	23%
M-0.5 SD X <M-0.5 SD	High	5	72%	8	62%
M -0.5 SD X M-0.5 SD	Moderate	1	14%	2	15%
Total		7	100%	13	100%

Based on the data in table six, it shows that the application of recovery hydration in the majority of coaches is high with a percentage of 72% while the majority of athletes are high with a percentage of 62%.

Table 7. Homogeneity and Normality

Sig	Test of Kolmogorov-Smirnov	
Variables	Recovery, sleep, hydration	
P	0.999	0.724

In table seven, it is known that the Kolmogorov-Smirnov significance value is sig recovery, sleep, and hydration (0.999) where the value is more than 0.05, which means that the normality test on this data is normally distributed. Meanwhile, to determine whether there is an equation between coach and athlete variables. P is known to be 0.724. based on the results of the Sig. 0.742, which value is more than 0.05, which means the data is declared homogeneous.

Table 8. Difference test using One Way Anova

Sig	One Way Anova
Variable	Coach and Athlete
P	0.781

The results of the One Way Anova difference test in table eight show the implementation of strategies recovery (recovery, sleep, hydration) on provincial level fencing trainers and athletes, it is known that the sig value is 0.781. Which ($P > 0.05$) indicates that there is no significant difference in the application of physiological recovery between coaches and fencing athletes during the pandemic covid-19.

RESULTS

After the difference is performed by one way anova and there is no significant difference between the application of physiological recovery coach and athlete. Data triangulation is also done for the equations of the application of physiological recovery. This indicates that the coach's instructions are already run by the athlete optimally so as to provide for the future performance of the athlete.

Based on research conducted on trainers and fencing athletes in East Java Province regarding physiological recovery with good categories then this can benefit sports physiology because it can support physical activity, minimize injury, and improve athlete performance (both physically and psychologically). Therefore, physiological quality affects achievement [16]. Often the training program provided by the coach contains several risks, especially if the athlete's physiology is not good so that it can cause physical injury. Lack of recovery time coupled with intense training results in both physical and psychological exhaustion. Athletes concentration will be disturbed from training to competition if they are stressed so that it has an impact on performance. The anxiety felt by athletes who do not have a history of injury is certainly different from athletes who have a history of injury [10].

Physiological measures can prevent athletes from drifting due to inadequate recovery, excessive stress, and potential illness. Coping with the stress experienced by athletes during the pandemic Covid-19 is one of the strategies recovery in this difficult time [11]. There are various kinds of research on the treatment and its benefits for fatigue, muscle injury, physiological recovery, and performance improvement. According to Joanna Vaile and Shona Holson in the book *Physiological Recovery*, there are several and appropriate recovery strategies, including: recovery, sleep, and hydration. Recovery consists of cryotherapy, massage, compressant garments, thermotherapy.

The application of *cryotherapy* to the whole body (WPC) as a recovery modality for elite athletes can reduce pain in inflammatory conditions. Given the very cold air temperature from -120C to -150C in 2-3 minutes it can cause strong physiological responses in

users [20]. Four research studies found a significant reduction in muscle soreness, in relation to improved muscle function. Five articles report that Cryotherapy (WBC) positively affects muscle function or performance capacity which in other ways adversely affects exercise mentally [29].

Compressent garment has been proposed to prevent performance degradation and improve recovery by speeding up nutrient delivery, improving post-workout edema, muscle soreness (DOMS), and muscle damage. The use of compressentgarment can minimize the symptoms of muscle damage due to exercise (EIMD) which usually occurs as a result of unusual or eccentric exercise [4]. According to Molly Winke et al [37] in their study, the muscle swelling assessed through changes in upper arm circumference was significantly lower intreatment compressed garment (1.7 vs 2.0 cm in compressed sleeves, $p = 0.012$). Interference ROM during Compressent Garments (PCD) treatment was lower and peak pain was 39% lower (27.5 mm in PCD compared to 45.2 mm in the compress arm (CS), $p < 0.05$). using compressent garments further reduces the peak disturbance and recovery time from DOMS (Delayed Onset Muscle Soreness).

There is a substantial treatment reduction in thermotherapy in fatigue. Through thermotherapy helps athletes to get relief from muscle pain and joints with soft tissue, metabolic rate and also blood flow with vasodilation. Due to increased blood flow, oxygen uptake increases and can heal damaged tissue. A temperature of 36 degrees Celsius in 8-10 minutes will give a feeling of calming to the muscles. Thermotherapy can be in the form of hot cloth, hot bottle, infrared, ultraviolet light, shortwave diathermy, heating pad, pussy tub water, sauna bath [31]. Faiz Shah Sharudin (30) in his research added that the data from fifteen studies were examined, and the overall quality. Immersion duration and water temperature showed an increase in recovery with a duration of 15 minutes with a water temperature of 15 $^{\circ}\text{C}$. The results showed good progress in athletes when the body temperature was more than 11 percent with the time of immersion. Thermotherapy (CWI) has a good effect on athletes including increased performance, reduced muscle soreness and fatigue.

Apart from cooling, sauna, compression to active recovery, the recovery strategy that is mostly used is massage. Massage can relieve muscle tension, reduce muscle soreness, and increase flexibility and range of motion (increases blood flow to muscles, improves cleansing of substances such as blood lactate or creatine kinase). Here techniques such as effleurage, petrissage, tapotement, are usually applied with the total duration of 10-30 minutes [26]. Houyong Zhong [38] in his study of 12 athletes received massage sessions for 20 minutes with the comparison group in bed for the same time. The visual analogy scale (VAS) on the normal scale for muscle fatigue, back muscle endurance, and pulse rate as parameters for stress index (SI), HRV indexes, SDNN, RMSSD, Pnn50, LF hf, DANJKA / HF were analyzed. The results were 12 athletes (HF) were significantly higher immediately after the intervention than the LF control group. Therefore massage can facilitate recovery from muscle fatigue after training or competition for athletes.

Fahmi & Ashadi [8] also added in his research that there was a significant difference in the ratio of reduction in lactic acid levels between the use of the method recovery with sport massage and cryotherapy with a p value < 0.05 . Where the average value of reducing lactic acid levels with sports massage was 4.92 nMol / L while cold water was 3.91 nMol / L. There are many strategies for recovery, but sleep is the most significant strategy in avoiding overtraining. During sleep growth hormone and androgens help for muscle repair, muscle building, and bone growth. Recommended sleep of 7-9 hours is sufficient to restore physiology (metabolism and inflammation), psychology (learning ability, motivation and memory). A 30-minute nap between 1:00 p.m. and 1:30 p.m., after only 4 hours of sleep a night leads to increased alertness and mental and physical performance. The drowsiness that is reported to be reduced after a nap and improved short-term memory can help restore physiological mechanisms to avoid significant muscle inflammation as well as maintenance of the immune system [21].

Lack of sleep negatively impacts the physical and psychological functions of the human body. Because sleep disturbances affect the release of hormones, hunger arises and leads to overeating. The balance of diet and body weight will be affected if athletes experience sleep deprivation for long periods of time. The immune system will be compromised and become weak, more diseases will become infected. In addition, poor sleep quality affects motor performance and increases the risk of injury. It negatively affects the circadian rhythm directly, as well as increased fatigue, decreased reaction time, and decreased maximal oxygen uptake ($\text{VO}_{2\text{max}}$). Humans who don't get enough sleep have negative moods such as depression and decreased motivation which causes psychological stress. Previous studies have shown that negative moods directly affect athlete's performance. The emotional state that is regulated by the brain is very sensitive to the quality and quantity of sleep [24].

Studies report that improved sleep quality in athletes is associated with improved performance and overall success. Apart from minimizing injuries, optimizing health and improving performance can be achieved through increased participation in training. However, the majority of athletes fail to get the recommended sleep hours, thus threatening performance and health [19].

Apart from implementing recovery and sleep patterns, it is important to keep the body well hydrated. If the body is dehydrated it can cause thirst, nausea, vomiting, discomfort during exercise and fainting. What is more fatal is the disruption of the metabolic system if it occurs over a long period of time. Therefore the recommendation to meet fluids during physical activity can prevent dehydration. Apart from physical activity, fulfillment of hydration must be done every time, because the human body's construction is 70% fluid. If the body is well hydrated, energy transport, metabolism, and enzymes can run according to their function smoothly [22]. If the fluids in the body are not fulfilled and experience a decrease, it can result in not optimal body condition [28].

AshadiKunjung et al [2] in their research also reported that the average urine scale was 3.1 points, which means that only 31.2% of athletes are dehydrated. However, after exercising, the urine color index showed a scale of 4.1. And 62.5% of athletes are dehydrated. The results showed that there was a significant change in the level of hydration before and after training. It can be concluded that prolonged exercise increases the risk of dehydration. It is important for athletes to meet the needs of body fluids so that there is no malfunction in the body during sports activities.

The amount of fluid that comes out can be influenced by temperature and humidity so it is necessary to calculate replacement fluids to keep the body hydrated. Therefore, it is necessary to know the right drinking pattern in order to reduce the bad effects of dehydration. [9] in their research stated that the drinking pattern that should be applied to a hot environment with an average temperature of 35.6 °C with aerobic activity is the ad libitum drink drinking pattern. Ad libitum drinking pattern is better than pool drinking planned drinks.

The application of physiological recovery, knowledge related to physiological recovery needs to be improved. Knowledge regarding physiological recovery affects athlete's performance because providing a less than optimal recovery program can reduce athlete's performance and can even lead to the risk of injury [6].

As additional information, there is an obstacle in this study, namely the application of a broad-scale questionnaire. Therefore, it is hoped that the limitations of this study can be used as a basis for further research to make it better and use a broad scale in the future.

CONCLUSION

Based on the results of the research data analysis, it can be seen that there is no significant difference in the application of the strategy physiological recovery carried out between coaches and fencing athletes in East Java Province during the Covid-19 pandemic ($P < 0.05$). Based on the above conclusions, coaches and athletes have implemented physiological recovery in the high category.

SUGGESTION

Based on the results of the research data analysis, it can be seen that the application of the strategy physiological recovery carried out by trainers and fencing athletes in East Java Province.

Based on the results of the discussion of coaches and fencing athletes in East Java Province regarding strategies physiological recovery during the pandemic Covid-19, efforts are needed to maintain these achievements, including providing variations to scheduling recovery. In addition, in order to maintain physical condition during a pandemic in order to maintain athlete performance, maintaining psychological health during a pandemic is necessary in order to reduce burnout during a pandemic. In addition to developing a training model, it is necessary to increase knowledge about recovery in order to obtain better results.

ACKNOWLEDGMENT

The aouthor would like to thank the Faculty of Sport Science, Universitas Negeri Surabaya who has helped in providing funds for the publication of this scientific article so that article can be completed smoothly.

REFERENCES

- 1 Ainun Rahma Mardiyah, Nunung. R Nurwati., 2020. Dampak Pandemi Covid-19 Terhadap Angka Pengganguran Di Indonesia. *Academia.edu*.
- 2 Ashadi Kunjung., ND Mirza., Siantoro G 2018. Hydration Status in adolescent runners: pre and post training. *IOP Conference Series: Materials Science and Engineering*.
- 3 Bekti Mulasih 2020. Penerapan AplikasiGoogle Classroom, Google Form, Dan Quizizz Dalam Pembelajaran Di Masa Pandemi Covid-19.*Jurnal Karya Ilmiah Guru*.
- 4 Brown Fredly., Gisanne Conor., Howatson Glyn., Van Ken Someren., Pedlar Charles., Hill Jessica 2017. Compressent Garments and Serveillance and Competition in Sport Protect Athelete and Staff During and After Pandemic. *Journal of Functional and Kinesiology*.
- 5 Dhecvangga Stainlause 2020. Perbandingan Hasil Tes Kondisi Fisik Atlet Anggar Tahun 2018 dan Tahun 2019 KONI Sidoarjo. *Jurnal Prestasi Olahraga*. Vol.3hal:3.
- 6 Dias Ghufon Abdullah., Ashadi Visits 2018. Pemahaman Pelatih Sekolah Sepakbola Se Kota MadiunTentang Physiological Recovery. *Jurnal Prestasi Olahraga*.
- 7 At SeleniaFronso., Costa Sergio., Montensano Cristina Fransisco at Grutolla 2020. The Effect COVID-19 Pandemic on Preceived Stress and Psycohobisoal States in Italy Athlete. *International Journal of Sport and Exercise Psychology*
- 8 Fahmi Hawin., Ashadi Visits 2019. Perbandingan Sport Massage dan Hydrotheraphy Cold Water Terhadap Physiological Recovery. *JurnalOlahraga*
- 9 Febiyanti Grisna., Ashadi Visits 2019. Perbandingan Jenis Dan Pola Minum Terhadap Status Hidrasi Pada Remaja Laki-laki Dan Perempuan. *Jurnal Olahraga*.
- 11 Hagen Joshua., D Jason Ston., Guy W Hornsby 2020. Covid -19 Serveillance and Competition in Sport: Utilizing Sport Science to Protect Athletes and Staff During and After Pandemic. *Journal of Foundational and Kinesiology*.
- 12 Heindri Johan., Beckman Jurgen., Bertolo Mauzuri., Brink Michel K., KallusWolfgang 2020. Multidimential Monitoring of Recovery Status and Implication for Performance. *In International Journal of Sport Physiology and Performance*.
- 13 Herprandika, RP, Yuliawan, D., Rizky M. Y 2019. The Analysis of Physical Conditions of Puslatkot Athletes

- of Kediri City for "Kediri Emas" in Porprov 2019. *Jurnal SPORTIF*.
- 14 Hossein Amir Mehsafar., Gazeni Oarisa., Moghadam Ali Zadeh., Carlos Jose Janes Sanchez 2020. Addressing Potential Impact of Covid-19 Pandemic on Physical and Mental Health of Elite Athlete. *Elsevier Public Health Emergency Collection*.
 - 15 JF Chan., SYuan., KH Kok, et al 2020. A Family Cluster of Pneumonia Associated With The 2019 Novel Corona Virus Indicating Person Transmission: A Study of A Family Cluster. *The Lancet Original Health*.
 - 16 Ketut Iwan Swades., Nyoman I Kanca 2020. Signifikansi Program Latihan Dalam Upaya Meningkatkan Performa Atlet. *Seminar Nasional Riset Inovatif 2020*.
 - 17 Kusumardhana., Buyung 2018. Perbandingan Metode Recovery Aktif dan Metode Konstabilitas Terhadap Asam Laktat. *Jurnal of Sport and Exercise Science*. Vol 1 (2).
 - 18 Latate Michele., E Grace Vincet., Duffield Rn.m D Roach Greory L Shaona Hlason., J Luke Heals., Sargent Charli 2018. Can Sleep Be Used as an Indicator of Excess and Excessive of Athlete Exercise ?. *Frontiers in Physiology*. Vol. 99 p: 436.
 - 19 M Adrew Watson 2017. Sleep and Athletic Performance. *Current Sport Medicine Report*. Vol. 16 p.: 413-418.
 - 20 M Emely Patriage., Cooke Julie., McKune Andrew., B David Pyne 2019. Whole-Body Cryotherapy: Potential to Enhance Athlete Preparation for Competition? *Frontiers in Physiology*.
 - 21 Marshall, JG Geoff MSc., Turner 2016. The Importance of Sleep for Athlete Performance *Strength and Conditioning Journal*. Vol. 38 page: 1.
 - 22 Mita Laily Andriana., Ashadi Kunjung 2019. Olahraga Di Lingkungan Indoor Pada Malam Hari Menghasilkan Rasio Keringat Lebih Banyak Dibandingkan Pagi Hari.
 - 23 Musta'in Mukhammad 2020. Manajemen Cedera Akibat Olahraga Pada Pelajar SMA Sdo Kecamatan Pringsurat. *Jurnal Pengabdian Masyarakat*. Vol. 6 (1).
 - 24 Mon-Lopez, D., de la Rubia Riazia, A., Hontoria Galan, M., & Refoyo Roman, I. 2020. The Impact of Covid-19 and the effect of psychological factors on training conditions of handball and public health. Vol. 17 page: 18.
 - 25 Nur Melly A 2020. Survey Aspek Kebugaran Jasmani Kecepatan, Kelincahan, Dan Daya Ledak Atlet Bola Basket Kabupaten Bangkalan Di Masa Pandemi Covid-19. *Prosiding SENOPATI*. Vol. 1 hal: 17-21.
 - 26 Poppendick Wigeand., Wagmann Melisa., Kellman Michael., Pfelfer Mark., Mater Team 2016. Massage and Recovery: A Meta-Analytical Review.

- 27 PriyaNuruddin Budi Santoso 2018. Analisis Gerakan Teknik Dasar Dalam Olahraga Anggar. *Prosiding Seminar Biomekanika Olahraga*.
- 28 Raditya Aprilia Mahayuni 2018. Profil Hidrasi Atlet Pencak Silat PUSLATCAB Kabupaten Bangkalan Pada Sesi Latihan. *Jurnal Prstasi Olahraga*.
- 29 Rose Catriona., M Kate Edewards., Siegler Jason 2017. Whole-Body Cryotherapy As A Recovery Teechinique After Exrcise: A Review of Literature. *International Journal of Sport Medicine*. Vol. 38 p: 14.
- 30 Shah Faiz Sharudin 2021. The Effect of Cold-Water Immersion for Recovery on Sport Performance: A Systematic Review. *Journal of Sports Medicine*.
- 31 Sharma Sanjeev 2019. Thermotherapy: A Boom in Sport Injuries. *International Journal of Yogic, Human, and Sport Science*. Vol. 4 things: 1.
- 32 Suharjo, 2017. Aktivitas Fisik dan *Recovery*.*Prosiding Seminar Hilirisasi Penelitian.Prosiding Seminar Hilirisasi Penelitian Untuk Kesejahteraan Masyarakat Lembaga Penellitian Universitas Negeri Medan*.
- 33 SupriyokoAgus., MahardikaWisnu 2018. .Kondisi Fisik Atlet Anggar Surakarta. *Jurnal Sportif: Jurna lPenelitian Pembelajaran*. Vol. 4hal:2.
- 34 SusantoNugroho 2020. Pengaruh Virus Covid-19 TerhadapBidang Olahraga Indonesia. *Jurnal Stamina*. Vol. 3hal:3.
- 35 Tavares Fransisco., Brett Tiaki Smith., Drillee Mtathew 2017. Fatigue and Recovery in Rugby: A Review. *Sports Medicine*.
- 36 Widiowati A., Dechline Grafitte 2018. Latihan Shadow Fencing AnggarProvinsi Jambi. *Journal Unja*.
37. Winke Molly. Williamson Sheleby 2018. Comparison of a pneumatic compression device a compression garment during recovery from DOMS. *International Journal of Exercise Science*. Vol.11hal:3.
- 38 Zhong Houyong., Eungpinichong Wichai., Wang Xingze 2018. Effect of Meechanical-Bed Massage on Exercise-Induced Back Fatigue in Athletes. *Journal of Physical Therapy Science*. Vol.30hal:3.

OF_STRATEGIES_FOR_PHYSIOLOGICAL_RECOVERY_COACH_A...

ORIGINALITY REPORT

9%

SIMILARITY INDEX

6%

INTERNET SOURCES

3%

PUBLICATIONS

4%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

1%

★ Submitted to University of St Mark and St John

Student Paper

Exclude quotes On

Exclude bibliography On

Exclude matches < 3 words