

Effect of Royal Jelly on Performance and Inflammatory Response to Muscle Damage: A Systematic Review

by Endang Sri Wahyuni

Submission date: 02-Feb-2023 03:46PM (UTC+0700)

Submission ID: 2004688590

File name: 10._Efek_royal_jelly._Sendy..pdf (357.44K)

Word count: 4627

Character count: 25630

18

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/366953530>

Effect of Royal Jelly on Performance and Inflammatory Response to Muscle Damage: A Systematic Review

Article · January 2023

DOI: 10.33263/BRIAC.135.479

CITATIONS

0

READS

27

5 authors, including:



Nining Kusnanik
Universitas Negeri Surabaya
72 PUBLICATIONS 126 CITATIONS

[SEE PROFILE](#)



Novadri Ayubi
Universitas Negeri Surabaya
26 PUBLICATIONS 35 CITATIONS

[SEE PROFILE](#)



Rizki Mulyawan
Universitas Negeri Yogyakarta
17 PUBLICATIONS 12 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Talent Identification in Sports [View project](#)

Effect of Royal Jelly on Performance and Inflammatory Response to Muscle Damage: A Systematic Review

Sendy Mohamad Anugrah ^{1*}, Nining Widyah Kusnanik ², Endang Sri Wahjuni ²,
Novadri Ayubi ¹, Rizki Mulyawan ³

¹ Doctoral Program of Sport Science, Postgraduate, Universitas Negeri Surabaya, Indonesia;
sendymohamad.21006@mhs.unesa.ac.id (S.M.A.); novadriayubii@yahoo.com (N.A.);

² Faculty of Sport Science, Universitas Negeri Surabaya, Indonesia; niningwidyah@unesa.ac.id (N.W.K.);
angwahjuni@unesa.ac.id (E.S.W.);

³ Faculty of Sport Science, Universitas Negeri Yogyakarta, Indonesia; rizkimulyawan@uny.ac.id (R.M.);

* Correspondence: sendymohamad.21006@mhs.unesa.ac.id (S.M.A.);

Scopus Author ID 57195430173 (N.W.K.)

57537497000 (N.A.)

Received: 21.09.2022; Accepted: 30.10.2022; Published: 3.01.2023

Abstract: Heavy and excessive physical activity can result in injury or muscle damage in ordinary people and athletes, hence diminishing their physical ability during training and competition. It has been established that royal jelly possesses antioxidant and anti-inflammatory effects and enhances athletic performance. This study aims to go through the published research on royal jelly to find evidence that improves performance and reduces inflammation after muscle injury. The method employed is Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA). The databases included Pubmed/MEDLINE, Scopus, Web of Science, and Embase. The ten papers' objectives, topic suitability, sample size, research technique, funding, and commentary were assessed. This research review reveals that royal jelly reduces the subjective sense of muscle pain intensity, reduces muscle damage via a decrease in creatine kinase (CK); improves muscle function; modulates pro-inflammatory cytokines such as TNF- α , IL-6, and IL-8; and has an antioxidant impact. Administration of royal jelly at doses ranging from 300 to 3000 mg/day before, during, and 72 hours after exercise enhances performance by lowering EIMD and regulating physical activity-induced inflammation. Conclusion Royal jelly enhances performance and the inflammatory response to muscle injury. Royal Jelly can be used as a dietary supplement by athletes and other physically active people whose activities cause muscle injury and inflammation.

Keywords: royal jelly; performance; muscle damage; exercise; inflammation.

© 2023 by the authors. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Excessive and strenuous physical activity can lead to muscular injury or damage in a sedentary individual or athlete (EIMD) [1–4]. Muscle injuries, such as inflammation and DOMS (delayed onset muscle soreness), are characterized by muscular rigidity as well as discomfort [2,5,6]. That often develops 24 hours after high-intensity physical exercise, with the peak of DOMS occurring 24-72 hours later and diminishing after 5-7 days [7,8]. Therefore, athletes or individuals who have DOMS cannot follow exercise routines to the fullest extent, affecting performance in both training and competition sessions and negatively affecting the athlete or individual's psychological state, performance, and training quality [6,9,10].

Numerous health, nutrition, and exercise professionals have conducted comprehensive research on strategies to lower the incidence of EIMD and DOMS, as well as taking non-steroidal anti-inflammatory drugs (NSAIDs) [3,11]. Nevertheless, despite the purported benefits of NSAIDs, clinical evidence suggests that NSAID interventions do not necessarily benefit recovery after EIMD or DOMS [12].

Previous research has demonstrated that nutritional interventions, such as green tea extract, curcumin, and cinnamon, enhance injury recuperation and biomarker changes in individuals and athletes [7,8,13,14]. However, the use of royal jelly to recover individuals and athletes following sports-related injuries is still limited. Royal jelly is known to produce the inflammatory markers CRP and anti-inflammatory adiponectin, decreasing biomarkers associated with EIMD and DOMS [15–17]. However, few studies have looked at the impact of royal jelly on working out, and those have found mixed results, despite periodic investigation. As a result, we assessed the available clinical study data on the effects of royal jelly consumption on EIMD.

1

2. Materials and Methods

This study follows the Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA). This study's analytical structure is compatible with a number of bibliometric studies undertaken on different topics. PICO is the approach used to search for articles. PICO is a library or research data-gathering approach that uses diverse library sources to survey study subjects (books, scientific journals). The population analyzed in academic writing utilizes royal jelly to enhance athletic performance and as a marker for muscle injury.

2.1. Methods for searching and selecting.

26

Databases are analyzed, including PubMed/MEDLINE, Scopus, Web of Science, and Embase. Journals were included if they addressed royal jelly, exercise, inflammation, performance, or muscular injury. Notes were compiled, titles and abstracts were screened, and full-text articles were verified using reference management software (Mendeley). A total of 40 papers published between 2010 and 2021 were identified.

29

2.2. Inclusion and exclusion criteria.

Four electronic databases were used in the study's data selection and analysis. Databases, including Pubmed/MEDLINE, Scopus, Web of Science, and Embase, are available. This study's analytic framework is consistent with others utilized to investigate a wide range of issues. Ten papers published between 2010 and 2021 were retrieved. To find all of the articles published about royal jelly between 2010 and 2021, we used a reference manager (Mendeley) to organize our notes, skimmed the titles and abstracts, and then read the full contents of all the papers that piqued our attention. Articles that only appear in abstract form or in publications that are not peer-reviewed are disqualified. Publications that met the inclusion criteria requirements were selected for this systematic review.




Figure 1. Research design.

3. Results and Discussion

The studies from Tables 1 and 2 are some of the findings from the research included in this literature review.

Table 1. Royal Jelly studies on animal impacts

Author	Sample Characteristics	Study Design	Intervention	Results
(Shirzad M, Yousofi M, Zamanzad B, Sedaghat A, Hosseini M, Shahinfard N, 2014) [18]	Sixty female mice were split into six groups of four, each containing eight-week-old mice. Take Group 1 as a baseline against which the rest of the experiment may be measured. The second group is good control. Subjects in Group 3 received RJ (200 mg/kg) once a day. Patients in Group 4 received RJ (200 mg/kg) twice weekly. RJ (300 mg/kg) was administered once daily to Group 5 and once every two days to Group 6.	Experiment	The sole negative control was a daily application of saline to the wounded area. Positive controls were treated with an ointment containing 2.0% nitrofurazone. The third group received 200 mg/kg of royal jelly daily, whereas the fourth group received the same amount. In the fifth group, 300 mg/kg of royal jelly was administered daily, and in the sixth group, 300 mg/kg of royal jelly was utilized daily. The findings of this study demonstrated that using RJ daily had a greater effect on wound healing than using Nitrofurazone or RJ every two days.	The results of this study reveal that daily consumption of RJ has a greater effect on wound healing than using Nitrofurazone and RJ every two days.
(Aslan & Aksoy, 2015) [19]	Five groups of 35 male Sprague-Dawley white rats weighing 300-350 g each were created. Group 1, Control condition Group 2: the EG group. RJ Group is Group 3. Group 4: RJ and EG. Group 5: EG and RJ	Experiment	Group 1 was given a standard meal and drinking water. The second group received normal feed and water containing 1% ethylene glycol. Group 3 was orally gavaged with 100 mg/kg of Royal Jelly. Group 4 was given water containing 1% ethylene	Royal Jelly's antioxidants inhibit ROS generation and assist the antioxidant system. It is believed that Royal Jelly has anti-inflammatory properties through modulating signaling pathways.

Author	Sample Characteristics	Study Design	Intervention	Results
			glycol. In addition, 27 rats were given 100 mg/kg of Royal Jelly daily by oral gavage. Group 5 was given 1% ethylene glycol-containing water for the first two weeks. During the previous two weeks, mice were given 100 mg/kg of Royal.	
(Takahashi et al., 2018) [20]	The mice with the same mean weight were then divided into four groups: the sedentary control group (Con + Sed) (n = 8), the control training group (Con + Tr) (n = 6), the treated sedentary group RJ (RJ + Sed) group (n = 7), and training treated with RJ (RJ + Tr) group (n = 7).	Experiment	Three weeks of RJ (1 mg/g body weight) or water were given to mice. Resistance training was administered to mice in the training group (20 m/min; 60 minutes; 5 times per week).	These data imply that RJ treatment elicits mitochondrial adaptation in the soleus muscle of ICR through resistance training and AMPK activation. mice

Table 2. Royal Jelly studies human consequences.

Author	Sample Characteristics	Study Design	Intervention	Results
(Morita et al., 2012) [21]	A total of 61 healthy volunteers aged 42-83 years were enrolled and randomly divided into a royal jelly group (n = 31) and a control group (n = 30).	A randomized placebo-controlled, double-blind trial.	3000 mg royal jelly (RJ) or placebo in 100 ml fluid/day taken for 6 months.	The consumption of RJ for six months in humans improved erythropoiesis, glucose tolerance, and mental health.
(Büyükepekçi et al., 2018) [22]	30 healthy 20- to 25-year-old undergraduate students were separated into two groups: the control group received corn starch, and the experimental group received honey.	Experiment	The experimental group received 5 grams of royal jelly and 45 grams of honey, for a total of 50 grams, 20 to 30 minutes before breakfast for eight weeks. In the same way, 50 grams of placebo (corn starch) were administered to the control group.	In adolescents who performed maximal strength training, supplementation with royal jelly and honey had no effect on the increase in weight lifted, which was attributed to the weight training but did elicit hormonal alterations.
(Tasdoğan et al., 2020) [23]	20 healthy, sedentary individuals aged 21-23 years, divided into two groups: the experimental group and the placebo group.	Placebo-controlled experimental design	For 15 days, the experimental group got 1000 mg/day of Royal Jelly, whereas the placebo group received 1000 mg/day of water. Aerobic and anaerobic power measures were taken in the laboratory one day before and after a 15-day interval in both groups.	It can be said that a royal jelly supplementation taken daily for 15 days at 1000 mg has a positive effect on the aerobic capacity of sedentary men.
(Saritas et al., 2011) [24]	Forty male swimmers, all between the ages of 18 and 25, were scouted. The swimmers were randomly split up into four groups of ten.	Placebo-controlled experimental design	Subjects in groups 1, 2, and 3 received up to 1 g and 500 mg of royal jelly, respectively, while group 4 received a placebo (corn starch). Except for the placebo group, royal jelly capsules were taken 20 to 30 minutes before breakfast once a day for four weeks.	Taking 500 mg, 1 g, or 2 g of royal jelly per day for 30 days did not improve swimming performance. Furthermore, because of its high amino acid content, BUN and creatinine levels tend to rise.
(NAZMI et al., 2014) [25]	Forty male swimmers, all between the ages of 18 and 25, were scouted. The	Placebo-controlled	The first group received 2 g/day, the second group received 1 g/day, the third	Four weeks of royal jelly supplementation in this study had no effect. It is

Author	Sample Characteristics	Study Design	Intervention	Results
	swimmers were randomly split up into four groups of ten.	experimental design	group received 500 mg/day, and the fourth group received a placebo. A total of 20 kilometers of swimming is completed in a four-week training period of two hours per day, five times each week.	suggested to be effective with higher doses and for a longer period of time.
(Pancar, 2021) [26]	Forty male participants were split into four groups before the trial began: a control group of smokers (n = 10), an inspiratory muscle training (BMI) group of 10 people, a group of RJ practitioners (n = 10), and a group of RJ practitioners with BMI practitioners (n = 10).	11 Randomized experimental study with a control group	The RJ-supplemented group received 1,000 mg/day of RJ in a glass vial between 8.00 and 10.00 am for 4 weeks.	BMI and RJ supplements improved these measures by modifying tobacco users' iron metabolism.
(Meng et al., 2017) [27]	The study's 194 participants were split into three groups: placebo, low-dose pRJ, and high-dose pRJ.	6 randomized, double-blind, placebo-controlled	The placebo group received no Royal Jelly, whereas the low-dose pRJ group received 1.2 g/day and the high-dose pRJ group received 4.8 g/day.	The intervention had no obvious effect on physical appearance. These data imply that pRJ therapy may not increase muscular strength in the elderly but rather slow its progression.

This literature evaluation set out to determine whether or not royal jelly may mitigate the negative effects of exercise-induced inflammation and enhance performance. Experiments on animals have shown that royal jelly, a honeybee product, has various beneficial pharmacological properties, including anti-tumor, anti-inflammatory, antibacterial, and performance-boosting benefits [20,28–30]. In humans, consuming royal jelly at specific doses can improve performance in athletes and active individuals and mental health and can be used to reduce the effects of inflammation after strenuous physical activity [21–23,27,31,32]. The main results showed that supplementation with a dose of 300 mg/bb royal jelly reduced various inflammatory reactions caused by muscle injury in experimental animals. In contrast, a 3000 mg/bb dose increased physical performance in active individuals and athletes. However, verifying the true effectiveness of royal jelly's antioxidant properties is difficult. In order to provide a clearer analysis, the following results have been categorized based on the varying measurement results from the various investigations [18,21]. The type of exercise, the dose of each supplement, and the intervention length can all impact the final result. Participants' age, gender, ethnicity, body composition, training level, training variations, diet, and health state may all have a role in the results.

3.1. Anti-inflammatory effects of royal jelly.

In a study conducted on animals, administration of royal jelly at doses of 200 mg/kg and 300 mg/kg for five consecutive days showed a greater effect on wound healing than nitrofurazone [18]. Another study found that a monthly dosage of 100 mg/kg of royal jelly reduced inflammatory indicators such as CRP and IL-6 [19]. Recent research on asymptomatic obese adults has shown anti-inflammatory benefits for patients. The capsule containing 333 mg/bb per day was taken orally to the patients for eight weeks. The treatment of RJ led to a drop in the inflammatory marker CRP, an increase in the anti-inflammatory adiponectin, and a decrease in the cytokine IL-6 [16]. Meanwhile, royal jelly reduces the adhesion of

Pseudomonas aeruginosa bacteria, which causes respiratory tract infections in older individuals. According to cell culture investigations, it protects epithelial cells against excessive inflammatory reactions to *P. aeruginosa* infection [15].

DNA and tissues can be safeguarded from damage due to royal jelly. These results point to RJ's potential as a natural antioxidant, able to counteract the inflammatory response caused by UVB rays and the resulting oxidative damage. The RJ group showed that the dose of RJ had a direct correlation with its ability to increase NRF2 levels. When a cell needs to defend itself against oxidative stress, it turns to a group of genes coordinated by the transcription factor NRF2. Antioxidant synthesis, mediated by NRF2, increases proportionally to the number of oxidants produced by cells. However, NRF2 expression is downregulated, and antioxidant synthesis is suppressed if cellular oxidant production is excessive and endogenous antioxidants are unable to compensate. Reduced levels of oxidants are one mechanism by which RJ inhibits inflammation; this decreases NF- κ B expression and, in turn, TNF- α production [33].

3.2. Performance effects of royal jelly.

Results from previous research showed that a daily dosage of 3 grams increased erythropoiesis, glucose tolerance, and mental health in participants aged 48-83 years when supplied for 6 months [21]. A further trial with teenage swimmers found that an intervention of 0.5-2 g/day of royal jelly had no apparent effect on their performance. Therefore, it was determined in this study that a combination of a larger intervention dose and a longer intervention period was optimal [24, 10]. A similar effect was seen in a trial of sedentary people who took royal jelly supplements at a dose of 1 g/day once a day for 15 days [23]. These improved outcomes indicate that more stimulation by RJ intervention is necessary to produce a rise in maximum mitochondrial enzymatic activity, which would have a significant impact on performance. Several components of RJ, including amino acids/proteins like leucine and 10-hydroxy-2-decenoic acid, which are unique fatty acids present in royal jelly, have been reported to induce the activation of AMPK (5'-AMP-activated protein kinase), the main mediator of mitochondrial biogenesis, in skeletal muscle, or myotubes. Meanwhile, royal jelly has a number of ingredients with potential antioxidant effects, including 10-hydroxydecanoic acid, free amino acids including proline, cystine, and cysteine, flavonoids, and phenolic compounds. Studies have revealed that oxidative stress is critical in activating AMPK (5'-AMP-activated protein kinase) in skeletal muscle, which can boost performance [20].

In the past, researchers have shown that taking a royal jelly supplement increases lipid peroxidation and prevents its growth. A significant improvement was seen after supplementation by reducing lipid peroxidation. These findings suggest that lipid peroxidation variables may contribute to the rise in aerobic power and can be employed to counteract this trend [23].

The objective of this study was to see how royal jelly affected healthy athletes' performance and how their bodies handled inflammation after muscle damage. Royal jelly reduces inflammation because it blocks the synthesis of inflammatory cytokines, including interleukin-6 (IL-6), interleukin-8 (IL-8), and tumor necrosis factor- α (TNF- α). Researchers have shown that taking royal jelly can help with pain relief, reduce inflammation indicators like C-reactive protein and creatine kinase, and improve exercise performance. From one thousand milligrams (mg) to three thousand (mg) per day is optimal. Therefore, this study demonstrates that royal jelly has many advantages, particularly as an anti-inflammatory substance, especially for the recovery process and its influence on performance. There have

been no reports of adverse reactions to consuming royal jelly; instead, the key is to determine the optimal dosage for yourself or your athlete to get the most beneficial effects.

4. Conclusions

There is no definitive proof that ingesting royal jelly can boost performance and reduce the inflammatory response to muscle damage. Those who put their bodies under extreme stress, such as athletes, may find that taking a Royal Jelly supplement helps reduce muscle soreness and inflammation.

1

Funding

This research did not receive external funding.

Acknowledgments

The authors would like to thank the support from Universitas Negeri Surabaya, Universitas Sultan Ageng Tirtayasa, Kemendikbudristek and LPDP.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Ayubi, N.; Kusnanik, N.W.; Herawati, L.; et al. Effects of Curcumin on Inflammatory Response During Exercise-Induced Muscle Damage. *2023*, *13*, 1–11, <https://biointerfaceresearch.com/wp-content/uploads/2022/03/BRIAC132.146.pdf>.
2. Tanabe, Y.; Fujii, N.; Suzuki, K. Dietary Supplementation for Attenuating Exercise-Induced Muscle Damage and Delayed-Onset Muscle Soreness in Humans. *Nutrients* **2021**, *14*, 1–19, <https://doi.org/10.3390/nu14010070>.
3. Xin, G.; Eshaghi, H. Effect of omega-3 fatty acids supplementation on indirect blood markers of exercise-induced muscle damage: Systematic review and meta-analysis of randomized controlled trials. *Food Sci Nutr* **2021**, *9*, 6429–6442, <https://doi.org/10.1002/fsn3.2598>.
4. Markus, I.; Constantini, K.; Hoffman, J.R.; Bartolomei, S.; Gepner, Y. Exercise-induced muscle damage: mechanism, assessment and nutritional factors to accelerate recovery. *Eur J Appl Physiol* **2021**, *121*, 969–992, <https://doi.org/10.1007/s00421-020-04566-4>.
5. Ayubi, N.; et al. Effect of acute omega 3 supplementation reduces serum tumor necrosis factor-alpha (TNF-α) levels, pain intensity, and maintains muscle strength after high-intensity weight training. *Retos* **2022**, *46*, 677–682, <https://doi.org/10.47197/retos.v46.93720>.
6. Akehurst, H.; Grice, J.E.; Angioi, M.; Morrissey, D.; Migliorini, F.; Maffulli, N. Whole-body vibration decreases delayed onset muscle soreness following eccentric exercise in elite hockey players: a randomised controlled trial. *J Orthop Surg Res* **2021**, *16*, 589, <https://doi.org/10.1186/s13018-021-02760-4>.
7. da, Silva, W.; Machado, Á.S.; Souza, M.A.; Mello-Carpes, P.B.; Carpes, F.P. Effect of green tea extract supplementation on exercise-induced delayed onset muscle soreness and muscular damage. *Physiol Behav* **2018**, *194*, 77–82, <https://doi.org/10.1016/j.physbeh.2018.05.006>.
8. Yoon, W.Y.; Lee, K.; Kim, J. Curcumin supplementation and delayed onset muscle soreness (DOMS): effects, mechanisms, and practical considerations. *Phys Act Nutr* **2020**, *24*, 39–43, <https://doi.org/10.20463/pan.2020.0020>.
9. Harty, S.; Cottet, M.L.; Malloy, J.K.; Kerksick, C.M. Nutritional and Supplementation Strategies to Prevent and Attenuate Exercise-Induced Muscle Damage: a Brief Review. *Sport Med* **2019**, *5*, 1–17, <https://doi.org/10.1186/s40798-018-0176-6>.
10. Zeng, C.; Luo, G.; Xu, S.; Li, Y. The Application of DOMS Mechanism and Prevention in Physical Education and Training. *J Health Eng* **2022**, *2022*, 1–5, <https://doi.org/10.1155/2022/9654919>.
11. Fraga, G.S.; Aidar, F.J.; Matos, D.G.; et al. Effects of ibuprofen intake in muscle damage, body temperature and muscle power in paralympic powerlifting athletes. *Int J Environ Res Public Health* **2020**, *7*, 1–9, <https://doi.org/10.3390/ijerph17145157>.
12. Duman, E.; Ceylan, K.C.; Akpınar, D.; et al. The effects of steroidal and non-steroidal anti-inflammatory

- drugs on tracheal wound healing in an experimental rat model. *Interact Cardiovasc Thorac Surg* **2021**, *30*, 646-651, <https://doi.org/10.1093/ICVTS/IVZ309>.
13. Costache, A.D.; Costache, I.; Miftode, R.; Ștefan, et al. Beyond the Finish Line: The Impact and Dynamics of Biomarkers in Physical Exercise-A Narrative Review. *J Clin Med* **2021**, *10*, 1-18, <https://doi.org/10.3390/jcm10214978>.
14. Kochman, J.; Jakubczyk, K.; Antoniewicz, J.; Mruk, H.; Janda, K. Health Benefits and Chemical Composition of Matcha Green Tea: A Review. *Molecules* **2021**, *26*, 1-11, <https://doi.org/10.3390/molecules26010085>.
15. Susilowati, H.; Murakami, K.; Yumoto, H.; et al. Royal Jelly Inhibits *Pseudomonas aeruginosa* Adherence and Reduces Excessive Inflammatory Responses in Human Epithelial Cells. *Biomed Res Int* **2017**, *2017*, 1-11, <https://doi.org/10.1155/2017/3191752>.
16. Petelin, A.; Kenig, S.; Kopinč, R.; Deželak, M.; Čermelič, Bizjak, M.; Jenko,.; Pražnikar, Z. Effects of royal jelly administration on lipid profile, satiety, inflammation, and antioxidant capacity in asymptomatic overweight adults. *Evidence-based Complement Altern Med* **2019**, *2019*, <https://doi.org/10.1155/2019/4969720>.
17. Collazo, N.; Carpena, M.; Nuñez-Estevez, B.; Otero, P.; Simal-Gandara, J.; Prieto, MA. Health Promoting Properties of Bee Royal Jelly: Food of the Queens. *Nutrients* **2021**, *13*, 1-25, <https://doi.org/10.3390/nu13020543>.
18. Shirzad, M.; Yousofi, M.; Zamanzad, B.; Sedaghat, A.; Hosseini, M.; Shahinfard, N.; et al. Effects of royal jelly on sterile skin cut repair. *J HerbMed Pharmacol* **2014**, *3*, 97-100, <https://core.ac.uk/download/pdf/143844734.pdf>.
19. Aslan, Z.; Aksoy, L. Anti-inflammatory effects of royal jelly on ethylene glycol induced renal inflammation in rats. *Int Braz J Urol* **2015**, *41*, 1008-1013, <https://doi.org/10.1590/S1677-5538.IBJU.2014.0470>.
20. Takahashi, Y.; Hijikata, K.; Seike, K.; et al. Effects of royal jelly administration on endurance training-induced mitochondrial adaptations in skeletal muscle. *Nutrients* **2018**, *10*, 1-12, <https://doi.org/10.3390/nu10111735>.
21. Morita, H.; Ikeda, T.; Kajita K, et al. Effect of royal jelly ingestion for six months on healthy volunteers. *Nutr J* **2012**, *11*, 1-7, <https://doi.org/10.1186/1475-2891-11-77>.
22. Büyükepekçi, S.; Sarıtaş, N.; Soylu, M.; Mistik, S.; Silici, S. Effects of royal jelly and honey mixture on some hormones in young males performing maximal strength workout. *Phys Educ students* **2018**, *22*, 308-315, <https://doi.org/10.15561/20755279.2018.0605>.
23. Tasdoğan, AM.; Vural, M.; Özdal, M.; Pancar, Z. Effect of royal jelly supplementation on aerobic power output and anaerobic power output. *Prog Nutr* **2020**, *22*, 281-287, <https://doi.org/10.23751/pn.v22i1.8747>.
24. Sarıtaş, N.; Yıldız, K.; Büyükepekçi, S.; Coskun, B. Effect of different levels of royal jelly on biochemical parameters of swimmers. *African J Biotechnol* **2011**, *10*, 10718-10723, <https://doi.org/10.5897/ajb11.1862>.
25. NAZMI S.; KADIR, Y.; BETÜL, C.; SERDAR, B.; BEKİR, Ç. Effect of Royal Jelly Ingestion for Four Weeks on Hematological Blood Markers on Swimmers. *Ovidius Univ Ann Ser Phys Educ Sport Mov Heal* **2014**, *14*, 196-202.
26. Pancar, Z. Effect of inspiratory muscle training with royal jelly supplement on iron metabolism in cigarette addicts. *Eurasian J Med* **2021**, *53*, 15-18, <https://doi.org/10.5152/eurasianjmed.2020.20271>.
27. Meng, G.; Wang, H.; Pei, Y.; et al. Effects of protease-treated royal jelly on muscle strength in elderly nursing home residents: A randomized, double-blind, placebo-controlled, dose-response study. *Sci Rep* **2017**, *7*, 6-14, <https://doi.org/10.1038/s41598-017-11415-6>.
28. Albalawi, A.E.; Althobaiti, N.A.; Alrdahe, S.S.; Alhasani, R.H.; Alaryani, F.S.; BinMowyna, M.N. Antitumor Activity of Royal Jelly and Its Cellular Mechanisms against Ehrlich Solid Tumor in Mice. *Biomed Res Int* **2022**, 1-11, <https://doi.org/10.1155/2022/7233997>.
29. Mohamed, H.K.; Mobasher, M.A.; Ebiya, R.A.; et al. Anti-Inflammatory, Anti-Apoptotic, and Antioxidant Roles of Honey, Royal Jelly, and Propolis in Suppressing Nephrotoxicity Induced by Doxorubicin in Male Albino Rats. *Antioxidants* **2022**, *11*, 1-15, <https://doi.org/10.3390/antiox11051029>.
30. Gevorgyan, S.; Schubert, R.; Yeranosyan, M.; et al. Antibacterial activity of royal jelly-mediated green synthesized silver nanoparticles. *AMB Express* **2021**, *11*, 1-8, <https://doi.org/10.1186/s13568-021-01213-9>.
31. Yuksel, S.; Akyol, S. The consumption of propolis and royal jelly in preventing upper respiratory tract infections and as dietary supplementation in children. *J Intercult Ethnopharmacol* **2016**, *5*, 308-311, <https://doi.org/10.5455/jice.20160331064836>.
32. Chen, Y.F.; Wang, K.; Zhang, Y.Z.; Zheng, Y.F.; Hu, F.L. In Vitro Anti-Inflammatory Effects of Three Fatty Acids from Royal Jelly. *Mediators Inflamm* **2016**, *2016*, 1-12, <https://doi.org/10.1155/2016/3583684>.
33. Fatmawati, F.; Erizka, E.; Hidayat, R. Royal jelly (Bee product) decreases inflammatory response in wistar rats induced with ultraviolet radiation. *Open Access Maced J Med Sci* **2019**, *7*, 2723-2727, <https://doi.org/10.3889/oamjms.2019.704>.

Effect of Royal Jelly on Performance and Inflammatory Response to Muscle Damage: A Systematic Review

ORIGINALITY REPORT

25%
SIMILARITY INDEX

22%
INTERNET SOURCES

20%
PUBLICATIONS

%
STUDENT PAPERS

PRIMARY SOURCES

1 www.mdpi.com Internet Source 3%

2 core.ac.uk Internet Source 2%

3 worldwidescience.org Internet Source 2%

4 doi.org Internet Source 1%

5 mdpi-res.com Internet Source 1%

6 www.ncbi.nlm.nih.gov Internet Source 1%

7 www.ajol.info Internet Source 1%

8 repository.unair.ac.id Internet Source 1%

9 nanobioletters.com Internet Source 1%

10	Aarti Yadav, Surender Singh Yadav, Sandeep Singh, Rajesh Dabur. "Natural products: Potential therapeutic agents to prevent skeletal muscle atrophy", European Journal of Pharmacology, 2022 Publication	1 %
11	www.eajm.org Internet Source	1 %
12	www.scielo.br Internet Source	1 %
13	N W Kusnanik, A Hariyanto, Y Herdyanto, A Satia. "Talent identification model for sprinter using discriminant factor", IOP Conference Series: Materials Science and Engineering, 2018 Publication	1 %
14	Yumiko Takahashi, Kamiyu Hijikata, Kohei Seike, Suguru Nakano, Mai Banjo, Yosuke Sato, Kenya Takahashi, Hideo Hatta. "Effects of Royal Jelly Administration on Endurance Training-Induced Mitochondrial Adaptations in Skeletal Muscle", Nutrients, 2018 Publication	1 %
15	fis.dshs-koeln.de Internet Source	1 %
16	mattioli1885journals.com Internet Source	1 %

17	aivpc41.vub.ac.be Internet Source	<1 %
18	gfgc.kar.nic.in Internet Source	<1 %
19	sportedu.org.ua Internet Source	<1 %
20	Zeyneb Aslan, Laçine Aksoy. "Anti-inflammatory effects of royal jelly on ethylene glycol induced renal inflammation in rats", International braz j urol, 2015 Publication	<1 %
21	hdl.handle.net Internet Source	<1 %
22	www2.mdpi.com Internet Source	<1 %
23	www.hindawi.com Internet Source	<1 %
24	cyberleninka.org Internet Source	<1 %
25	efsupit.ro Internet Source	<1 %
26	pubag.nal.usda.gov Internet Source	<1 %
27	Andreea Bălan, Marius Alexandru Moga, Lorena Dima, Sebastian Toma, Andrea Elena	<1 %

Neculau, Costin Vlad Anastasiu. "Royal Jelly—
A Traditional and Natural Remedy for
Postmenopausal Symptoms and Aging-
Related Pathologies", *Molecules*, 2020

Publication

28

docplayer.biz.tr

Internet Source

<1 %

29

vuir.vu.edu.au

Internet Source

<1 %

30

"Poster Özetleri / Poster Abstracts", *Turkish
Journal of Biochemistry*, 2015

Publication

<1 %

31

AKSOY, Laçine and ASLAN, Zeyneb. "Ratlarda
etilen glikolün indüklediği nefropatide arı
sütünün nefroprotektif ve antioksidatif
etkileri", *Ankara Üniversitesi*, 2017.

Publication

<1 %

32

Joung-Kyue Han, Gangmi Kim. "Role of
physical exercise in modulating the insulin-like
growth factor system for improving breast
cancer outcomes: A meta-analysis",
Experimental Gerontology, 2021

Publication

<1 %

33

Mischoulon, D.. "A double-blind dose-finding
pilot study of docosahexaenoic acid (DHA) for
major depressive disorder", *European
Neuropsychopharmacology*, 200809

Publication

<1 %

34	oamjms.eu Internet Source	<1 %
35	www.science.gov Internet Source	<1 %
36	Diego Fernández-Lázaro, Juan Mielgo-Ayuso, Jesús Seco Calvo, Alfredo Córdova Martínez et al. "Modulation of Exercise-Induced Muscle Damage, Inflammation, and Oxidative Markers by Curcumin Supplementation in a Physically Active Population: A Systematic Review", <i>Nutrients</i> , 2020 Publication	<1 %
37	Hiroyuki Morita, Takahide Ikeda, Kazuo Kajita, Kei Fujioka, Ichiro Mori, Hideyuki Okada, Yoshihiro Uno, Tatsuo Ishizuka. "Effect of royal jelly ingestion for six months on healthy volunteers", <i>Nutrition Journal</i> , 2012 Publication	<1 %
38	Kashif Maroof, Siew Hua Gan. "Bee products and diabetes mellitus", Elsevier BV, 2022 Publication	<1 %
39	Maryam Lohrasbi, Farzaneh Taghian, Khosro Jalali Dehkordi, Seyed Ali Hosseini. "The functional mechanisms of synchronizing royal jelly consumption and physical activity on rat with multiple sclerosis-like behaviors hallmarks based on bioinformatics analysis,	<1 %

and experimental survey", BMC Neuroscience, 2022

Publication

40

S. Büyükipekçi, N. Sarıtaş, M. Soylu, S. Mıstık, S. Silici. "Effects of royal jelly and honey mixture on some hormones in young males performing maximal strength workout", Physical education of students, 2018

Publication

<1 %

41

"Bee Products - Chemical and Biological Properties", Springer Science and Business Media LLC, 2017

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On